

2006 Update



Zimbabwe

EPIDEMIOLOGICAL FACT SHEETS ON HIV/AIDS AND SEXUALLY TRANSMITTED INFECTIONS

December 2006



HIV/AIDS estimates

The estimates and data provided in the following tables relate to 2005 unless stated otherwise. These estimates have been produced and compiled by UNAIDS/WHO. They have been shared with national AIDS programmes for review and comments, but are not necessarily the official estimates used by national governments. In order to calculate regional totals, older data or regional models were used to produce minimum estimates for these countries. The estimates are given in rounded numbers. However, unrounded numbers were used in the calculation of rates and regional totals, so there may be minor discrepancies between the regional/global totals and the sum of the country figures. The new estimates in this report are presented together with ranges, called 'plausibility bounds'. These bounds reflect the certainty associated with each of the estimates. The wider the bounds, the greater the uncertainty surrounding an estimate. The extent of uncertainty depends mainly on the type of epidemic, and the quality, coverage and consistency of a country's surveillance system. The general methodology and tools used to produce the country-specific estimates in the table have been described in a series of papers in *Sexually Transmitted Infections* 2006, 82 (Suppl x). The estimates produced by UNAIDS/WHO are based on methods and on parameters that are informed by advice given by the UNAIDS Reference Group on HIV/AIDS Estimates, Modelling and Projections.

Estimated number of adults and children living with HIV/AIDS, end of 2003 and 2005

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS.

| | 2003 | 2005 |
|---------------------------|-----------|-----------|
| Adults (15+) and children | 1 700 000 | 1 700 000 |
| Low estimate | 1 200 000 | 1 100 000 |
| High estimate | 2 300 000 | 2 200 000 |
| Adults (15+) | 1 600 000 | 1 500 000 |
| Low estimate | 1 100 000 | 1 000 000 |
| High estimate | 2 100 000 | 2 000 000 |
| Children (0-14) | 160 000 | 160 000 |
| Low estimate | 51 000 | 54 000 |
| High estimate | 340 000 | 340 000 |
| Adult rate (15-49) (%) | 22.1 | 20.1 |
| Low estimate | 14.6 | 13.3 |
| High estimate | 30.4 | 27.6 |
| Women (15+) | 930 000 | 890 000 |
| Low estimate | 540 000 | 520 000 |
| High estimate | 1 300 000 | 1 300 000 |

Source: 2006 Report on the global AIDS epidemic

| Estimates 2005 | Men | Women |
|----------------------------------|-----|-------|
| Prevalence among 15-24 year olds | 4.4 | 14.7 |
| Low estimate | 2.3 | 7.7 |
| High estimate | 6.9 | 23.2 |

Source: 2006 Report on the global AIDS epidemic

HIV prevalence among young people

| | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|---------------------------------------|------|------|------|------|------|------|
| Prevalence among 15-24 year olds | | | | | | |
| Prevalence among 15-24 pregnant women | | | | | | |

Source: 2006 Report on the global AIDS epidemic

Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS:

| | 2003 | 2005 |
|---------------------|---------|---------|
| Adults and children | 200 000 | 180 000 |
| Low estimate | 130 000 | 120 000 |
| High estimate | 270 000 | 250 000 |

Source: 2006 Report on the global AIDS epidemic

Estimated number of orphans due to AIDS

Nb: only for generalized epidemics

Estimated number of children who have lost their mother or father or both parents to AIDS and who were alive and under age 17 at the end of 2005:

| Estimated number of orphans | 2003 | 2005 |
|-----------------------------|-----------|-----------|
| Current living orphans | 1 000 000 | 1 100 000 |
| Low estimate | 720 000 | 780 000 |
| High estimate | 1 200 000 | 1 300 000 |

Source: 2006 Report on the global AIDS epidemic

| | 2003 | 2005 |
|------------------|-----------|-----------|
| Maternal orphans | 900 000 | 1 000 000 |
| Low estimate | 650 000 | 750 000 |
| High estimate | 1 100 000 | 1 200 000 |
| Paternal orphans | 620 000 | 680 000 |
| Low estimate | 450 000 | 500 000 |
| High estimate | 750 000 | 820 000 |
| Dual orphans | 570 000 | 680 000 |
| Low estimate | 410 000 | 510 000 |
| High estimate | 700 000 | 820 000 |

Source: 2006 Report on the global AIDS epidemic

| | Year | Total |
|--------------------------|------|-------|
| Education ratio | | |
| External support for OVC | | |

Source:

The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, is the coordination and implementation mechanism for UNAIDS and WHO to compile and improve the quality of data needed for informed decision-making and planning at national, regional and global levels. The primary objective of the working group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the working group collaborates closely with WHO Regional Offices, national AIDS programmes and a number of national and international institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional, and global levels. The Epidemiological Fact Sheets are one of the products of this close collaboration across the globe.

Within this framework, the Fact Sheets collate the most recent country specific data on HIV/AIDS prevalence and incidence, together with information on behaviour (e.g.; casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed upon indicators was not available for many countries in 2005. However these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The fact Sheets may also be instrumental in identifying potential partners when planning and implementing surveillance systems.

The Fact Sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the Working Group would like to encourage all programme managers as well as national and international experts to communicate additional information to them whenever such information becomes available. The Working Group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

Basic indicators

For consistency reasons the data in the table below are taken from official UN publications.

| DEMOGRAPHIC DATA | YEAR | ESTIMATE | SOURCE |
|---|-----------|----------|-------------------------------|
| Total population (thousands) | 2005 | 13 009 | UN Population Division |
| Population aged 15-49 (thousands) | 2005 | 6490 | UN Population Division |
| Female population aged 15-24 (thousands) | 2005 | 1651 | UN Population Division |
| Annual population growth rate (%) | 1995-2004 | 0.9 | UN Population Division |
| % of population in urban areas | 2005 | 35.9 | UN Population Division |
| Crude birth rate (births per 1000 pop.) | 2005 | 29.5 | UN Population Division |
| Crude death rate (deaths per 1000 pop.) | 2005 | 23.4 | UN Population Division |
| Maternal mortality rate (per 100 000 live births) | 2000 | 1100 | World Health Report 2006, WHO |
| Life expectancy at birth (years) | 2004 | 36 | World Health Report 2006, WHO |
| Total fertility rate (per woman) | 2004 | 3.4 | World Health Report 2006, WHO |
| Infant mortality rate (per 1000 live births) | 2004 | 78 | UNICEF / WHO |
| Under 5 mortality rate (per 1000 live births) | 2004 | 129 | World Health Report 2006, WHO |

| SOCIO-ECONOMIC DATA | YEAR | ESTIMATE | SOURCE |
|--|-----------|----------|------------------------------------|
| Gross national income, ppp, per capita (Int.\$) | 2004 | 2180 | World Bank |
| Per capita total expenditure on health (Int.\$) | 2003 | 132 | WHO |
| UN Human Development Index (ranking) | 2005 | 145 | UNDP Human Development Report 2005 |
| General government expenditure on health as % of total expenditure on health | 2003 | 35.9 | WHO |
| Adult literacy rate (%) | 2000-2004 | N/A | UNESCO |
| Male literacy rate (%) | 2000-2004 | N/A | UNESCO |
| Female literacy rate (%) | 2000-2004 | N/A | UNESCO |
| Net primary school enrolment ratio, male (%) | 1998-2004 | 79 | World Bank |
| Net primary school enrolment ratio, female (%) | 1998-2004 | 80 | World Bank |
| Human Poverty Index (ranking) | 2005 | 89 | UNDP Human Development Report 2005 |

| | 2001 | 2002 | 2003 | 2004 | 2005 |
|--|------|------|-----------|-----------|------------|
| National funds spent by governments on HIV/AIDS from domestic sources (US\$) | | | 6 300 000 | 9 900 000 | 12 052 578 |

Source: UNGASS CR and SIDALAC

Contact address

UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

20, Avenue Appia

CH - 1211 Geneva 27

Switzerland

Fax: +41-22-791-4834

email: hivstrategicinfo@who.int

estimates@unaids.org

website: <http://www.who.int/hiv>

<http://www.unaids.org>

Extracts of the information contained in these fact sheets may be reviewed, reproduced or translated for research or private study but not for sale or for use in conjunction with commercial purposes. Any use of information in these fact sheets should be accompanied by the following acknowledgment "UNAIDS/WHO Epidemiological Fact Sheets on HIV/AIDS and Sexually Transmitted Infections, 2006 Update".

HIV prevalence in different populations

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV database maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences are compiled. To provide a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study from which the medians were calculated are printed at the end of this fact sheet. The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and - where applicable - other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.



HIV sentinel surveillance prevalence

| Group | Area | | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | |
|-------------------------|---------------------------|---------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Pregnant women | Major urban areas | N-Sites | 4 | 1 | 3 | 1 | 1 | 2 | 1 | 6 | | 1 | 3 | 3 | 3 | | 3 | | |
| | | Minimum | 16 | 17.1 | 26 | 25.8 | 41.7 | 30 | 30.4 | 22.7 | | 31.1 | 30 | 27.9 | 20.3 | | 18.1 | | |
| | | Median | 18.6 | 17.1 | 28 | 25.8 | 41.7 | 31 | 30.4 | 30 | | 31.1 | 31.1 | 30.6 | 24.8 | | 19.7 | | |
| | | Maximum | 23.8 | 17.1 | 34 | 25.8 | 41.7 | 32 | 30.4 | 34 | | 31.1 | 33.5 | 30.7 | 28.1 | | 20.3 | | |
| | Outside major urban areas | N-Sites | 4 | 15 | 18 | 16 | 24 | 12 | 4 | 30 | 3 | | 18 | 19 | 16 | | 16 | | |
| | | Minimum | 7.6 | 7.7 | 6.6 | 13.7 | 14 | 19 | 34 | 7 | 29 | | 13 | 8.3 | 18.8 | | 10.3 | | |
| | | Median | 16 | 20 | 19.9 | 20.6 | 24.8 | 36.5 | 41.6 | 28.6 | 31 | | 33.1 | 28.5 | 24.9 | | 22.4 | | |
| | | Maximum | 31.6 | 33.8 | 42.1 | 27 | 54 | 70.2 | 59 | 53.3 | 37 | | 70.7 | 45.3 | 36.3 | | 35.8 | | |
| Sex workers | Major urban areas | N-Sites | | | | | | 1 | | | | | | | | | | | |
| | | Minimum | | | | | | 86 | | | | | | | | | | | |
| | | Median | | | | | | 86 | | | | | | | | | | | |
| | | Maximum | | | | | | 86 | | | | | | | | | | | |
| | Outside major urban areas | N-Sites | | | | | | | | | | | | 1 | | | | | 1 |
| | | Minimum | | | | | | | | | | | | 50 | | | | | 57.2 |
| | | Median | | | | | | | | | | | | 50 | | | | | 57.2 |
| | | Maximum | | | | | | | | | | | | 50 | | | | | 57.2 |
| Injecting drug users | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | |
| STI patients | Major urban areas | N-Sites | 1 | 2 | 3 | 1 | | 2 | | | | | | | | | | | |
| | | Minimum | 52 | 39 | 52 | 60.3 | | 50.7 | | | | | | | | | | | |
| | | Median | 52 | 45 | 56 | 60.3 | | 71 | | | | | | | | | | | |
| | | Maximum | 52 | 51 | 70 | 60.3 | | 71.2 | | | | | | | | | | | |
| | Outside major urban areas | N-Sites | 3 | 14 | 13 | 9 | 8 | 7 | 1 | | | | | | | | | | |
| | | Minimum | 24.3 | 24.5 | 39 | 25.6 | 48 | 43 | 71.8 | | | | | | | | | | |
| | | Median | 28.7 | 45.6 | 44.6 | 52 | 54 | 65 | 71.8 | | | | | | | | | | |
| | | Maximum | 59.7 | 65.3 | 59.5 | 71.8 | 75.4 | 87.7 | 71.8 | | | | | | | | | | |
| Men having sex with men | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | |
| Tuberculosis patients | Major urban areas | N-Sites | | 1 | | 1 | | | | 1 | | 1 | | | | | | | |
| | | Minimum | | 56.5 | | 59.7 | | | | 74 | | 73.6 | | | | | | | |
| | | Median | | 56.5 | | 59.7 | | | | 74 | | 73.6 | | | | | | | |
| | | Maximum | | 56.5 | | 59.7 | | | | 74 | | 73.6 | | | | | | | |

| Group | Area | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|-----------------------|---------------------------|---------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Tuberculosis patients | Outside major urban areas | N-Sites | | | 1 | | | 1 | | | | | | | | | |
| | | Minimum | | | 64.8 | | | 84.3 | | | | | | | | | |
| | | Median | | | 64.8 | | | 84.3 | | | | | | | | | |
| | | Maximum | | | 64.8 | | | 84.3 | | | | | | | | | |

Maps & charts

Mapping the geographical distribution of HIV prevalence among different population groups may assist in interpreting both the national coverage of the HIV surveillance system as well in explaining differences in levels of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the WHO Public Health Mapping and GIS Team, Communicable Diseases, is producing maps showing the location and HIV prevalence in relation to population density, major urban areas and communication routes. For generalized epidemics, these maps show the location of prevalence of antenatal surveillance sites. Trends in antenatal sentinel surveillance for higher prevalence countries, or in prevalence among selected populations for countries with concentrated epidemics, are a new addition. These are presented for those countries where sufficient data exist.



The boundaries and names shown and the designations used on the map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. WHO 2006, all rights reserved.

Reported HIV/AIDS cases

Reported AIDS cases

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases are aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections. Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of Anti-Retroviral Therapy (ART).

| | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | Total |
|---------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-------|
| Males | | | | | | | | | | | | | | | | | |
| Females | | | | | | | | | | | | | | | | | |
| Total | | | | | | | | | | | | | | | | | |

Reported HIV cases

A case of HIV infection is defined as an individual with HIV infection irrespective of clinical stage (including severe or stage 4 clinical disease) confirmed by laboratory criteria according to country definitions and requirements.

| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | Total |
|---------|------|------|------|------|------|------|------|------|------|------|-------|
| Males | | | | | | | | | | | |
| Females | | | | | | | | | | | |
| Total | | | | | | | | | | | |

Source:

Note: In some instances, the number in the total column is not the sum of the individual years due to differing reporting, estimation processes or available data.

Sexually transmitted infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STIs are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STIs facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Thus, detection and treatment of individuals with STIs is an important part of an HIV control strategy. In summary, if the incidence/prevalence of STIs is high in a country, then there is the possibility of high rates of sexual transmission of HIV. Monitoring trends in STIs provides valuable insight into the likelihood of the importance of sexual transmission of HIV within a country, and is part of second generation surveillance. These trends also assist in assessing the impact of behavioural interventions, such as delaying sexual debut, reducing the number of sex partners and promoting condom use. Clinical services offering STI care are an important access point for people at high risk for both STIs and HIV. Identifying people with STIs allows for not only the benefit of treating the STI, but for prevention education, HIV testing, identifying HIV-infected persons in need of care, and partner notification for STIs or HIV infection. Consequently, monitoring different components of STI prevention and control can also provide information on HIV prevention and control activities within a country.

STI syndromatic reporting

Genital discharge

| Reported cases | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|
| Males | | | | | | | | | | | |

Source:

Genital ulcers

| Reported cases | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|
| Males | | | | | | | | | | | |
| Females | | | | | | | | | | | |
| Total | | | | | | | | | | | |

Source:

STI etiological reporting

Chlamydia

| Reported cases | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|
| Males | | | | | | | | | | | |
| Females | | | | | | | | | | | |
| Total | | | | | | | | | | | |

Source:

Gonorrhoea

| Reported cases | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|
| Males | | | | | | | | | | | |
| Females | | | | | | | | | | | |
| Total | | | | | | | | | | | |

Source:

Syphilis

| Reported cases | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|
| Males | | | | | | | | | | | |
| Females | | | | | | | | | | | |
| Total | | | | | | | | | | | |

Source:

Herpes simplex

| Reported cases | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|----------------|------|------|------|------|------|------|------|------|------|------|------|
| Males | | | | | | | | | | | |
| Females | | | | | | | | | | | |
| Total | | | | | | | | | | | |

Source:

Syphilis prevalence, women

Percent of blood samples taken from pregnant women aged 15-49 that test positive for syphilis - positive reaginic and treponema test-during routine screening at selected antenatal clinics.

Syphilis prevalence, ANC women

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|-------|------|------|------|------|------|------|------|------|------|------|------|
| Total | | | | | | | | | | | |

Prevalence of curable STIs among specific populations

Prevalence of curable STIs among female sex workers

| | Year | Area | Rate | Range |
|----------------|------|------|------|-------|
| Chlamydia | | | | |
| <i>Source:</i> | | | | |

| | Year | Area | Rate | Range |
|----------------|------|------|------|-------|
| Gonorrhoea | | | | |
| <i>Source:</i> | | | | |

| | Year | Area | Rate | Range |
|----------------|------|------|------|-------|
| Syphilis | | | | |
| <i>Source:</i> | | | | |

| | Year | Area | Rate | Range |
|----------------|------|------|------|-------|
| Trichomoniasis | | | | |
| <i>Source:</i> | | | | |

Prevalence of curable STIs among other specific populations

Specific populations according to the epidemic pattern of the country

| | Year | Area | Rate | Range |
|----------------|------|------|------|-------|
| Chlamydia | | | | |
| <i>Source:</i> | | | | |

| | Year | Area | Rate | Range |
|----------------|------|------|------|-------|
| Gonorrhoea | | | | |
| <i>Source:</i> | | | | |

| | Year | Area | Rate | Range |
|----------------|------|------|------|-------|
| Syphilis | | | | |
| <i>Source:</i> | | | | |

| | Year | Area | Rate | Range |
|----------------|------|------|------|-------|
| Trichomoniasis | | | | |
| <i>Source:</i> | | | | |

Health service and care indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS - related issues.

| Access to health care | | | |
|--|------|----------|------------|
| Indicators | Year | Estimate | Source |
| % of population with access to health services - total | | | |
| % of population with access to health services - urban | | | |
| % of population with access to health services - rural | | | |
| Contraceptive prevalence rate (%) | 1999 | 53.5 | UNPOP |
| Percentage of contraceptive users using condoms | | | |
| % of births attended by skilled health personnel | 1999 | 73 | UNICEF |
| % of 1-yr-old children fully immunized - DPT | 2004 | 85 | WHO/UNICEF |
| % of 1-yr-old children fully immunized - Measles | 2004 | 80 | WHO/UNICEF |
| % of ANC clinics where HIV testing is available | | | |

Estimated number of adults (15+) in need of treatment

Total number of adults needing antiretroviral therapy

| | 2003 | 2005 |
|---------------|---------|---------|
| Both sexes | 340 000 | 320 000 |
| Low estimate | 220 000 | 210 000 |
| High estimate | 450 000 | 430 000 |

Source: WHO and UNAIDS, March 2006

Estimated number of people receiving antiretroviral therapy

Total number of people receiving antiretroviral therapy at end of each year

| | 2003 | 2005 |
|------------|------|--------|
| Males | | |
| Females | | |
| Both sexes | 3500 | 25 000 |

Source: Based on the most recent calculated ART need estimates by WHO and UNAIDS, as of March 2006.

| Coverage | 2003 | 2005 |
|------------|------|------|
| Both sexes | 1% | 7% |

Source: WHO and UNAIDS, March 2006

Comments: See also the paediatrics estimates section on the next page, as the ART need among children should also be taken into account for estimating ART coverage.

Services providing antiretroviral therapy

Reported number of sites that are providing antiretroviral therapy

| | 2003 | 2005 |
|---------|------|------|
| Public | | |
| Private | | |
| Total | | 48 |

Source: (total 2005) Annex 3: Progress on Global Access to HIV Antiretroviral Therapy, A Report on "3 by 5" and Beyond. Geneva, WHO and UNAIDS, March 2006.

Comments:

Paediatrics estimates, 2005

| | Total | Source |
|-----------------------------------|---------|-----------------------------------|
| Children living with HIV | | |
| Low estimate | | |
| High estimate | | |
| Children in need of ART | 44 000 | |
| Low estimate | 18 000 | |
| High estimate | 83 000 | <i>WHO and UNAIDS, March 2006</i> |
| Children receiving ART | | |
| Children in need of cotrimoxazole | 250 000 | |
| Low estimate | 100 000 | |
| High estimate | 470 000 | <i>WHO and UNAIDS, March 2006</i> |
| Children receiving cotrimoxazole | | |

Comments:

Coverage of HIV testing and counselling

Number of public, private and NGO sites providing testing and counselling services.

| | Year | Area | Total number of sites |
|----------------|------|------|-----------------------|
| Public sector | | | |
| Private sector | | | |
| NGOs | | | |
| Total | | | |

Source:

Number of people counselled and tested over time

Number of people who have been tested and counselled in the country.

| | 2003 | 2004 | 2005 |
|------------|------|------|------|
| Males | | | |
| Females | | | |
| Both sexes | | | |

Source:

Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, injecting drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of a standard set of indicators defined in the National Guide (Source: National AIDS Programmes, A Guide to Monitoring and Evaluation, UNAIDS/00.17) and regular behavioural surveys in order to monitor trends in behaviours and to target interventions. The indicators on knowledge and misconceptions are an important prerequisite for prevention programmes to focus on increasing people's knowledge about sexual transmission, and, to overcome the misconceptions that act as a disincentive to behaviour change. Indicators on sexual behaviour and the promotion of safer sexual behaviour are at the core of AIDS programmes, particularly with young people who are not yet sexually active or are embarking on their sexual lives, and who are more amenable to behavioural change than adults. Finally, higher risk male-male sex reports on unprotected anal intercourse, the highest risk behaviour for HIV among men who have sex with men.

Knowledge of HIV prevention methods

Prevention indicator: Percentage of young people 15-24 who both correctly identify two ways of preventing the sexual transmission of HIV and who reject three misconceptions about HIV transmission.

| | Total | Urban | Rural | Year |
|---------|-------|-------|-------|------|
| Males | | | | |
| Females | | | | |

Source:

Reported condom use at last higher risk sex (young people 15-24)

Prevention indicator: Proportion of young people reporting the use of a condom during sex with a non-regular partner.

| | Total | Urban | Rural | Year |
|---------|-------|-------|-------|------|
| Males | | | | |
| Females | | | | |

Source:

Age-mixing in sexual partnerships among young women

The proportion of young women who have sex in the last 12 months with a partner who is 10 or more years older than themselves.

| | Total | Urban | Rural | Year |
|---------|-------|-------|-------|------|
| Females | | | | |

Source:

Reported non regular sexual partnerships

Prevention indicator: Proportion of young people 15-24 having at least one sex partner other than a regular partner in the last 12 months.

| Year | Males | Females |
|------|-------|---------|
| | | |

Source:

Ever used a condom

Percentage of people who ever used a condom.

| | Age | Total | Urban | Rural | Year |
|---------|-----|-------|-------|-------|------|
| Males | | | | | |
| Females | | | | | |

Source:

Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child.

| | Year | Percentage |
|--|------|------------|
| | | |

Source:

Age at first sexual experience

Percentage of 15-19 year olds who have had sex before age 15.

| | Year | Males | Females |
|--|------|-------|---------|
| | | | |

Source:

Prevention indicators

Prevention of mother-to-child transmission (PMTCT) nationwide

Infection of HIV from an HIV-positive mother to her child during pregnancy, labour, delivery of breastfeeding is called mother-to-child transmission (MTCT). An estimated 530 000 (410 000 - 660 000) children were newly infected in 2006, mainly through mother-to-child transmission. The vast majority of these infections are preventable, yet coverage levels are remarkably low in most resource-limited countries.

Prevention mother-to-child transmission

| | Total | Year | Comment |
|---|---------|------|---------|
| Antenatal care coverage (%), 1997--2005* | 93 | 2005 | |
| Number of pregnant women counselled on PMTCT services | 159 829 | 2005 | |
| Estimated number of HIV-infected pregnant women | 98 000 | 2005 | |
| Number of HIV-infected pregnant women who received ARVs for PMTCT | 8461 | 2005 | |
| % of HIV-infected pregnant women who received ARVs for PMTCT | 9 | 2005 | |

* Data refer to the most recent year available during the period specified.

Source: UNAIDS/Unicef/WHO. *Children and AIDS: A stocktaking report, Actions and progress during the first year of "Unite for Children, Unite against AIDS". New York, 2007.*

Prevention indicators among injecting drugs users

| Availability of harm reduction services | Number of centers | Number of people attending services | Estimation of coverage | Year |
|---|-------------------|-------------------------------------|------------------------|------|
| Needle exchange programs | | | | |
| Opioid substitute therapy | | | | |

Source:

| | Estimated number of IDUs aged 15-65 | IDU prevalence(%) | Year |
|--------------------------|-------------------------------------|-------------------|------|
| Needle exchange programs | | | |
| | | | |

Source:

Screening of blood transfusions nationwide

Blood safety programs aim to ensure that the majority of blood units are screened for HIV and other infectious agents. This indicator gives an idea of the overall percentage of blood units that have been screened to high enough standards that they can confidently be declared free of HIV.

| | Percentage |
|--|------------|
| Percentage of blood units transfused in the last 12 months that have been adequately screened for HIV according to national or WHO guidelines. | 100% |

Sources

Data presented in this Epidemiological Fact Sheet come from several sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

- Annex 3: Progress on Global Access to HIV Antiretroviral Therapy, A Report on "3 by 5" and Beyond. Geneva, WHO and UNAIDS, March 2006.
- 2006 Report on the global AIDS epidemic
- Based on the most recent calculated ART need estimates by WHO and UNAIDS, as of March 2006.
- Census
- Coverage Survey
- United Nations Population Division
- UNAIDS/Unicef/WHO. Children and AIDS: A stocktaking report, Actions and progress during the first year of "Unite for Children, Unite against AIDS". New York, 2007.
- UNDP Human Development Report 2005
- United Nations Educational, Scientific and Cultural Organization
- UNGASS CR
- UNGASS CR and SIDALAC
- UNICEF Global Database on Skilled Attendant at Delivery. The United Nations Children's Fund. (<http://www.childinfo.org/areas/deliverycare/countrydata.php>)
- UNICEF / WHO
- World Contraceptive Use 2005 database. Population Division, Department of Economic and Social Affairs, United Nations.
- UNPOP Dept. Of Economic and Social Affairs
- World Health Organization
- World Health Organization, 3 by 5
- WHO and UNAIDS, March 2006
- WHO/UNICEF estimates of national coverage for year 2004 (as of September 2005). (http://www.who.int/immunization_monitoring/routine/immunization_coverage/en/index4.html)
- World Bank
- World Health Report 2006, WHO
- Armstrong, K., G. Woelk, S. Gloyd 2000 Socio-Economic Factors Related to HIV Prevalence among Antenatal Clinic Attendees at Sentinel Surveillance Sites Throughout ... XIII International AIDS Conference, Durban, South Africa, 7/9-14, Poster TuPeC3434.
- Cowan, F., L. Langhaug, T. Swarthout, et al. 2001 Interaction of HSV-2 and HIV in Rural Zimbabwean Sex Workers (SW) Presented at International Congress of Sexually Transmitted Infections, ISSTD/JUSTI, Berlin, Germany, 24-27 June 2001, Abstract in International Journal of STD & AIDS, vol. 12, suppl. 2, p.160.
- Cowan, F. M., J. W. Hargrove, L. F. Langhaug, et.al 2005 The Appropriateness of Core Group Interventions Using Presumptive Periodic Treatment Among Rural Zimbabwean Women Who ... Journal of Acquired Immune Deficiency Syndromes, 38(2): 202-207.
- De Muylder, X., M. Laga, C. Tennstedt, et al. 1990 The Role of Neisseria Gonorrhoeae and Chlamydia Trachomatis in Pelvic Inflammatory Disease and Its Sequelae in Zimbabwe Journal of Infectious Diseases, vol. 162, pp. 501-505.
- Easterbrook, P., D. Lamprecht, N. Ives, et al. 2000 Transmission Patterns of Tuberculosis in Harare, Zimbabwe: A Molecular Epidemiological Analysis XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract WePeC4436.
- Guevara, H., E. Johnston, L. Zijenah, et al. 2000 Prenatal Transmission of Subtype C HIV-1 in Zimbabwe: HIV-1 RNA and DNA in Maternal and Cord Blood Journal of Acquired Immune Deficiency Syndromes, vol. 25, no. 5, pp. 390-397.
- Jackson, H. 1993 AIDS Update in Zimbabwe: AIDS Cases ZAINET AIDS News, vol. 1, no. 3, pp. 10-13.
- Kububa, P., L. B. Dube, S. Midzi, et al. 2002 First Suggestion of Decline in HIV Prevalence in Zimbabwe XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Poster LbPeC9034.
- Le Bacq, F., P.R. Mason, L. Gwanzura, et al. 1993 HIV and Other Sexually Transmitted Diseases at a Rural Hospital in Zimbabwe Genitourinary Medicine, vol. 69, pp. 352-256.
- Louie, L. G., W. E. Hartogensis, R. P. Jackman, et al. 2004 Mycobacterium Tuberculosis/HIV-1 Coinfection and Disease: Role of Human Leukocyte Antigen Variation Journal of Infectious Diseases, vol. 189, no. 6, pp. 1084-1090.
- Mahomed, K., J. Kasule, D. Makuyana, et al. 1991 Seroprevalence of HIV Infection amongst Antenatal Women in Greater Harare, Zimbabwe Central African Journal of Medicine, vol. 37, no. 10, pp. 322-325.
- Mahari, M., W. Legg, S. Houston, et al. 1990 Association of Tuberculosis and HIV Infection in Zimbabwe VI International Conference on AIDS, San Francisco, 6/20-24, Abstract Th.B.494.
- Mason, P., S. Ray, C. Mapushere, et al. 1995 Use of Female and Male Condoms by Commercial Sex Workers: Impact on STD Transmission IX International Conference on AIDS and STD in Africa, Kampala, Uganda, 12/10-14, Poster TuC614.

- Olayinka, B. A., C. L. OBI 1999 Symptomatic HIV-Infection in Infants According to Serostatus of Mothers during Pregnancy East African Medical Journal, vol. 76, no. 10, pp. 566-570.
- Tswana, S. A., L. Nystrom, S. R. Moyo, et al. 1995 Hospital-Based Study of Sexually Transmitted Diseases at Murewa Rural District Hospital, Zimbabwe 1991-1992 Sexually Transmitted Diseases, vol. 22, no. 1, pp. 1-6.
- Zimbabwe Ministry of Health 1990 HIV Sentinel Surveillance 1990 Report AIDS Control Programme and Health Information Unit, Ministry of Health, report.
- Zimbabwe Ministry of Health 1991 HIV and AIDS Surveillance Quarterly Report : January, February, March 1991 AIDS Control Programme and Health Information Unit, Ministry of Health, report.
- Zimbabwe Ministry of Health and Child Welfare 1998 HIV, AIDS, STD and TB Fact Sheet Monitoring and Evaluation Unit, National AIDS Coordination Programme, November, report, pp. 1-18.
- Zimbabwe Ministry of Health 1998 HIV Sentinel Surveillance Data Ministry of Health & CW EDC-NHIS Unit, documents.
- Zimbabwe Ministry of Health and Child Welfare 2000 National Survey of HIV and Syphilis Prevalence among Women at Antenatal Clinics in Zimbabwe, 2000 Republic of Zimbabwe, Health Information and Surveillance Unit, Department of Disease Prevention and Control, unpublished report.
- Zimbabwe Ministry of Health and Child Welfare 2002 National Survey of HIV and Syphilis Prevalence among Women Attending Antenatal Clinics in Zimbabwe, 2002 (Based on 2004 ... Ministry of Health and Child Welfare, Zimbabwe National Family Planning Control, unpublished report. (Based on 2001 publication Z0117).
- Zimbabwe Ministry of Health and Child Welfare 2004 National Survey of HIV and Syphilis Prevalence among Women Attending Antenatal Clinics in Zimbabwe, 2004 Ministry of Health and Child Welfare, Health Information and Surveillance Unit, Department of Disease Prevention Control, AIDS & TB Programme, Preliminary report.
- Asamoah-Odei, E. 1999 Surveillance of HIV/AIDS and Related Diseases WHO Travel Report Summary, Zimbabwe, 12 July, unpublished trip report.
- Eriki, P. 1992 HIV Seroprevalence among Patients with Tuberculosis in Selected African Countries VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-11, Poster W.PS.Congo.
- Gregson, S., T. Zhuwau, R. M. Anderson, et al. 1995 Age and Religion Selection Biases in HIV-1 Prevalence Data from Antenatal Clinics in Manicaland, Zimbabwe Central African Journal of Medicine, vol. 41, no. 11, pp. 339-346.
- Legg, W., M. Mahari, S. Houston, et al. 1989 Association of Tuberculosis and HIV Infection in Zimbabwe V International Conference on AIDS, Montreal, 6/4-9, Abstract Th.G.O. 6.
- Latif, A. S. 1995 A Report on a Study to Determine the Aetiology and Pattern of STD amongst Men and Women Presenting to Health Centres in ... University of Zimbabwe Medical School, Dept. of Medicine, September STD report, unpublished.
- Mbizvo, E. M., S. E. Msuya, B. Stray-Pedersen, et al. 2001 HIV Seroprevalence and Its Associations with the Other Reproductive Tract Infections in Asymptomatic Women in Harare, Zimbabwe International Journal of STD and AIDS, vol. 12, no. 8, pp. 524-531.
- Schoch, O. D., H. L. Rieder 1996 Characteristics of Sputum Smear-Positive Tuberculosis Patients with and without HIV Infection in a Hospital in Zimbabwe European Respiratory Journal, vol. 9, pp. 284-287.
- Ticconi, C., M. Maptumo, M. Dorrucchi, et al. 2003 Effect of Maternal HIV and Malaria Infection on Pregnancy and Perinatal Outcome in Zimbabwe Journal of Acquired Immune Deficiency Syndromes, vol. 34, no. 3, pp. 289-294.
- Zhuwau, T. 1997 HIV Trends and Behaviour Changes Presented at the UNAIDS Regional Workshop on "Evidence of Behavioural Change in the Context of HIV Decline in Uganda." 10-13 February, Nairobi, Kenya.
- Zimbabwe Ministry of Health and Child Welfare 1994 HIV, STD, and AIDS Surveillance Workshop: Monomotapa Hotel Health Information Unit, Department of Epidemiology and Disease Control, Ministry of Health and Child Welfare, 7-8 July, report.
- Zimbabwe Ministry of Health and Child Welfare 1997 Summary Results of HIV Sentinel Surveillance AIDS Coordination Programme, fax from Sunday Manyenya, 21 April.

Websites

- www.aids.africa.com

Annex: HIV surveillance prevalence by site

| Group | Area | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | |
|----------------------------|---------------------------|------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Pregnant women | Major urban areas | Attending clinics | | | | | 32 | | | | | 30 | 30.6 | | | | | |
| | | | | | | | | | | | | 31.1 | 27.9 | | | | | |
| | | Budiriro area | | | | | | | 30.4 | | | | | | | | | |
| | | Bulawayo | | 17.1 | | 25.8 | | 30 | | 24 | | | | | | | | |
| | | Chitungwiza | | | | | 41.7 | | | | | | | | | | | |
| | | Edith Opperman Clinic | 17.9 | | | | | | | 28 | | | | | | | | |
| | | Glenview Clinic | 19.4 | | | | | | | | | | | | | | | |
| | | Harare | 23.8 | | | | | | | | 22.7 | | | | | | | |
| | | Harare Hospital | 16 | | | | | | | | 33.3 | | | | | | | |
| | | Kuwadzana Clinic | | | | | | | | | | | | | 20.3 | | 19.7 | |
| | | Mabvuku & Mbare Clinics | | | | | | | | | | | 31.1 | | | | | |
| | | Nkulumane Clinic | | | | | | | | | | | | | 24.8 | | 18.1 | |
| | | Seke North Clinic | | | 34 | | | | | | 32 | | | | | | | |
| | | St. Mary Clinic | | | 28 | | | | | | 34 | | | 33.5 | 30.7 | 28.1 | | 20.3 |
| | | Zengeza clinic | | | 26 | | | | | | | | | | | | | |
| | Outside major urban areas | Antelope Hospital | | | | | 15 | | | 28 | | | | | | | | |
| | | Attending clinic | | | | | | | | 24 | | | | | | | | |
| | | Attending clinics | | | | | 24.3 | | | | | | | | | | | |
| | | | | | | | 14 | | | | | | | | | | | |
| | | Banket | | | | | | | | | | | | 32.6 | | | | |
| | | Banket District Hospital | | | | | | | | | | | | | 30.3 | | 30.3 | |
| | | Beitbridge District Hospital | | | | | | | | | | | | | 28.8 | | 26.3 | |
| | | | | | | | | 23 | 44 | 59 | 46 | 37 | | | | | | |
| | | Beitbridge Dulibadzimu | | | | | | | | | | | 41.4 | 28.5 | | | | |
| | | Bindura Chipadze | | | | | | | | | | | 32.1 | 25.8 | | | | |
| | | Bindura Chipadze Clinic | | | | | | | | | | | | | 32 | | 24.6 | |
| | | Bindura Government Hospital | | | | | 40 | 27 | | 29.3 | | | | | | | | |
| | | Binga District Hospital | | | 17.5 | | 14.4 | | | | 9.2 | | | | | 19.6 | | 10.3 |
| | | Birchenough Bridge | | | 15.5 | 22.3 | | | | | | | | | | | | |
| | | Buhera | | | | | | | | | 50.8 | | | | | | | |
| | | Chinotimba | | | | | | | | | 42.6 | | | 55.7 | 45.3 | | | |
| | | Chinotimba Clinic | | | | | | | | | | | | | 36.3 | | 35.8 | |
| | | Chipinge Hospital | | | | 13.7 | | | | | | | | | | | | |
| Chiradzi District Hospital | | | | | | | | | | | | | 28.8 | | | | | |
| Chiredzi | | | 39.7 | | | | 70.2 | 46.7 | | | | 70.7 | 32.1 | | | | | |

| Group | Area | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|---------------------------|----------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Pregnant women | Outside major urban areas | | | | | | | | | | | | | | | 17.8 | |
| | Chiredzi District Hospital | | | | | | | | | | | | | | | | |
| | Chiredzi Hospital | | | | | | | | 25 | | | | | | | | |
| | Chitsungo Mission Hospital | | | | | | | | 37.3 | | | | | | | | |
| | Commercial farms | | | 23 | 22.5 | 36.2 | | | | | | | | | | | |
| | Eastern Highlands | | | | 15.8 | | | | | | | | | | | | |
| | Gokwe district | | 22.4 | | | | | | | | | | | | | | |
| | Growth Point | | | 22 | 27 | 34 | | | | | | | | | | | |
| | Gutu Mission | | | | | 20.5 | 39.5 | | 25 | | | | 27.4 | 22.7 | | | |
| | Gutu Mission Hospital | | | | | | | | | | | | | | | | 23.4 |
| | Gwanda District Hospital | | | | | | | | | | | | | | | | 26 |
| | | | 16 | 21.1 | 25 | 25 | 33 | 34 | 48 | 31 | | | | | | | |
| | Gwanda Pakama | | | | | | | | | | | | 30.4 | 41.9 | | | |
| | Gwanda Provincial Hospital | | | | | | | | | | | | | | 32.4 | | |
| | Gweru | | 28.1 | 24 | 20.9 | 34.5 | | | | | | | | | | | |
| | Gweru Provincial Hospital | | | | | | | | | | | | | | 26 | | 19.9 |
| | Hauna Growth Point | | | | | 17.9 | | | | | | | | | | | |
| | Hoyuyu | | | | | | 32 | | | | | | | | | | |
| | Hwange Hospital | 12.3 | 24.5 | 19.9 | | | | | | 18.8 | | | | | | | |
| | Hwange St. Patricks | | | | | | | | | | | | 32.6 | | | | |
| | Kadoma District Hospital | | | | | | | | | | | | | | 23.9 | | 21.9 |
| | Karanda Hospital | | | | | 20 | 22.5 | 26 | | 23.9 | | | 18.3 | 19.2 | 19 | | 12.1 |
| | Kariyangwe | | | | | | | | | 7 | | | | | | | |
| | | | | 6.6 | | | | | | | | | 13 | 19.2 | | | |
| | Katsukunye | | | | | | 54 | | | | | | | | | | |
| | Kezi | | | | | | 16 | | | | | | | | | | |
| | Kuwadzana | | | 15.7 | | | | | | | | | 27.8 | 30.4 | | | |
| | Kwekwe district | | 22.4 | 22 | 25 | | | | | | | | | | | | |
| | Makoni | | | | | | | | | 29.3 | | | | | | | |
| | Makosa | | | | | | 34.1 | | | | | | | | | | |
| | Mandava | | 16 | | | | | | | | | | | | | | |
| | Mandava Clinic | | | | | | | | | 44 | | | | | | | |
| Maphisa District Hospital | | | | | | 15 | | | 27.8 | 29 | | | | | | | |
| Mary Mount Hospital | | | | | | | | | 34.9 | | | | | | | | |
| Mashoko Mission Hospital | | | | | 19.1 | 18.1 | 19 | | 30 | | | | | | | | |
| Masvingo | | 30.9 | 42.1 | | | 35.1 | | 36.5 | | | | | | | | | |

| Group | Area | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | | |
|----------------------------------|---------------------------|-------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--|
| Pregnant women | Outside major urban areas | Masvingo Province | | | | | 41.7 | | | | | | | | | | | | |
| | | Mbizo 11 Clinic | | | | | | | 7 | | | | | | | | | | |
| | | Mines | | | | | 24.5 | | | | | | | | | | | | |
| | | Mkoba 1 | | | | | | | | 19.2 | | | | | | | | | |
| | | Mkoba Polyclinic | | | | | | | | 30.7 | | | 36.4 | 28.5 | | | | | |
| | | Mneme & Musume | 7.6 | 7.7 | 7.7 | 20 | 25.5 | | | 30 | | | | | | | | | |
| | | Murambinda Hospital | | | | | | | | | | | | | 21.3 | | 19.7 | | |
| | | Murambinda Mission Hospital | 19.7 | 13.9 | | | | 41.4 | | | | | | 25.7 | 26.4 | | | | |
| | | Musume Mission Hospital | | | | | | | | | | | 54.5 | 26.1 | | | | | |
| | | | | | | | | | | | | | | | | 22.8 | | 22.9 | |
| | | Mutare | | | | 25.1 | | 33.6 | | | 37.7 | | | | | | | | |
| | | Mutoko | | | | | 34.4 | | | | | | | 33.7 | 33.1 | | | | |
| | | Mutoko District Hospital | | | 25.9 | | | | | | 19.1 | | | | | 18.8 | | 20.5 | |
| | | Mutoko Hospital | | | | 25.9 | | | | | | | | | | | | | |
| | | Mutorashanga | | | 15 | | | | | | | | | | | | | | |
| | | Mvurwi District Hospital | | | | | | | | | 27 | | | | | | | | |
| | | Nkulumane | | | | | | | | | | | | 31.1 | 27.9 | | | | |
| | | Nyamuzuwe | | | | | | 38 | | | | | | | | | | | |
| | | Plumtree Hospital | | 9.9 | 11.1 | | | | | | | | | | | | | | |
| | | Rusape Hospital | 31.6 | 33.8 | | | | | 67 | | | | | | | | | | |
| | | Sadza District Hospital | | | | | | | | | 31.4 | | | 42.7 | 31.5 | 34.2 | | | |
| | | | | | | | | | | | | | | | | 34.2 | | 22.9 | |
| | | Sakubva | | | | | | | | | | | | 33.9 | 21.3 | | | | |
| | | Sakubva Clinic | | | | | | | | | | | | | | 22.6 | | 17.7 | |
| | | Sanyati Hospital | | 20 | 18.5 | 18.5 | | | | | | | | 36.9 | 33.9 | | | | |
| | | Shabanie Mines | | 17 | | | | | | | | | | | | | | | |
| | | Shurugwi Mines | | 23.7 | 20 | 17 | | | | | | | | | | | | | |
| | | St. Alberts | | | | | | | 23 | | 23 | | | | | | | | |
| | | St. Alvert's Mission Hospital | | | | | | | | | | | | | 8.3 | | | | |
| | | Venghere Clinic | | | | | | | | | 53.3 | | | | | | | | |
| Wankie Colliery Company Hospital | | 11.1 | | | | 22.9 | | | | | | | | | | | | | |
| Zvimba/Kadoma | | | | | | | | | 24.3 | | | | | | | | | | |
| Sex workers | Major urban areas | | | | | | 86 | | | | | | | | | | | | |
| | Outside major urban areas | | | | | | | | | | | | | | | | 57.2 | | |
| | | | | | | | | | | | | | 50 | | | | | | |
| Injecting drug users | | | | | | | | | | | | | | | | | | | |

| Group | Area | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | |
|----------------------------------|---------------------------|------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--|
| STI patients | Major urban areas | Bulawayo | | 39 | | 60.3 | | 71 | | | | | | | | | | |
| | | Harare | 52 | 51 | | | | 61 | | | | | | | | | | |
| | | Seke North Clinic | | | 52 | | | | | | | | | | | | | |
| | | St. Mary Clinic | | | 70 | | | | | | | | | | | | | |
| | | Zengeza clinic | | | 56 | | | | | | | | | | | | | |
| | Outside major urban areas | Beitbridge District Hospital | | | | | 73 | 65 | | | | | | | | | | |
| | | Bindura Government Hospital | | | | | 56 | 43 | | | | | | | | | | |
| | | Binga District Hospital | | | 44.6 | | | | | | | | | | | | | |
| | | Commercial farms | | | | | 51 | | | | | | | | | | | |
| | | Gokwe district | | 48.8 | 49.4 | 32.2 | | | | | | | | | | | | |
| | | Gutu | | | | 71.8 | | | | | | | | | | | | |
| | | Gwanda District Hospital | | 33.1 | 40 | | 66 | 66 | | | | | | | | | | |
| | | Gweru | | 48.2 | 48 | 52 | 52 | | | | | | | | | | | |
| | | Hwange Hospital | 28.7 | 42 | 42.3 | | | | | | | | | | | | | |
| | | Karoi district | | 62.9 | | | | | | | | | | | | | | |
| | | Kuwadzana | | | 42.4 | | | | | | | | | | | | | |
| | | Kwekwe | | | 48.8 | | | | | | | | | | | | | |
| | | Kwekwe district | | 45.6 | | 55.7 | | | | | | | | | | | | |
| | | Mashoko Mission Hospital | | | | | 51.9 | 58.4 | | | | | | | | | | |
| | | Masvingo | | 58.5 | 59.5 | | 75.4 | | 71.8 | | | | | | | | | |
| | | Mines | | | | | 48 | | | | | | | | | | | |
| | | Mneme & Musume | 24.3 | 24.5 | | 46.8 | | | | | | | | | | | | |
| | | Murambinda Mission Hospital | | | | | | 69.3 | | | | | | | | | | |
| | | Murewa District Hospital | | | 46 | | | | | | | | | | | | | |
| | | Mutare | | | | 25.6 | | 56.6 | | | | | | | | | | |
| | | Mutoko District Hospital | | | 51.6 | | | | | | | | | | | | | |
| | | Mutoko Hospital | | | | 52.1 | | | | | | | | | | | | |
| | | Mutorashanga | | | 39 | | | | | | | | | | | | | |
| | | Plumtree Hospital | | 33.3 | 43.1 | | | | | | | | | | | | | |
| | | Rusape Hospital | 59.7 | 56.4 | | | | 87.7 | | | | | | | | | | |
| | | Sanyati Hospital | | 45.6 | | 62.2 | | | | | | | | | | | | |
| | | Shabanie Mines | | 46.7 | | | | | | | | | | | | | | |
| Shurugwi Mines | | 39.6 | 43 | 41 | | | | | | | | | | | | | | |
| Wankie Colliery Company Hospital | | 32.6 | | | | | | | | | | | | | | | | |

Men having sex with men

| Group | Area | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|-----------------------|---------------------------|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Tuberculosis patients | Major urban areas | Beatrice Road Hospital | | | | | | | 74 | | | | | | | | |
| | | Beatrice Road Infectious Diseases Hospital | | | | | | | | | 73.6 | | | | | | |
| | | Bulawayo | | 56.5 | | 59.7 | | | | | | | | | | | |
| | Outside major urban areas | Harare | | | | | | | | | | | | | | | |
| | | Driefontein Tuberculosis Sanatorium | | | 64.8 | | | | | | | | | | | | |
| | | Not specified | | | | | | | | | | | | | | | |
| | | Rusape Hospital | | | | | | 84.3 | | | | | | | | | |