

# CLIMATE AND HEALTH COUNTRY PROFILE – 2015

## IRAN



United Nations  
Framework Convention on  
Climate Change



### OVERVIEW

The Islamic Republic of Iran has a diverse geography with vast coastlines, mountains, forested areas, plains and deserts. It is predominantly a dry climate with over 80% of the country being arid or semi-arid.<sup>a</sup> In past decades, environmental resources and ecosystems have been threatened due to lack of precipitation, persistent drought, heat-stress, air pollution, water pollution, soil erosion and loss of biodiversity.<sup>a</sup> Iran is highly vulnerable to the impacts of climate change, including rising temperatures, sea-level rise and increased frequency of natural disasters. Climate change could result in increased heat-related mortality, food insecurity, disruption of livelihoods, internal migration and changing distribution patterns of infectious and vector-borne diseases.

Iran is yet to develop an official national climate change action plan, which could prove beneficial for mitigation and adaptation strategies that target public health in the context of climate change. Iran has envisioned climate change within the broader objective of achieving sustainable development.

### SUMMARY OF KEY FINDINGS

- In Iran, under a high emissions scenario, mean annual temperature is projected to rise by about 6.2°C on average from 1990 to 2100. If global emissions decrease rapidly, the temperature rise is limited to about 1.7°C [page 2].
- In Iran, under a high emissions scenario, the number of days of warm spell<sup>b</sup> is projected to increase from about 10 days in 1990 to about 215 days on average in 2100. If global emissions decrease rapidly, the days of warm spell are limited to about 50 on average [page 2].
- In Iran, under a high emissions scenario, and without large investments in adaptation, an annual average of about

184,700 people are projected to be affected by flooding due to sea level rise between 2070 and 2100. If global emissions decrease rapidly and there is a major scale up in protection the annual affected population could be limited to about 200 people [page 3].

- in Iran, under a high emissions scenario heat-related deaths in the elderly [65+ years] are projected to increase to almost 70 deaths per 100,000 by 2080 compared to the estimated baseline of under 6 deaths per 100,000 annually between 1961 and 1990. A rapid reduction in global emissions could limit heat-related deaths in the elderly to about 16 deaths per 100,000 in 2080 [page 4].

### OPPORTUNITIES FOR ACTION

Iran has conducted a national assessment of climate change impacts, vulnerability and adaptation for health. Additionally, Iran is currently implementing projects on health adaptation to climate change and is taking action to build institutional and technical capacities to work on climate change and health. Country reported data [see section 6] indicate there are further opportunities for action in the following areas:

#### 1) Adaptation

- Implement activities to increase climate resilience of health infrastructure.
- Estimate the costs to implement health resilience to climate change and include these costs in planned allocations.

#### 2) Mitigation

- Conduct a valuation of co-benefits to health of climate change mitigation policies.

#### 3) National policy implementation

- Develop a national health adaptation strategy.

### DEMOGRAPHIC ESTIMATES

Population [2013] <sup>c</sup>	77.15 million
Population growth rate [2013] <sup>c</sup>	1.3 %
Population living in urban areas [2013] <sup>d</sup>	72.3 %
Population under five [2013] <sup>c</sup>	8.8 %
Population aged 65 or older [2013] <sup>c</sup>	4.9 %

### ECONOMIC AND DEVELOPMENT INDICATORS

GDP per capita [current US\$, 2013] <sup>e</sup>	6,631 USD
Total expenditure on health as % of GDP [2013] <sup>f</sup>	6.7 %
Percentage share of income for lowest 20% of population [2012] <sup>g</sup>	NA
HDI [2013, +/- 0.01 change from 2005 is indicated with arrow] <sup>h</sup>	0.749 ▲

### HEALTH ESTIMATES

Life expectancy at birth [2013] <sup>i</sup>	74 years
Under-5 mortality per 1000 live births [2013] <sup>i</sup>	17

a Iran's 2nd national communication to UNFCCC <http://unfccc.int/resource/docs/natc/iranc2.pdf>

b A 'warm spell' day is a day when maximum temperature, together with that of at least the 6 consecutive previous days, exceeds the 90th percentile threshold for that time of the year.

c World Population Prospects: The 2015 Revision, UNDESA [2015]

d World Urbanization Prospects: The 2014 Revision, UNDESA [2014]

e World Development Indicators, World Bank [2015]

f Global Health Expenditure Database, WHO [2014]

g United Nations Development Programme, Human Development Reports [2014]

h Global Health Observatory, WHO; 2014

i Levels & Trends in Child Mortality Report 2015, The UN Inter-agency Group for Child Mortality Estimation [2015]

# 1

## CURRENT AND FUTURE CLIMATE HAZARDS


Due to climate change, many climate hazards and extreme weather events, such as heat waves, heavy rainfall and droughts, could become more frequent and more intense in many parts of the world.

Outlined here are country-specific projections up to the year 2100 for climate hazards under a 'business as usual' high emissions scenario compared to projections under a 'two-degree' scenario with rapidly decreasing global emissions. Most hazards caused by climate change will persist for many centuries.

### COUNTRY-SPECIFIC CLIMATE HAZARD PROJECTIONS


The model projections below present climate hazards under a high emissions scenario, Representative Concentration Pathway 8.5 [RCP8.5] (in orange) and a low emissions scenario, [RCP2.6] (in green).<sup>a</sup> The text boxes describe the projected changes averaged across about 20 models (thick line). The figures also show each model individually as well as the 90% model range (shaded) as a measure of uncertainty and, where available, the annual and smoothed observed record (in blue).<sup>b,c</sup>

#### MEAN ANNUAL TEMPERATURE




Under a high emissions scenario, mean annual temperature is projected to rise by about 6.2°C on average from 1990 to 2100. If emissions decrease rapidly, the temperature rise is limited to about 1.7°C.

#### DAYS OF WARM SPELL ('HEAT WAVES')




Under a high emissions scenario, the number of days of warm spell<sup>d</sup> is projected to increase from about 10 days in 1990 to about 215 days on average in 2100. If emissions decrease rapidly, the days of warm spell are limited to about 50 on average.

#### DAYS WITH EXTREME RAINFALL ('FLOOD RISK')



Under a high emissions scenario, the number of days with very heavy precipitation [20 mm or more] could increase by an average of about 2 days on average from 1990 to 2100. Some models indicate larger increases but the total number of such days remains low. If emissions decrease rapidly, the increase in risk is somewhat reduced.

#### CONSECUTIVE DRY DAYS ('DROUGHT')



Under both high and low emissions scenarios, the longest dry spell is not indicated to change much from an average of about 90 days, with continuing large year-to-year variability.

<sup>a</sup> Model projections are from CMIP5 for RCP8.5 (high emissions) and RCP2.6 (low emissions). Model anomalies are added to the historical mean and smoothed.

<sup>b</sup> Observed historical record of mean temperature is from CRU-TSV.3.22; observed historical records of extremes are from HadEX2.


<sup>c</sup> Analysis by the Climatic Research Unit and Tyndall Centre for Climate Change Research, University of East Anglia, 2015.

<sup>d</sup> A 'warm spell' day is a day when maximum temperature, together with that of at least the 6 consecutive previous days, exceeds the 90th percentile threshold for that time of the year.

## CURRENT AND FUTURE HEALTH RISKS DUE TO CLIMATE CHANGE

Human health is profoundly affected by weather and climate. Climate change threatens to exacerbate today's health problems – deaths from extreme weather events, cardiovascular and respiratory diseases, infectious diseases and malnutrition – whilst undermining water and food supplies, infrastructure, health systems and social protection systems.

### ANNUAL EXPOSURE TO FLOODING DUE TO SEA LEVEL RISE (PERIOD 2070–2100)



\* Medium ice melting scenario


\*\* Values rounded to nearest '00

Under a high emissions scenario, and without large investments in adaptation, an annual average of about 184,700 people are projected to be affected by flooding due to sea level rise between 2070 and 2100. If global emissions decrease rapidly and there is a major scale up in protection [i.e. continued construction/raising of dikes] the annual affected population could be limited to about 200 people. Adaptation alone will not offer sufficient protection, as sea level rise is a long-term process, with high emissions scenarios bringing increasing impacts well beyond the end of the century.

Source: Human dynamics of climate change, technical report, Met Office, HM Government, UK, 2014.

### INFECTIOUS AND VECTOR-BORNE DISEASES


#### Estimated number of deaths due to diarrhoeal disease in children under 15 years in Iran (base case scenario for economic growth)



In the baseline year of 2008 there were an estimated 5,900 diarrhoeal deaths in children under 15 years old. Under a high emissions scenario, diarrhoeal deaths attributable to climate change in children under 15 years old are projected to be approximately 12.7% of about 2,300 diarrhoeal deaths projected in 2030. Although diarrhoeal deaths are projected to decline to approximately 1,100 by 2050, the proportion of deaths attributable to climate change will rise to about 17.5%.

Source: Lloyd, S., 2015.<sup>c</sup>

#### Mean relative vectorial capacity for dengue fever transmission in Iran



#### KEY IMPLICATIONS FOR HEALTH

In addition to deaths from drowning, flooding causes extensive indirect health effects, including impacts on food production, water provision, ecosystem disruption, infectious disease outbreak and vector distribution. Longer term effects of flooding may include post-traumatic stress and population displacement.



#### KEY IMPLICATIONS FOR HEALTH

Some of the world's most virulent infections are also highly sensitive to climate: temperature, precipitation and humidity have a strong influence on the life-cycles of the vectors and the infectious agents they carry and influence the transmission of water and food-borne diseases.<sup>a</sup>

Socioeconomic development and health interventions are driving down burdens of several infectious diseases, and these projections assume that this will continue. However, climate conditions are projected to become significantly more favourable for transmission, slowing progress in reducing burdens, and increasing the populations at risk if control measures are not maintained or strengthened.<sup>b</sup>

Under both high and low emissions scenarios, mean relative vectorial capacity remains relatively constant (about 0.23–0.25) towards 2070. Co-factors such as urbanization, development and population movements may modify the disease burdens associated with dengue, and make the disease cross new subnational borders.

Source: Rocklöv, J., Quam, M. et al., 2015.<sup>c</sup>


a Atlas of Health and Climate, World Health Organization and World Meteorological Organization, 2012.

b Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s. Geneva: World Health Organization, 2014.

c Country-level analysis, completed in 2015, was based on health models outlined in the Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s. Geneva: World Health Organization, 2014. The mean of impact estimates for three global climate models are presented. Models assume continued socio-economic trends [SSP2 or comparable].

## HEAT-RELATED MORTALITY

Heat-related mortality in population 65 years or over, Iran (deaths / 100,000 population 65+ yrs)



Under a high emissions scenario heat-related deaths in the elderly (65+ years) are projected to increase to almost 70 deaths per 100,000 by 2080 compared to the estimated baseline of under 6 deaths per 100,000 annually between 1961 and 1990. A rapid reduction in global emissions could limit heat-related deaths in the elderly to about 16 deaths per 100,000 in 2080.

Source: Honda et al., 2015.<sup>a</sup>



## KEY IMPLICATIONS FOR HEALTH

Climate change is expected to increase mean annual temperature and the intensity and frequency of heat waves resulting in a greater number of people at risk of heat-related medical conditions.

The elderly, children, the chronically ill, the socially isolated and at-risk occupational groups are particularly vulnerable to heat-related conditions.

## UNDERNUTRITION

Climate change, through higher temperatures, land and water scarcity, flooding, drought and displacement, negatively impacts agricultural production and causes breakdown in food systems. These disproportionately affect those most vulnerable to hunger and can lead to food insecurity. Vulnerable groups risk further deterioration into food and nutrition crises if exposed to extreme weather events.<sup>b</sup>

Without considerable efforts made to improve climate resilience, it has been estimated that globally the risk of hunger and malnutrition could increase by up to 20 percent by 2050.<sup>b</sup>

In Iran, the prevalence of underweight children under age 5 was 4.1% in 2011.<sup>c</sup>

<sup>a</sup> Country-level analysis, completed in 2015, was based on health models outlined in the Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s. Geneva: World Health Organization, 2014. The mean of impact estimates for three global climate models are presented. Models assume continued socio-economic trends [SSP2 or comparable].

<sup>b</sup> World Food Project 2015 <https://www.wfp.org/content/two-minutes-climate-change-and-hunger>


<sup>c</sup> World Health Organization, Global Database on Child Growth and Malnutrition [2015 edition]. Child malnutrition estimates are for % underweight, defined as: Percentage of children aged 0–59 months who are below minus two standard deviations from median weight-for-age of the World Health Organization (WHO) Child Growth Standards. Note, % underweight estimates for Iran for 2011 are pending re-analysis.

## CURRENT EXPOSURES AND HEALTH RISKS DUE TO AIR POLLUTION

Many of the drivers of climate change, such as inefficient and polluting forms of energy and transport systems, also contribute to air pollution. Air pollution is now one of the largest global health risks, causing approximately seven million deaths every year. There is an important opportunity to promote policies that both protect the climate at a global level, and also have large and immediate health benefits at a local level.

### OUTDOOR AIR POLLUTION EXPOSURE

Outdoor air pollution in cities in Iran  
annual mean PM<sub>2.5</sub> (µg/m<sup>3</sup>) 2010\*



The cities for which there was air pollution data available in 2010 had annual mean PM<sub>2.5</sub> levels that were above the WHO guideline value of 10 µg/m<sup>3</sup>.

Source: Ambient Air Pollution Database, WHO, May 2014. \* A standard conversion has been used for the city of Tehran, see source for further details.



### KEY IMPLICATIONS FOR HEALTH

Outdoor air pollution can have direct and sometimes severe consequences for health.

Fine particles which penetrate deep into the respiratory tract subsequently increase mortality from respiratory infections, lung cancer and cardiovascular disease.

### OUTDOOR AIR POLLUTION EXPOSURE AND SHORT-LIVED CLIMATE POLLUTANTS

Short-lived climate pollutants such as black carbon, methane and tropospheric ozone – released through inefficient use and burning of biomass and fossil fuels for transport, housing, power production, industry, waste disposal (municipal and agricultural) and forest fires – are responsible for a substantial fraction of global warming as well as air pollution related deaths and diseases.

Since short-lived climate pollutants persist in the atmosphere for weeks or months while CO<sub>2</sub> emissions persist for years, significant reductions of SLCP emissions could result in immediate health benefits and health cost savings,<sup>a</sup> and generate very rapid climate benefits – helping to reduce near-term climate change by as much as 0.5°C before 2050.<sup>a</sup>

In Iran, it is estimated that a reduction in SLCPs\* could prevent about 5,800 premature deaths attributed to outdoor air pollution per year, from 2030 onwards (Shindell, D., Science, 2012).

\* Through implementation of 14 reduction measures: 7 targeting methane emissions and the rest, emissions from incomplete combustion. See source for further detail.

<sup>a</sup> United Nations Environment Programme. Reducing Climate-related Air Pollution and Improving Health: Countries can act now and reap immediate benefits. <http://www.unep.org/ccac/Media/PressReleases/ReducingClimate-relatedAirPollution/tabid/131802/language/en-US/Default.aspx>

## CO-BENEFITS TO HEALTH FROM CLIMATE CHANGE MITIGATION: A GLOBAL PERSPECTIVE

Health co-benefits are local, national and international measures with the potential to simultaneously yield large, immediate public health benefits and reduce the upward trajectory of greenhouse gas emissions. Lower carbon strategies can also be cost-effective investments for individuals and societies.

Presented here are examples, from a global perspective, of opportunities for health co-benefits that could be realised by action in important greenhouse gas emitting sectors.<sup>a</sup>

### Transport

Transport injuries lead to 1.2 million deaths every year, and land use and transport planning contribute to the 2–3 million deaths from physical inactivity. The transport sector is also responsible for some 14% [7.0 GtCO<sub>2</sub>e] of global carbon emissions. The IPCC has noted significant opportunities to reduce energy demand in the sector, potentially resulting in a 15%–40% reduction in CO<sub>2</sub> emissions, and bringing substantial opportunities for health: A modal shift towards walking and cycling could see reductions in illnesses related to physical inactivity and reduced outdoor air pollution and noise exposure; increased use of public transport is likely to result in reduced GHG emissions; compact urban planning fosters walkable residential neighborhoods, improves accessibility to jobs, schools and services and can encourage physical activity and improve health equity by making urban services more accessible to the elderly and poor.



### Electricity Generation

Current patterns of electricity generation in many parts of the world, particularly the reliance on coal combustion in highly polluting power plants, contribute heavily to poor local air quality, causing cancer, cardiovascular and respiratory disease. Outdoor air pollution is responsible for 3.7 million premature deaths annually. High-income countries still have work to do in transitioning to cleaner and healthier energy sources.

The health benefits of transitioning from fuels such as coal to lower carbon sources, including ultimately to renewable energy, are clear: Reduced rates of cardiovascular and respiratory disease such as stroke, lung cancer, coronary artery disease, and COPD; cost-savings for health systems; improved economic productivity from a healthier and more productive workforce.



### Food and Agriculture

Agricultural emissions account for some 5.0–5.8 GtCO<sub>2</sub>e annually, with food and nutrition constituting an important determinant of health. Many high-income countries are feeling the burden of poor diet and obesity-related diseases, with some 1.9 billion adults overweight globally.

A wide range of interventions designed to reduce emissions from agriculture and land-use will also yield positive benefits for public health. For example, policy and behavioural interventions to encourage a reduction in red meat consumption and a shift towards local and seasonal fruit and vegetables, which tend to have lower carbon emissions associated with their production, will improve diets and result in reductions in cardiovascular disease and colorectal cancer.



### Healthcare Systems

Health care activities are an important source of greenhouse gas emissions. In the US and in EU countries, for example, health care activities account for between 3–8% of greenhouse gas [CO<sub>2</sub>-eq] emissions. Major sources include procurement and inefficient energy consumption. Modern, on-site, low-carbon energy solutions (e.g. solar, wind, or hybrid solutions) and the development of combined heat and power generation capacity in larger facilities offer significant potential to lower the health sector's carbon footprint, particularly when coupled with building and equipment energy efficiency measures. Where electricity access is limited and heavily reliant upon diesel generators, or in the case of emergencies when local energy grids are damaged or not operational, such solutions can also improve the quality and reliability of energy services. In this way, low carbon energy for health care could not only mitigate climate change, it could enhance access to essential health services and ensure resilience.




<sup>a</sup> For a complete list of references used in the health co-benefits text please see the Climate and Health Country Profile Reference Document, <http://www.who.int/globalchange/en/>

# 5

## EMISSIONS AND COMMITMENTS

Global carbon emissions increased by 80% from 1970 to 2010, and continue to rise.<sup>a,b</sup> Collective action is necessary, but the need and opportunity to reduce greenhouse gas emissions varies between countries. Information on the contribution of different sectors, such as energy, manufacturing, transport and agriculture, can help decision-makers to identify the largest opportunities to work across sectors to protect health, and address climate change.

### IRAN ANNUAL GREENHOUSE GAS EMISSIONS (metric tonnes CO<sub>2</sub> equivalent)



A 2°C upper limit of temperature increase relative to pre-industrial levels has been internationally agreed in order to prevent severe and potentially catastrophic impacts from climate change. Reductions are necessary across countries and sectors. In order to stay below the 2°C upper limit it is estimated that global annual CO<sub>2</sub>-energy emissions, currently at 5.2 tons per capita, need to be reduced to 1.6 tons per capita.<sup>c</sup>

The most recent greenhouse gas emissions data for Iran is from the year 2000. At that time, carbon emissions were increasing across most sectors, with the largest contributions from the energy, transport and 'other' sectors. Through intersectoral collaboration, the health community can help to identify the best policy options not only to eventually stabilize greenhouse gas emissions, but also to provide the largest direct benefits to health.

Source: UNFCCC Greenhouse Gas Data Inventory, UNFCCC [2015].

### NATIONAL RESPONSE

**1992<sup>d</sup>**

**IRAN SIGNED THE UNFCCC**

**2005<sup>d</sup>**

**IRAN RATIFIED THE KYOTO PROTOCOL**

**2030<sup>e</sup>**

**ON THE BASIS OF NATIONAL CAPABILITIES, FINANCIAL RESOURCES AVAILABLE AND REQUIREMENTS OF THE NATIONAL DEVELOPMENT PROGRAM, TAKING INTO ACCOUNT GHGS EMISSION SCENARIOS, THE ISLAMIC REPUBLIC OF IRAN INTENDS TO PARTICIPATE BY MITIGATING ITS GHGS EMISSION IN 2030 BY 4% COMPARED TO THE BUSINESS AS USUAL (BAU) SCENARIO. SUBJECT TO TERMINATION AND NON-EXISTENCE OF SANCTIONS, AVAILABILITY OF INTERNATIONAL RESOURCES IN THE FORM OF FINANCIAL SUPPORT AND TECHNOLOGY TRANSFER, EXCHANGE OF CARBON CREDITS, ACCESSIBILITY OF BILATERAL OR MULTILATERAL IMPLEMENTATION MECHANISMS, TRANSFER OF CLEAN TECHNOLOGIES AS WELL AS CAPACITY BUILDING, THE ISLAMIC REPUBLIC OF IRAN HAS THE POTENTIAL OF MITIGATING ADDITIONAL GHGS EMISSION UP TO 8% AGAINST THE BAU SCENARIO (I.E. 12% IN TOTAL).**

a Boden, T.A., G. Marland, and R.J. Andres [2010]. Global, Regional, and National Fossil-Fuel CO<sub>2</sub> Emissions. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tenn., U.S.A. doi 10.3334/CDIAC/00001\_V2010.  
 b IPCC [2014] Blanco G., R. Gerlagh, S. Suh, J. Barrett, H.C. de Coninck, C.F. Diaz Morejon, R. Mathur, N. Nakicenovic, A. Ofosu Ahenkora, J. Pan, H. Pathak, J. Rice, R. Richels, S.J. Smith, D.J. Stern, F.L. Toth, and P. Zhou, 2014: Drivers, Trends and Mitigation. In: Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Edenhofer, O., R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S. Brunner, P. Eickemeier, B. Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, T. Zwickel and J.C. Minx (eds.)]. Cambridge University Press, Cambridge, United Kingdom  
 c Pathways to deep decarbonization, Sustainable development Solutions Network, 2014 report.  
 d Columbia Law School, 'Climate Change Laws Of The World'. N.p., 2015.  
 e Iran INDC, 19 November 2015.

The following table outlines the status of development or implementation of climate resilient measures, plans or strategies for health adaptation and mitigation of climate change [reported by countries].<sup>a</sup>

<b>GOVERNANCE AND POLICY</b>	
Country has identified a national focal point for climate change in the Ministry of Health	✓
Country has a national health adaptation strategy approved by relevant government body	✗
The National Communication submitted to UNFCCC includes health implications of climate change mitigation policies	✓
<b>HEALTH ADAPTATION IMPLEMENTATION</b>	
Country is currently implementing projects or programmes on health adaptation to climate change	✓
Country has implemented actions to build institutional and technical capacities to work on climate change and health	✓
Country has conducted a national assessment of climate change impacts, vulnerability and adaptation for health	✓
Country has climate information included in Integrated Disease Surveillance and Response (IDSR) system, including development of early warning and response systems for climate-sensitive health risks	✓
Country has implemented activities to increase climate resilience of health infrastructure	✗
<b>FINANCING AND COSTING MECHANISMS</b>	
Estimated costs to implement health resilience to climate change included in planned allocations from domestic funds in the last financial biennium	✗
Estimated costs to implement health resilience to climate change included in planned allocations from international funds in the last financial biennium	✗
<b>HEALTH BENEFITS FROM CLIMATE CHANGE MITIGATION</b>	
The national strategy for climate change mitigation includes consideration of the health implications (health risks or co-benefits) of climate change mitigation actions	✓
Country has conducted valuation of co-benefits of health implications of climate mitigation policies	✗

a Supporting monitoring efforts on health adaptation and mitigation of climate change: a systematic approach for tracking progress at the global level. WHO survey, 2015.

For further information please contact:

**World Health Organization**

20 Avenue Appia  
1211 Geneva 27  
Switzerland

Tel.: +41 22 791 3281 | Fax: +41 22 791 4853  
<http://www.who.int/globalchange/en/>



**United Nations**  
Framework Convention on  
Climate Change

© World Health Organization 2016

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

The estimates and projections provided in this document have been derived using standard categories and methods to enhance their cross-national comparability. As a result, they should not be regarded as the nationally endorsed statistics of Member States which may have been derived using alternative methodologies.

To ensure readability, health estimates and projections have been presented without the margins of uncertainty which are available upon request.