

19. Wagner AK, Graves AJ, Fan Z, Walker S, Zhang F, Ross-Degnan D. Need for and access to health care and medicines: are there gender inequities? *PLoS ONE* 2013;8:e57228. doi: <http://dx.doi.org/10.1371/journal.pone.0057228> PMID:23505420
20. *World health report – health systems financing: the path to universal coverage*. Geneva: World Health Organization; 2010. Available from: <http://www.who.int/whr/2010/en/index.html> [accessed 26 July 2013].
21. Bustreo F, Hunt P, Gruskin S, Eide A, McGoey L, Rao S et al. *Women's and children's health: evidence of impact of human rights*. Geneva: World Health Organization; 2013. Available from: <http://www.ifhro.org/news-a-events/435-womens-and-childrens-health-and-human-rights.html> [accessed 26 July 2013].
22. Broker TR. Global prevention and management of Human Papillomavirus related diseases: the pressing challenges and the compelling opportunities. *Vaccine* 2012;30:vii–x. doi: <http://dx.doi.org/10.1016/j.vaccine.2012.07.071> PMID:23199970

Corrigendum

In Volume 91, Issue 8, August 2013, on page 612: the city for affiliation “c” should be spelt “Ghent”.

In Volume 91, Issue 8, August 2013, on page 566: the group sizes need to be added for sexual assault in Table 5.

Table 5. Fraction of homicides related to child abuse and neglect or to sexual assault, by sex and age of victim, South Africa 2009

Age (years)	Percentage (95% CI)			P
	All	Males	Females	
Child abuse and neglect	(n = 454) ^a	(n = 174)	(n = 276)	
<5	73.8 (66.4–81.2)	45.4 (39.5–51.5)	54.5 (48.5–60.5)	0.0647
5–9	11.7 (7.0–16.3)	24.8 (9.3–51.4)	75.2 (48.6–90.7)	0.0005
10–14	8.0 (3.9–12.0)	22.4 (6.9–52.8)	77.6 (47.2–93.1)	0.0000
15–17	6.6 (3.6–9.7)	8.3 (1.5–35.6)	91.7 (64.4–98.5)	0.0001
Sexual assault	(n = 102)	(n = 10)	(n = 92)	
<5	17.9 (2.8–33.1)	22.2 (6.3–54.8)	77.8 (45.2–93.7)	0.0832
5–9	30.9 (15.7–46.0)	0.0	100	0.0000
10–14	29.1 (18.3–39.8)	13.7 (2.4–50.8)	86.3 (49.2–97.6)	0.0002
15–17	22.1 (11.0–33.3)	8.0 (1.4–34.9)	92.0 (65.1–98.6)	0.0000

CI, confidence interval.

^a This value includes four cases that were excluded from the subgroup analysis because the sex and cause of death could not be determined owing to the decomposition of the body.

In Volume 91, Issue 4, April 2013, on page 242: the first sentence in the forth paragraph of the third column should read: “According to Dr Olayinka Ogunleye, who works in the hypertension unit at that hospital, it costs around 1000 Nigerian Naira (US\$ 6.36) for a patient to have his or her blood pressure checked, although there are opportunities to get blood pressure checked for free in Nigeria as well.”

continues ...