

REPORT OF BOSNIA AND HERZEGOVINA

TABLE OF CONTENTS

1	Introduction
2	Methodology
2.1	Sampling
2.2.1	Geographic Component of the Survey
2.2.2	The map
2.3	Survey Metrics
2.3.1	Sample deviation index
2.3.2	HH Sample deviation index
2.3.3	Response rate
2.3.4	Missing data
2.3.5	Reliability
3	Socio-Demographic Profile
3.1	Household composition by residence and age groups according to income quintile
3.2	Selected age distributions by income quintile
3.3	Characteristics of head of household by income quintile
3.4	Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex
3.5	Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education
4	Risk Factors
4.1	Prevalence of smoking and average daily tobacco consumption
4.2	Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers
4.3	Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity
4.4	Access to improved drinking-water: percent distribution of households by

- type of access to drinking water
- 4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility
- 4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used
- 5 Coverage
 - 5.01 Tuberculosis
 - 5.02 HIV Prevention and Screening
 - 5.03 Antenatal Care and Mother to Child Transmission of HIV
 - 5.04 Birth Delivery Care
 - 5.05 Child's Health - Common Illness
 - 5.06 Child's Health - Danger Signs
 - 5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card
 - 5.08 Vaccination Status of Children 12-23 Months
 - 5.09 Women's Cancer Screening
 - 5.10 Oral Health and Injuries
 - 5.11 Self Reported Non-Communicable Conditions
 - 5.12 Mental Health Conditions
 - 5.13 Cataracts
- 6 Health State Descriptions
 - 6.01 In general, rate your health today - q2000
 - 6.02 Difficulty with 'work and household activities' in last 30 days - q2001
 - 6.03 Difficulty with 'moving around' in last 30 days - q2010
 - 6.04 Difficulty with 'vigorous activities' in last 30 days - q2011
 - 6.05 Difficulty with 'self-care' in last 30 days - q2020
 - 6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021
 - 6.07 Amount of 'bodily aches or pains' in last 30 days - q2030

- 6.08 Amount of 'bodily discomfort' in last 30 days - q2031
- 6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050
- 6.10 Difficulty with 'learning a new task' in last 30 days - q2051
- 6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060
- 6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061
- 6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071
- 6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072
- 6.15 Problem with 'sleeping' in last 30 days - q2080
- 6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081
- 6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090
- 6.18 Problem with 'worry or anxiety' in last 30 days - q2091
- 6.19 Adjusted for vignette responses
- 7 Health System Responsiveness
 - 7.1 Self-reported utilization of health services and perceived unmet need
 - 7.2 Patient assessed characteristics of ambulatory health services
 - 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1
 - 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2
 - 7.4 Patient assessed characteristics of hospital inpatient health services
 - 7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1
 - 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2
 - 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"
- 8 Health Expenditure, Insurance and Human Resources for Health

- 8.0 Expenditure Background Indicators (2002)
- 8.1 Distribution of health payments across different population groups
- 8.2 Percent distribution of structure of out-of-pocket health payments
- 8.3 Financial sources used by households for paying for health services
- 8.4 Insurance Coverage
- 8.5 Human resources for health
- 9.1 Adult Mortality Chart 1
- 9.2 Adult Mortality Chart 2
- 9.3 Adult Mortality Chart 3
- 9.4 Adult Mortality Chart 4
- 10 Stress and Social Capital
- 10.1 Inability to control the important things in life - q8000
- 10.2 Inability to cope - q8001
- 10.3 Satisfaction with your health - q8002
- 10.4 Feeling safe when alone at home - q8011
- 10.5 Feeling safe when walking down the street alone after dark - q8012

1. Introduction

The World Health Organization initiated the World Health Survey (WHS) as a *data collection platform* to obtain information on the health of populations and on health systems. The WHS responds to the need of countries for a detailed and sustainable health information system and gathers data through surveys to measure essential population health parameters. The WHS uses standard survey procedures and instruments for general population surveys to gather comparable data across WHO Member States. These methods and instruments have been refined through scientific review of literature, extensive consultations with international experts and large-scale pilot tests. WHS is designed to evolve through its implementation by continuous input from collaborators, including policy makers, survey institutions, scientists and other interested parties. The data is jointly owned by the countries and the WHO, and there is a commitment for a long-term data collection, building local capacity and using this information for health policy.

The objectives of the World Health Survey were to:

- develop a **means** of providing:
 - **valid, relevant, reliable** and **comparable** information on **outcomes, functions** and **inputs** to health systems
 - to **supplement** the information provided by routine **HIS**
- build the **evidence base** necessary for policy-makers:
 - to **routinely monitor** if health systems are achieving the **desired goals**
 - and in some countries, to assess if the **increase in health resources** is having the desired effect

The introductory chapter of each country report will provide details on the current sources of data from routine health information systems and ad hoc surveys. It will describe the health situation in the country and identify gaps. It will also present a brief overview on the health system organization within the country.

2. Sampling

The WHO sampling guidelines emphasized scientific principles of survey sampling as explicit standards for quality, included examples of good sampling plans, and identified quality assurance standards for countries to adhere to. WHO and technical advisers provided technical support to countries when needed. The following principles were used for sampling in the WHS:

1. The sample should target the *de facto* population (i.e. all people living in that country including guest workers, immigrants and refugees) and not *de jure* population (i.e. the citizens of that country alone) with complete geographic coverage of the country, if possible, or a representative geographic segment of the country.
2. A *sample frame* with 90% coverage that included all key subgroups of interest. Enumeration or listing of households to update the frame was recommended.
3. The WHS sample identified all adult members of the general population aged 18 years or older as its sample. Households are selected using a random, stratified sampling procedure. One individual per household is then selected through a random selection procedure using the Kish table method.
4. The WHS used a sampling strategy which encompasses a known non-zero selection probability for any individual included in the study. This makes it possible to extrapolate the data to the whole population. Households and individuals are selected with a known probability in order to provide selection probabilities for each household and individual in the data set.
5. The WHS sampling strategy was without replacement and categories of non-respondents were clearly defined.
6. The WHS used GPS devices in certain countries and digitized maps to geocode the data within specified guidelines.

Each national report will present a detailed description of the sampling strategy and the allocation tables

2.2.1 Geographic Component of the Survey

Geographic Information refers to any information associated with a location on the Earth's surface. This location can be associated to a coordinate (latitude/longitude), a place's name, a postal address or any other geographic object.

Household Surveys represent an important source of information. If the geographic location of the surveyed clusters is collected at the same time that the interview is conducted users are then able to integrate the spatial dimension in their analysis

The geographic component of the WHO WHS is stored in two sections of the questionnaire:


- 0100 Sampling Information allowing the link between the location of each cluster and the sampling frame used for the survey. This section has also been used in order to collect the type of setting in which each household was located at the time of the survey. This section of the questionnaire has been used for all the countries part of the WHO WHS
- 0200 Geocoding Information for the collection of the latitude and longitude of each household. This section has only been used in 27 countries where GPS devices were available or have been provided. The choice of these countries has been done on the basis of the availability of digital maps delimiting the surveyed clusters, the field data collection planning and their distribution within the different WHO regions. Specific training and field material have been prepared and used by the interviewers or field supervisors in these countries in order to collect the latitude and longitude of each of the surveyed household.

As the sampling design may not always include the administrative structure of the country and as the codes entered in the data set for the sampling level needed to be labelled, a key correspondence table containing this information has been collected after the survey itself.

Once all the survey data and the key correspondence table available a specific protocol has been applied to the two sections mentioned earlier in order to clean eventual data collection or data entry mistakes as well as to fill the gaps in close collaboration with the survey institution.


The questionnaire as well as the training material and data cleaning protocols used can be downloaded from the WHO WHS web site (<http://www.who.int/healthinfo/survey/en/index.html>) in the "Survey Instrument and Related Documents" section.

2.3.1 Sample deviation index



SDI


2.3.2 HH Sample deviation index



HDI

2.3.3 Response rate

2.3.4 Missing data



2.3.5 Reliability

Insufficient / No data available

3.1 Household composition by residence and age groups according to income quintile

	Percent Rural	SE	Mean Number of									
			Household members	SE	Children < 2 years	SE	Children < 5years	SE	Children 5 -17 years	SE	Persons 18+ years	SE
Q1(poorest)	92.8	3.79	3.1	0.17	0.0	0.02	0.1	0.03	0.7	0.12	2.3	0.10
Q2	62.8	6.31	3.4	0.16	0.0	0.02	0.1	0.04	0.6	0.08	2.6	0.12
Q3	54.4	4.68	3.8	0.16	0.1	0.02	0.2	0.04	0.8	0.08	2.7	0.12
Q4	41.0	4.99	3.7	0.14	0.0	0.02	0.2	0.03	0.8	0.10	2.6	0.09
Q5(richest)	30.6	7.17	4.0	0.13	0.1	0.03	0.2	0.04	0.9	0.09	2.9	0.08
All	53.4	3.33	3.6	0.07	0.1	0.01	0.2	0.02	0.8	0.04	2.7	0.05

Note: table based on de jure members of household i.e. usual residents

3.2 Selected age distributions by income quintile

	Females Age 18–49	Persons Age 50+	Persons Age 60+
Q1 (poorest)	95	167	115
Q2	152	243	153
Q3	190	208	136
Q4	216	170	101
Q5 (richest)	250	185	75
All	903	973	581

Note: table based on de jure members of household i.e. usual residents

3.3 Characteristics of head of household by income quintile

	Mean Age of Head	SE	Median Education of Head	Percent Female Head	SE
Q1 (poorest)	50	1.85	3	31.5	5.00
Q2	53	1.58	3	22.5	3.88
Q3	48	1.18	4	22.7	3.73
Q4	47	0.98	4	18.3	2.99
Q5 (richest)	45	0.96	4	18.7	2.89
All	48	0.61	4	22.1	1.88

Education

1=no formal education

2=less than primary school

3=primary school completed

4=secondary school completed

5=high school (or equivalent) completed

6=college/pre-university/university completed

7=post graduate degree completed

Note: table based on de jure members of household i.e. usual residents

3.4 Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex

	Male	SE	Female	SE	Missing	Total	SE
Age							
0-4	5.5	0.60	5.2	0.60		5.40	0.40
5-14	17.8	1.20	15.8	1.10		16.80	0.70
15-29	21.8	1.10	21.8	1.10		21.80	0.60
30-44	23.4	1.00	23.1	0.80		23.30	0.70
45-59	17.3	1.00	17.0	1.00		17.10	0.80
60-69	9.0	1.10	10.4	0.80		9.70	0.70
70-79	4.8	0.70	5.5	0.60		5.20	0.50
80+	0.3	0.10	1.1	0.30		0.70	0.20
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Residence							
Urban	41.4	3.90	41.8	3.40		41.60	3.60
Rural	58.6	3.90	58.2	3.40		58.40	3.60
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Marital_Status							
Never	44.9	1.20	37.2	1.20		40.90	0.90
Currently	51.7	1.20	48.3	1.10		49.90	0.80
Separated	0.1	0.10	0.4	0.20		0.30	0.10
Divorced	0.3	0.10	2.2	0.40		1.30	0.20
Widowed	2.2	0.40	10.9	0.90		6.70	0.50
CoHabitng	0.3	0.20	0.4	0.20		0.40	0.20
Missing	0.5	0.10	0.5	0.20		0.50	0.10
Education							
NoFormal	14.1	1.10	20.8	1.40		17.60	0.80
LessthanPrimary	15.0	1.20	18.4	1.60		16.80	1.20
Primary	19.4	1.40	24.4	1.50		22.00	1.00
Secondary	44.5	1.90	30.5	1.80		37.20	1.60
HighSchool	3.9	0.60	3.1	0.50		3.50	0.40
College	2.4	0.50	2.3	0.50		2.40	0.50
PostGraduate	0.4	0.20	0.2	0.10		0.30	0.20
Missing	0.3	0.10	0.3	0.10		0.30	0.10
Total	100.0		100.0			100.00	
Number	1,701.0		1,810.0			3,511.00	
Number of Households						1,028.00	

3.5 Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education

	Male	SE	Female	SE	Missing	Total	SE
Age							
18-29	24.0	3.00	21.7	2.40		22.80	1.80
30-44	33.1	2.70	30.9	3.40		32.00	2.40
45-59	25.6	2.60	25.3	2.40		25.40	1.60
60-69	10.4	2.00	12.0	1.50		11.20	0.90
70-79	5.8	1.40	7.8	1.60		6.90	1.10
80+	1.1	0.60	2.2	1.10		1.60	0.60
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Residence							
Urban	43.9	4.30	45.2	4.20		44.50	3.60
Rural	56.1	4.30	54.8	4.20		55.50	3.60
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Marital_Status							
Never	24.8	3.00	15.0	1.80		19.80	1.70
Currently	70.5	3.00	64.2	3.00		67.30	2.10
Separated	0.6	0.60	0.1	0.10		0.40	0.30
Divorced	0.5	0.30	2.9	0.80		1.70	0.50
Widowed	3.4	1.00	16.9	1.90		10.30	1.10
CoHabiting	0.2	0.20	1.0	1.00		0.60	0.50
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Education							
NoFormal	2.9	1.40	14.7	2.80		9.00	1.70
LessthanPrimary	7.9	1.80	10.4	1.70		9.20	1.20
Primary	17.4	2.20	23.5	3.20		20.50	1.80
Secondary	61.6	3.20	44.3	3.50		52.70	2.80
HighSchool	6.2	1.50	3.8	1.10		5.00	0.80
College	3.8	1.00	3.4	1.10		3.60	0.90
PostGraduate	0.2	0.20	0.0	0.00		0.10	0.10
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Total	100.0		100.0			100.00	
Number	502.0		526.0			1,028.00	

4.1 Prevalence of smoking and average daily tobacco consumption

	Current daily smoker	SE	Smoker, not daily	SE	Non-smoker	SE	Missing	Average daily tobacco consumption*	SE	Number of respondents
Sex										
Male	46.4	4.10	7.6	3.40	45.6	3.60	0.4	23.4	0.70	502
Female	24.9	3.40	9.3	2.10	65.8	5.10	0.0	19.9	1.30	526
Missing										0
Residence										
Urban	40.3	4.20	11.7	4.00	47.5	5.80	0.4	22.2	1.40	458
Rural	31.5	3.10	5.9	1.40	62.7	3.20	0.0	22.0	0.90	570
Missing										0
Income Quintile										
Q1	37.1	5.20	3.9	1.30	59.0	5.50	0.0	23.8	1.30	137
Q2	37.8	4.20	7.9	2.20	54.1	5.00	0.2	21.0	1.30	208
Q3	33.1	4.20	5.2	2.90	61.7	5.40	0.0	20.9	1.80	206
Q4	36.5	4.70	6.7	2.60	56.0	5.30	0.7	20.7	0.90	223
Q5	33.7	6.50	15.1	5.80	51.3	5.70	0.0	24.6	1.70	252
Missing										3
Age										
18-29	36.0	4.20	11.3	4.60	52.7	5.80	0.0	18.9	1.10	235
30-44	39.0	3.80	9.2	2.20	51.7	4.50	0.0	21.6	1.00	329
45-59	39.4	4.90	9.8	4.40	50.1	4.80	0.6	26.4	1.70	262
60-69	29.4	4.30	2.9	1.80	67.4	4.30	0.3	22.0	1.80	116
70-79	18.5	6.00	1.7	1.00	79.7	6.20	0.0	15.2	2.50	70
80+	6.3	5.50	0.0	0.00	93.7	5.50	0.0	11.9	2.10	17
Missing										0
All	35.4	2.70	8.5	2.10	55.9	3.40	0.2	22.1	0.80	1,028

* Denominator used is current daily smoker

4.2 Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers

	Life time abstainers	SE	Non-heavy drinkers	SE	Infrequent Heavy Drinkers*	SE	Frequent Heavy drinkers**	SE	Missing	Number of respondents
Sex										
Male	53.6	4.90	43.9	4.80	1.8	0.90	0.3	0.30	0.4	502
Female	84.1	4.70	15.9	4.70	0.0	0.00	0.0	0.00	0.0	526
Missing										0
Residence										
Urban	59.5	5.00	39.5	5.00	0.6	0.30	0.0	0.00	0.4	458
Rural	77.0	3.40	21.6	3.30	1.1	0.80	0.3	0.30	0.0	570
Missing										0
Income Quintile										
Q1	79.8	4.40	18.9	4.50	0.3	0.20	1.0	1.10	0.0	137
Q2	69.2	8.80	30.4	8.80	0.2	0.00	0.0	0.00	0.2	208
Q3	80.7	4.10	18.9	4.10	0.5	0.30	0.0	0.00	0.0	206
Q4	70.6	4.40	27.8	4.10	0.8	0.60	0.0	0.00	0.7	223
Q5	53.6	5.80	44.1	5.70	2.3	1.70	0.0	0.00	0.0	252
Missing										3
Age										
18-29	68.4	5.40	31.2	5.40	0.4	0.30	0.0	0.00	0.0	235
30-44	70.2	4.50	28.5	4.40	1.3	1.10	0.0	0.00	0.0	329
45-59	62.6	4.90	35.5	4.80	0.7	0.50	0.5	0.50	0.6	262
60-69	71.3	6.60	26.5	6.80	1.9	1.80	0.0	0.00	0.3	116
70-79	83.7	5.30	16.3	5.30	0.0	0.00	0.0	0.00	0.0	70
80+	90.3	6.20	9.7	6.20	0.0	0.00	0.0	0.00	0.0	17
Missing										0
All	69.2	3.10	29.6	3.10	0.9	0.40	0.1	0.10	0.2	1,028

*Infrequent heavy drinkers: 1-2 days with 5 or more standard drinks per week(last 7 days)

**Frequent heavy drinkers: 3 or more days with 5 or more standard drinks per week(last 7 days)

4.3 Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

	Fruit and vegetable intake					Physical Activity					Number of respondents
	*Insufficient	SE	Sufficient	SE	Missing	**Insufficient	SE	Sufficient	SE	Missing	
Sex											
Male	84.2	2.70	15.8	2.70	0.0	8.9	1.80	91.1	1.80	0.0	502
Female	82.8	3.10	17.2	3.10	0.0	10.7	1.80	89.3	1.80	0.0	526
Missing											0
Residence											
Urban	84.2	4.30	15.8	4.30	0.0	9.5	1.80	90.5	1.80	0.0	458
Rural	82.9	2.70	17.1	2.70	0.0	10.2	2.30	89.8	2.30	0.0	570
Missing											0
Income Quintile											
Q1	76.1	6.40	23.9	6.40	0.0	11.7	2.60	88.3	2.60	0.0	137
Q2	93.4	1.90	6.6	1.90	0.0	14.0	3.90	86.0	3.90	0.0	208
Q3	83.8	2.80	16.2	2.80	0.0	6.1	2.90	93.9	2.90	0.0	206
Q4	86.3	2.60	13.7	2.60	0.0	14.6	3.20	85.4	3.20	0.0	223
Q5	76.7	5.90	23.3	5.90	0.0	4.4	1.80	95.6	1.80	0.0	252
Missing											3
Age											
18-29	80.5	4.20	19.5	4.20	0.0	6.6	2.80	93.4	2.80	0.0	235
30-44	85.3	2.60	14.7	2.60	0.0	2.5	0.90	97.5	0.90	0.0	329
45-59	81.5	3.90	18.5	3.90	0.0	8.9	2.30	91.1	2.30	0.0	262
60-69	80.9	4.50	19.1	4.50	0.0	12.8	2.90	87.2	2.90	0.0	116
70-79	95.4	2.70	4.6	2.70	0.0	40.7	6.90	59.3	6.90	0.0	70
80+	89.4	7.90	10.6	7.90	0.0	64.9	16.00	35.1	16.00	0.0	17
Missing											0
All	83.5	2.40	16.5	2.40	0.0	9.9	1.50	90.1	1.50	0.0	1,028

*Insufficient intake of fruit/vegetables: less than 5 servings of fruit/vegetables per one typical day

**Insufficient physical activity: less than 150 minutes per week(last 7 days) spent on walking/moderate activity/vigorous activity

4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water

	Piped into households	SE	other improved sources	SE	Total improved sources	SE	No improved sources	SE	Missing	Number of households
Residence										
Urban	95.9	1.60	3.5	1.40	99.3	0.50	0.7	0.50	0.0	479
Rural	79.9	3.60	13.2	2.80	93.1	2.10	6.9	2.10	0.0	549
Missing										0
Income_Quintile										
Q1(poorest)	67.1	5.60	19.4	4.40	86.5	4.40	13.5	4.40	0.0	157
Q2	83.4	4.30	10.5	3.30	93.9	2.10	6.1	2.10	0.0	208
Q3	88.6	3.10	10.0	3.10	98.6	0.60	1.4	0.60	0.0	202
Q4	92.0	1.60	6.0	1.80	98.1	1.30	1.9	1.30	0.0	223
Q5(richest)	98.6	0.70	1.4	0.70	100.0	0.00	0.0	0.00	0.0	234
Missing										3
Total	87.3	2.00	8.7	1.60	96.0	1.10	4.0	1.10	0.0	1,028

other improved sources: public standpipe/protected tube well/borehole/protected dug well/protected spring/rainwater collection

no improved sources: unprotected well/unprotected spring/water taken directly from a pond or stream/tanker truck water/vendor provided water

4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility

	Flush toilet to sewage system	SE	Other improved toilet facilities	SE	Total improved sanitation	SE	No improved sanitation	SE	Missing	Number of households
Residence										
Urban	85.5	5.50	13.9	5.20	99.3	0.50	0.7	0.50	0.0	479
Rural	49.0	4.00	48.0	4.10	97.0	1.60	3.0	1.60	0.0	549
Missing										0
Income_Quintile										
Q1(poorest)	20.8	5.20	69.7	5.40	90.5	4.10	9.5	4.10	0.0	157
Q2	52.7	5.90	46.1	5.70	98.9	1.10	1.1	1.10	0.0	208
Q3	70.3	5.20	28.7	5.30	99.0	1.00	1.0	1.00	0.0	202
Q4	82.8	4.10	17.1	4.10	99.8	0.20	0.2	0.20	0.0	223
Q5(richest)	87.7	3.60	12.3	3.60	100.0	0.00	0.0	0.00	0.0	234
Missing										3
Total	65.9	3.50	32.1	3.40	98.1	0.90	1.9	0.90	0.0	1,028

other improved sanitation facilities: connection to septic system/pour-flush latrine/covered dry latrine(with privacy) [provided facilities not shared]
no improved sanitation: uncovered dry latrine(without privacy)/bucket latrine/no facilities(open defecation)

4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used

	Electricity, Gas	SE	Kerosene	SE	Solid fuels on improved stove	SE	Solid fuels on traditional stove	SE	Total solid fuel use	SE	Missing	Number of households
Residence												
Urban	79.0	5.10	0.0	0.00	9.9	3.70	11.1	2.00	21.0	5.10	0.0	479
Rural	25.3	4.50	0.0	0.00	61.5	4.30	13.2	2.70	74.7	4.50	0.0	549
Missing												0
Income_ Quintile												
Q1(poorest)	2.8	1.20	0.0	0.00	76.4	5.00	20.9	5.00	97.2	1.20	0.0	157
Q2	25.8	3.90	0.0	0.00	54.3	6.60	20.0	4.60	74.2	3.90	0.0	208
Q3	48.3	6.10	0.0	0.00	39.0	4.90	12.7	3.70	51.7	6.10	0.0	202
Q4	63.9	6.40	0.0	0.00	28.0	5.00	8.1	3.90	36.1	6.40	0.0	223
Q5(richest)	92.1	3.40	0.0	0.00	4.5	1.50	3.4	2.20	7.9	3.40	0.0	234
Missing												3
Total	50.3	3.50	0.0	0.00	37.4	2.90	12.3	1.70	49.7	3.50	0.0	1,028

5.01 Tuberculosis

		Tuberculosis			
	Total	Need	Coverage	SE	M
Sex					
Female	526	4.1	22.8	7.8	0.0
Male	502	4.4	26.1	10.3	0.0
Missing	0				
Residence					
Urban	458	5.3	30.7	9.5	0.0
Rural	570	3.4	16.7	6.4	0.0
Missing	0				
Income_Quintile					
Q1(poorest)	137	10.3	20.8	8.2	0.0
Q2	208	2.7	10.0	8.5	0.0
Q3	206	2.8	27.7	22.0	0.0
Q4	223	5.1	37.9	18.3	0.0
Q5(richest)	252	2.5	20.3	10.4	0.0
Missing	3				
Age					
18-29	235	3.8	18.2	13.4	0.0
30-44	329	3.6	41.4	16.7	0.0
45-59	262	3.0	16.8	14.9	0.0
60-69	116	9.5	14.9	9.2	0.0
70-79	70	3.7	13.9	13.2	0.0
80+	17	8.4	58.8	34.3	0.0
Missing	0				
Total	1,028	4.2	24.5	6.1	0.0

M = % Missing of Covered SE= Standard Error of % Covered

Tuberculosis Indicators:

"Need" = % of all respondents with self-reported continuous cough lasting 3 weeks, OR had blood in phlegm, OR had coughed blood in the last 12 months

"Coverage" = % of respondents with self reported symptoms who indicated receiving a TB screening test in the last 12 months.

5.02 HIV Prevention and Screening

	Total (persons of reproductive age 18-49)	Voluntary Counselling and Testing for HIV/AIDS				Condom Usage			
		Need	Coverage	SE	M	Need	Coverage	SE	M
Sex									
Females	339	79.0	0.5	0.5	20.8	6.5	48.2	20.9	0.0
Males	344	86.1	0.0	0.0	37.5	23.8	57.3	8.5	0.9
Missing									
Residence									
Urban	289	83.1	0.6	0.6	15.5	20.5	49.4	8.2	0.9
Rural	394	82.2	0.0	0.0	40.0	11.3	63.1	13.0	0.4
Missing									
Income_Quintile									
Q1(poorest)	83	75.5	0.0	0.0	53.2	10.9	45.7	26.1	1.8
Q2	113	72.4	0.0	0.0	26.3	9.0	85.3	9.9	5.5
Q3	141	81.4	0.0	0.0	32.5	14.3	35.5	15.6	0.0
Q4	166	85.8	1.0	1.0	23.9	14.2	67.8	14.8	0.0
Q5(richest)	179	90.2	0.0	0.0	25.3	22.9	52.6	8.7	0.0
Missing	2								
Total	683	82.6	0.3	0.3	29.6	15.2	55.3	7.5	0.7

M = % Missing of Covered SE= Standard Error of % Covered

HIV Prevention and Screening Indicators:

"Total" = number of respondents of reproductive age.

"Voluntary Counselling and Testing for HIV/AIDS Need" = % of respondents of reproductive age who are sexually active, i.e. have had sexual intercourse in the last 12 months.

"Voluntary Counselling and Testing for HIV/AIDS Coverage" = % of respondents of reproductive age who are sexually active who indicated having taken an HIV test and receiving the results of test.

"Condom Usage Need" = % of respondents of reproductive age whose last sexual encounter during the last 12 months is classified as risky sexual behavior, i.e. having had sexual intercourse with a non-cohabitating partner .

"Condom Usage Coverage" = % of respondents of reproductive age who used a condom last time had risky sex during the last 12 months (with non-cohabitating partner).

5.03 Antenatal Care and Mother to Child Transmission of HIV

	Antenatal Care											Maternal to Child HIV Trans.			
	Women 18-49	Birth in last 5 years	Cov	SE	Visit 1x	Visit 2x	Visit 3x	Blood pressure checked	Blood sample taken	Told Signs of Pregnancy Complications	M	Birth in last 2 years	Cov	SE	M
Residence															
Urban	147	20.7	10.0	7.0	61.9	61.9	61.9	83.2	77.9	14.3	38.1	10.0	0.0	0.0	32.0
Rural	191	19.3	19.2	8.0	65.7	65.7	65.7	91.9	90.4	30.4	34.3	6.6	20.4	17.1	6.1
Missing															
Income_Quintile															
Q1(poorest)	43	12.4	32.6	19.3	64.9	64.9	64.9	71.0	71.0	32.6	35.1	3.0	0.0	0.0	0.0
Q2	67	18.9	27.5	16.5	92.8	92.8	92.8	98.3	93.0	27.5	7.2	4.3	0.0	0.0	7.9
Q3	74	23.2	18.3	14.2	92.5	92.5	92.5	92.5	92.5	18.3	7.5	17.3	20.3	18.5	0.0
Q4	82	19.5	11.3	6.0	35.7	35.7	35.7	71.0	65.1	18.1	64.3	7.5	0.0	0.0	55.4
Q5(richest)	72	22.9	0.0	0.0	39.4	39.4	39.4	97.3	94.0	26.6	60.6	6.0	0.0	0.0	42.8
Missing	2														
Total	339	19.9	15.0	5.3	64.0	64.0	64.0	88.0	84.8	23.1	36.0	8.1	9.4	8.9	20.0

M = % Missing of Covered SE= Standard Error of %Covered

Antenatal Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least 3 times or more by health care professional, received blood pressure, blood sample taken, and told signs about complications for their last pregnancy (aggregate antenatal care indicator).

"Visit 1x" "Visit 2x" "Visit 3x" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least once, twice or three times or more respectively by a health care professional during their last pregnancy.

"Blood pressure checked" = % of female respondents of reproductive age with live birth in the last 5 years who had blood pressure taken when they saw health care professional during their last pregnancy.

"Blood sample taken" = % of female respondents of reproductive age with live birth in the last 5 years who had a blood sample taken when they saw health care professional during their last pregnancy.

"Told Signs of Pregnancy Complications" = % of female respondents of reproductive age with live birth in the last 5 years who had been told signs of pregnancy complications when they saw health care professional during their last pregnancy.

Maternal to Child HIV Transmission Indicators:

"Birth in last 2 years" = % of female respondents of reproductive age indicating having had a live birth in the last 2 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 2 years who indicated having been counseled and tested and given results for HIV during their last pregnancy.

5.04 Birth Delivery Care

	Delivery Care					Place of Delivery						
	Women 18-49	Need	Coverage	SE	M	Public Health Facility	SE	Private Health Facility	SE	Home	SE	M
Residence												
Urban	147	20.7	83.2	10.8	16.8	83.2	10.8	0.0	0.0	0.0	0.0	0.0
Rural	191	19.3	97.9	1.6	2.1	97.9	1.6	0.0	0.0	0.0	0.0	0.0
Missing												
Income Quintile												
Q1(poorest)	43	12.4	96.5	3.6	3.5	96.5	3.6	0.0	0.0	0.0	0.0	0.0
Q2	67	18.9	97.9	2.2	2.1	97.9	2.2	0.0	0.0	0.0	0.0	0.0
Q3	74	23.2	92.5	7.5	7.5	92.5	7.5	0.0	0.0	0.0	0.0	0.0
Q4	82	19.5	76.7	14.4	23.3	76.7	14.4	0.0	0.0	0.0	0.0	0.0
Q5(richest)	72	22.9	97.3	2.8	2.7	97.3	2.8	0.0	0.0	0.0	0.0	0.0
Missing	2											
Total	339	19.9	91.3	4.9	8.7	91.3	4.9	0.0	0.0	0.0	0.0	0.0

M = % Missing of Covered SE= Standard Error of % Covered Birth Delivery Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Coverage" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having received assistance with birth delivery from a health care professional (doctor, nurse or midwife) during their last birth.

"Public Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a public health facility.

"Private Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a private health facility.

"Home" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered home or in the field (etc.)

5.05 Child's Health - Common Illness

	Total Children under Age of 5	Fever	More liquid	SE	Same or more food	SE	M	Cough	More liquid	SE	Same or more food	SE	M	Diarrhea	More liquid	SE	Same or more food	SE	M
Sex																			
Females	101	22.9	46.4	17.2	28.0	11.7	0.0	20.5	45.1	16.8	21.3	22.9	0.0	1.4	100.0	0.0	0.0	0.0	0.0
Males	101	37.4	44.5	13.4	32.4	13.8	1.2	22.3	42.8	17.1	44.1	37.4	2.0	10.1	53.6	23.8	2.3	2.5	4.5
Missing	0																		
Residence																			
Urban	63	46.6	58.0	14.9	28.8	18.4	0.0	34.5	47.6	17.7	39.0	46.6	0.0	11.4	73.6	21.4	3.3	3.7	0.0
Rural	139	22.9	33.5	15.2	32.4	10.6	1.4	15.6	40.3	14.9	27.5	22.9	2.1	3.2	36.0	25.0	0.0	0.0	10.3
Missing																			
Income_Quintile																			
Q1(poorest)	17	34.1	10.7	8.6	39.3	25.9	0.0	15.2	11.8	13.6	88.2	34.1	0.0	3.7	100.0	0.0	0.0	0.0	0.0
Q2	28	29.9	36.9	21.1	3.2	3.4	0.0	19.5	56.5	24.4	4.9	29.9	0.0	15.0	0.0	0.0	0.0	0.0	0.0
Q3	53	40.5	33.7	16.2	51.3	19.9	0.0	41.2	33.2	16.4	39.0	40.5	0.0	0.0	0.0	0.0	0.0	0.0	
Q4	50	15.4	56.2	20.7	37.8	19.7	6.0	8.4	64.4	24.7	24.5	15.4	11.0	2.9	68.1	30.7	0.0	0.0	31.9
Q5(richest)	54	32.6	70.0	14.2	12.9	8.1	0.0	16.9	61.7	18.8	24.8	32.6	0.0	9.7	100.0	0.0	4.4	5.5	0.0
Missing	0																		
Total	202	30.2	45.2	11.0	30.7	10.4	0.8	21.4	43.9	11.6	33.2	30.2	1.1	5.7	59.2	20.8	2.0	2.2	4.0

M = % Missing of Covered SE= Standard Error of % Covered

Child's Common Illness Indicators:

Total = Number of children under the age of 5 years at the time of the interview.

"Fever"/"Cough"/"Diarrhea" = % of children under the age of 5 with reported episode of (cough, fever, or diarrhea) in the last 3 months.

"More Liquid" = % of children under 5 with either fever, cough, or diarrhea in the last 3 months, that were given more liquids to drink during their illness.

"Same or More Food" = % of children under 5 with either fever, cough, or diarrhea in the last 3 months, that were given the same amount or more food to eat during their illness.

5.06 Child's Health - Danger Signs

	Children under 5	Fast Breath	Seeking Treatment						Vomit	Seeking Treatment						No Drink or Eat	Seeking Treatment					
			Yes	SE	Public	Private	Trad	M		Yes	SE	Public	Private	Trad	M		Yes	SE	Public	Private	Trad	M
Sex																						
Females	101	0.1	0.0	0.0	100.0	0.0	0.0	0.0	3.3	100.0	0.0	69.1	0.0	30.9	0.0	7.1	100.0	0.0	85.8	0.0	14.2	0.0
Males	101	0.1	4.7	5.1	92.2	7.8	0.0	0.0	4.6	71.0	25.4	76.6	23.4	0.0	0.0	12.2	89.1	10.8	100.0	0.0	0.0	0.0
Missing	0																					
Residence																						
Urban	63	0.1	0.0	0.0	87.0	13.0	0.0	0.0	6.4	100.0	0.0	55.4	19.3	25.3	0.0	12.4	100.0	0.0	86.8	0.0	13.2	0.0
Rural	139	0.1	4.5	4.9	100.0	0.0	0.0	0.0	2.8	65.6	30.2	100.0	0.0	0.0	0.0	8.5	88.5	11.0	100.0	0.0	0.0	0.0
Missing																						
Income_Quintile																						
Q1(poorest)	17	0.1	0.0	0.0	100.0	0.0	0.0	0.0	15.1	100.0	0.0	100.0	0.0	0.0	0.0	27.2	100.0	0.0	100.0	0.0	0.0	0.0
Q2	28	0.1	0.0	0.0	100.0	0.0	0.0	0.0	7.8	100.0	0.0	100.0	0.0	0.0	0.0	21.8	100.0	0.0	100.0	0.0	0.0	0.0
Q3	53	0.1	0.0	0.0	100.0	0.0	0.0	0.0	1.9	100.0	0.0	0.0	0.0	100.0	0.0	1.9	100.0	0.0	0.0	0.0	100.0	0.0
Q4	50	0.1	16.4	17.2	69.0	31.0	0.0	0.0	2.7	0.0	0.0			0.0	0.0	2.7	0.0	0.0		0.0	0.0	0.0
Q5(richest)	54	0.0	0.0	0.0	100.0	0.0	0.0	0.0	1.4	100.0	0.0	0.0	100.0	0.0	0.0	11.8	100.0	0.0	100.0	0.0	0.0	0.0
Missing	0																					
Total	202	0.1	2.9	3.1	95.3	4.7	0.0	0.0	3.9	83.0	15.9	72.9	11.7	15.4	0.0	9.7	93.1	6.8	94.4	0.0	5.6	0.0

M = % Missing of those Seeking Treatment SE= Standard Error of % Seeking Treatment

Child Danger Signs Indicators:

Total = Number of children under the age of 5 years at the time of the interview.

"Fast Breathing"/"Vomiting"/"No Drink or Eat" = % of children under the age of 5 with reported episodes of (fast breathing, vomiting, or no eating or drinking) in the last 3 months.

"Seeking Treatment" = % of children under 5 with episodes of fast breathing, vomiting, or no eating or drinking in the last 3 months that were taken for care/treatment at either a private health facility, public health facility, or traditional healer (other).

Note: large number of missings are usually due to not having information type of facility (either a no answer or a don't know answer).

5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card

	Children 12-23 months with card	DPT 3 Immunization			Measles		
		%Cov	SE	M	%Cov	SE	M
Sex							
Females	9	80.4	14.6	0.0	69.0	18.9	0.0
Males	16	33.1	17.6	0.0	55.4	23.5	0.0
Missing	0						
Residence							
Urban	10	23.7	23.7	0.0	23.7	23.7	0.0
Rural	15	67.8	17.2	0.0	84.3	9.0	0.0
Missing	0						
Income Quintile							
Q1(poorest)	1	54.3	30.5	0.0	69.6	26.4	0.0
Q2	5	89.5	10.9	0.0	89.5	10.9	0.0
Q3	10	36.3	31.6	0.0	36.3	31.6	0.0
Q4	5	78.4	21.0	0.0	60.5	13.3	0.0
Q5(richest)	5	15.0	16.1	0.0	76.1	23.6	0.0
Missing	0						
Total	26	50.4	17.2	0.0	60.4	17.6	0.0

M = % Missing of Covered SE= Standard Error of % Covered

Vaccinations Meeting WHO Timeline for Children 12 - 23 months Indicators:

Total = Children 12 - 23 months with a vaccination card seen by the interviewer (does not include children less than 12 months to account for censored observations).

"DPT 3" = % of children 12 - 23 months with a vaccination card seen by the interviewer who received 3 inoculations of DTP not sooner than 6, 10 and 14 weeks respectively and before 12 months of age, and that was verified by the interviewer as being recorded on the card.

"Measles" = % of children 12 - 23 months with a vaccination card seen by the interviewer who received who received 1 inoculation of measles not sooner than 8 months of age and before 12 months of age.

5.08 Vaccination Status of Children 12-23 Months

	Children 12-23 months	Card seen			DPT 3 Immunization							Measles						
		Seen	SE	M	Total		Card		No Card		M	Total		Card		No Card		M
					%Cov	SE	%Cov	SE	%Cov	SE		%Cov	SE	%Cov	SE	%Cov	SE	
Sex																		
Females	16	59.4	12.9	0.0	51.5	14.7	100.0	0.0	0.0	0.0	7.9	43.5	16.3	100.0	0.0	0.0	0.0	7.9
Males	29	56.1	15.8	0.0	56.1	15.8	100.0	0.0	0.0	0.0	26.8	61.2	15.5	91.6	8.4	8.4	8.4	21.7
Missing	0																	
Residence																		
Urban	19	53.4	19.5	0.0	46.9	20.4	100.0	0.0	0.0	0.0	32.0	46.9	20.4	100.0	0.0	0.0	0.0	32.0
Rural	26	60.1	12.1	0.0	60.1	12.1	100.0	0.0	0.0	0.0	11.5	61.0	11.9	90.5	9.0	9.5	9.0	5.7
Missing	0																	
Income_Quintile																		
Q1(poorest)	1	88.6	11.8	0.0	88.6	11.8	100.0	0.0	0.0	0.0	0.0	88.6	11.8	100.0	0.0	0.0	0.0	0.0
Q2	5	89.6	10.5	0.0	89.6	10.5	100.0	0.0	0.0	0.0	0.0	89.6	10.5	100.0	0.0	0.0	0.0	0.0
Q3	15	65.0	18.4	0.0	65.0	18.4	100.0	0.0	0.0	0.0	19.7	74.9	15.6	86.7	13.3	13.3	13.3	9.8
Q4	7	60.7	21.6	0.0	60.7	21.6	100.0	0.0	0.0	0.0	0.0	43.9	14.8	100.0	0.0	0.0	0.0	0.0
Q5(richest)	15	33.8	18.7	0.0	25.7	18.6	100.0	0.0	0.0	0.0	39.4	25.7	18.6	100.0	0.0	0.0	0.0	39.4
Missing	0																	
Total	45	57.3	10.8	0.0	54.5	11.2	100.0	0.0	0.0	0.0	20.2	55.0	11.2	93.9	6.0	6.1	6.0	16.8

M = % Missing (% without adequate information to estimate Total DPT3 or Measles Coverage)

SE= Standard Error of % Total, DPT3 or Measles Coverage

Explanation of Indicators for Vaccination Status of Children 12 - 23 Months:

"Card Seen" = % of children 12 - 23 months of age who had a vaccination card seen by the interviewer.

"DPT 3 Immunization": does not account for WHO timeline recommendations.

"Total" = % of all children 12 - 23 months of age with DPT3 vaccination either verified by card or by recall of respondent.

"Card" = % of all DPT3 vaccinations among children 12-23 months that were verified by card only.

"No Card" = % of all DPT3 vaccinations among children 12-23 months that were based on recall of the respondent only.

"Measles": does not account for WHO timeline recommendations.

"Total" = % of all children 12 - 23 months of age with measles vaccination either verified by card or by recall of respondent.

"Card" = % of all Measles vaccinations among children 12-23 months that were verified by card only.

"No Card" = % of all Measles vaccinations among children 12-23 months that were based on recall of the respondent only.

5.09 Women's Cancer Screening

	Total Females	Percent Females 18+	PAP Smear	SE	M	Percent Females 40+	Mammography / Breast Exam	SE	M
Residence									
Urban	237	92.0	35.9	5.2	4.9	56.0	18.6	4.1	2.8
Rural	288	88.4	29.4	2.7	11.4	42.2	3.6	1.7	6.0
Missing	0								
Income_Quintile									
Q1(poorest)	71	79.3	23.0	6.7	16.4	38.5	2.9	2.2	5.5
Q2	118	89.3	25.9	6.8	8.6	50.7	1.0	1.0	5.7
Q3	117	89.1	30.9	4.3	10.8	40.5	5.8	2.8	6.6
Q4	108	92.5	35.7	5.8	4.5	46.9	17.3	6.8	3.9
Q5(richest)	110	96.3	41.0	5.3	5.5	63.6	23.2	6.2	1.4
Missing	3								
Age									
18-29	114	21.7	39.6	5.9	3.5				
30-39	104	19.8	48.8	6.6	4.9				
40-49	120	22.8	30.0	5.7	7.6	22.8	13.4	4.6	1.6
50-59	71	13.5	18.9	3.9	5.6	13.5	10.6	3.3	3.8
60-69	63	12.0	12.5	2.4	27.5	12.0	8.5	4.4	10.1
70+	53	10.0		0.0		10.0	0.0	0.0	
Missing	0								
Total	526	90.0	32.4	2.7	8.4	48.4	11.4	2.3	4.3

M = % Missing of those with Pap Smear/Mammography SE= Standard Error of % with Pap Smear/Mammography

***Note: Values in the age row under the "Females 18 plus" and "Females 40 plus" columns are column percent values.

Women's Cancer Screening Indicators:

"Papsmear need"= % of all female respondents aged 18-69

"Papsmear" coverage = % of those female respondents aged 18-69 who self-reported receiving a pap smear test during pelvic examination in the last 3 years.

"Mammography need" = % of all female respondents aged 40-69

"Mammography" coverage = % of those female respondents aged 40-69 who self-reported receiving a mammography test or breast examination in the last 3 years.

****In several countries women above age 69 were asked the cancer screening questions, and for this reason they are included in the table. For countries that did not ask this questions of women above age 69, these values are expected to be missing.

5.10 Oral Health and Injuries

	Oral Health Problems					Road Traffic Accidents				Injury			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
Sex													
Female	526	29.9	76.9	5.7	0.0	1.8	65.2	22.3	0.0	4.8	58.7	15.8	0.0
Male	502	36.8	71.8	4.7	0.0	2.2	100.0	0.0	0.0	5.1	69.9	12.2	0.0
Missing	0												
Residence													
Urban	458	41.0	80.8	5.9	0.0	3.1	78.8	17.6	0.0	6.5	68.0	17.6	0.0
Rural	570	27.0	66.0	3.4	0.0	1.0	96.6	3.9	0.0	3.8	59.4	7.5	0.0
Missing	0												
Income_ Quintile													
Q1(poorest)	137	36.5	61.3	8.5	0.0	4.1	100.0	0.0	0.0	10.7	52.7	14.6	0.0
Q2	208	33.2	71.0	9.7	0.0	0.8	87.2	15.8	0.0	4.0	63.0	24.6	0.0
Q3	206	29.1	62.8	8.6	0.0	0.0	0.0	0.0	0.0	2.6	51.9	22.6	0.0
Q4	223	32.8	75.3	6.1	0.0	1.2	100.0	0.0	0.0	5.6	54.6	20.7	0.0
Q5(richest)	252	35.7	90.2	4.4	0.0	4.1	70.8	21.9	0.0	4.1	100.0	0.0	0.0
Missing	3												
Age													
18-29	235	38.5	84.1	5.6	0.0	3.2	100.0	0.0	0.0	7.0	81.9	16.2	0.0
30-44	329	34.5	75.9	6.6	0.0	3.4	72.5	19.4	0.0	1.4	13.3	14.2	0.0
45-59	262	35.5	75.5	7.3	0.0	0.5	100.0	0.0	0.0	7.1	86.1	8.6	0.0
60-69	116	22.2	59.6	9.9	0.0	0.2	0.0	0.0	0.0	8.6	16.2	11.6	0.0
70-79	70	25.5	32.3	10.2	0.0	0.0	0.0	0.0	0.0	2.1	82.7	16.0	0.0
80+	17	10.0	18.8	18.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Missing	0												
Total	1,028	33.3	74.1	4.0	0.0	2.0	83.9	12.4		5.0	64.4	11.2	

M = % Missing of % Covered SE= Standard Error of % Covered

Indicators:

Oral Health:

"Need" = % of all respondents indicating any problems in the last 12 months with their mouth and/or teeth.

"Coverage" = % of respondents reporting problems with their mouth and teeth who indicated receiving dental or medical care in last 12 months.

Road Traffic Accident:

"Need" = % of all respondents indicating having suffered bodily injury from a road traffic accident in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury from a road traffic accident in last 12 months who indicating having received treatment for their injuries on-site or at health facility within 1 hour of accident .

Injuries from Non-traffic Accidents:

"Need" = % of all respondents indicating having suffered bodily injury due to non-traffic accidents in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury due to non-traffic accidents in last 12 months who indicating having received treatment for injuries on-site or at health facility within 1 hour of accident

5.11 Self Reported Non-Communicable Conditions

	Total	Angina				Osteo-Arthritis				Asthma				Diabetes			
		Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
Sex																	
Female	526	8.7	85.2	6.7	0.0	17.8	55.9	7.7	1.1	3.1	66.5	20.5	0.0	5.2	86.8	7.0	0.8
Male	502	6.7	84.3	6.3	0.0	5.5	47.4	13.6	0.0	3.9	64.3	9.9	0.0	4.4	87.8	8.4	0.0
Missing	0																
Residence																	
Urban	458	7.5	76.1	10.0	0.0	12.6	59.2	10.6	1.7	4.5	52.5	13.7	0.0	5.8	87.1	8.4	0.0
Rural	570	8.0	91.4	3.8	0.0	11.2	49.2	5.9	0.0	2.7	82.4	9.3	0.0	4.0	87.3	7.6	1.0
Missing	0																
Income_Quintile																	
Q1(poorest)	137	13.1	89.4	6.4	0.0	13.5	29.0	7.2	0.0	4.6	72.8	12.0	0.0	9.5	88.5	7.9	1.8
Q2	208	7.5	92.9	6.7	0.0	11.0	70.2	10.1	0.0	3.2	98.3	1.8	0.0	2.6	88.3	7.5	0.0
Q3	206	10.9	84.3	9.3	0.0	14.8	62.8	15.1	0.0	4.7	74.4	14.8	0.0	6.2	83.7	10.9	0.0
Q4	223	4.4	83.9	10.6	0.0	12.0	34.8	12.3	3.7	4.2	33.9	20.7	0.0	4.6	86.7	7.1	0.0
Q5(richest)	252	5.3	70.3	16.4	0.0	8.1	73.0	10.4	0.0	1.5	49.2	31.0	0.0	3.1	90.7	10.2	0.0
Missing	3																
Age																	
18-29	235	0.2	100.0	0.0	0.0	5.8	4.8	5.2	0.0	0.5	39.8	33.9	0.0	0.2	90.5	8.3	0.0
30-44	329	1.7	83.4	14.2	0.0	5.5	61.8	16.4	0.0	2.0	21.4	20.6	0.0	2.3	98.1	1.7	0.0
45-59	262	8.4	73.2	11.9	0.0	12.0	44.6	11.0	3.1	1.5	74.5	22.2	0.0	4.7	82.5	9.0	2.7
60-69	116	19.1	84.5	7.9	0.0	32.0	59.9	8.9	0.0	12.2	61.0	11.8	0.0	17.6	79.1	11.5	
70-79	70	30.3	92.3	5.5	0.0	29.5	83.9	6.4	0.0	13.4	97.8	1.7	0.0	12.0	0.0	0.0	
80+	17	51.5	96.4	4.4	0.0	2.7	0.0	0.0	0.0	5.0	100.0	0.0	0.0	0.0	0.0	0.0	
Missing	0																
Total	1,028	7.8	84.8	5.1	0.0	11.8	53.9	5.9	0.8	3.5	65.3	10.3	0.0	4.8	87.2	5.7	0.5

M = % Missing for Covered

SE= Standard Error of %Covered

Self Reported Non-Communicable Conditions Indicators:

"Need" = % of total respondents indicating having received a diagnosis of the condition (angina, arthritis, asthma, diabetes) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (angina, arthritis, asthma, diabetes) indicating they have received treatment or screening for the condition in the last 2 weeks.

5.12 Mental Health Conditions

	Total	Depression				Schizophrenia			
		Need	Coverage	SE	M	Need	Coverage	SE	M
Sex									
Female	526	9.1	50.5	12.9	0.0	0.1	0.0	0.0	0.0
Male	502	3.4	46.5	17.3	0.0	0.1	100.0	0.0	0.0
Missing	0								
Residence									
Urban	458	7.1	41.3	17.5	0.0	0.1	100.0	0.0	0.0
Rural	570	5.7	57.5	11.2	0.0	0.1	26.1	27.3	0.0
Missing	0								
Income Quintile									
Q1(poorest)	137	9.3	51.3	13.5	0.0	0.1	100.0	0.0	0.0
Q2	208	6.1	86.9	7.4	0.0	0.1	0.0	0.0	0.0
Q3	206	6.6	73.4	15.4	0.0	0.0	0.0	0.0	0.0
Q4	223	10.0	20.6	15.7	0.0	0.3	100.0	0.0	0.0
Q5(richest)	252	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Missing	3								
Age									
18-29	235	1.3	53.6	34.0	0.0	0.0	100.0	0.0	0.0
30-44	329	7.5	34.9	17.2	0.0	0.0	0.0	0.0	0.0
45-59	262	3.8	42.8	14.8	0.0	0.3	67.1	31.2	0.0
60-69	116	8.3	50.5	19.7	0.0	0.0	0.0	0.0	0.0
70-79	70	17.2	59.3	20.8	0.0	0.0	0.0	0.0	
80+	17	33.5	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Missing	0								
Total	1,028	6.3	49.5	10.8	0.0	0.1	70.5		0.0

M = % Missing for Covered

SE= Standard Error of %Covered

Mental Health Conditions Indicators:

"Need" = % of total respondents indicating having received a diagnosis of the condition (depression, schizophrenia) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (depression, schizophrenia) indicating they have received treatment or screening for the condition in the last 2 weeks.

5.13 Cataracts

		Cataracts			
	Total 60+	Need	Coverage	SE	M
Sex					
Female	116	10.6	24.1	15.8	0.0
Male	87	13.7	53.5	21.8	0.0
Missing	0				
Residence					
Urban	92	18.5	54.7	18.6	0.0
Rural	111	6.4	0.0	0.0	0.0
Missing	0				
Income_Quintile					
Q1(poorest)	39	10.1	0.0	0.0	0.0
Q2	58	1.9	10.4	11.5	0.0
Q3	46	13.9	20.4	18.0	0.0
Q4	36	23.3	50.5	22.8	0.0
Q5(richest)	24	16.2	96.3	4.2	0.0
Missing	1				
Age					
60-69	116	8.1	42.2	19.9	0.0
70-79	70	15.6	14.2	11.7	0.0
80+	17	22.8	100.0	0.0	0.0
Missing	0				
Total	203	11.9	38.6	11.8	0.0

M = % Missing of Covered SE= Standard Error of %Covered

Cataracts Indicators:

Total = Number of respondents 60 years of age or more.

"Need" = % in need (respondents 60 years of age or more with self-reported diagnosis of cataracts in the last 5 years).

"Coverage" = % respondents 60 years of age or more with self-reported diagnosis of cataracts having indicated receiving eye surgery to remove cataracts in the last 5 years)

6.01 In general, rate your health today - q2000

	Very Good	SE	Good	SE	Moderate	SE	Bad	SE	Very Bad	SE	Missing	SE	N
Gender													
Male	31.6	4.3	36.5	4.3	24.0	2.5	6.1	1.3	1.8	0.7	0.0	0.0	502
Female	26.5	2.7	33.0	3.2	26.1	2.3	13.0	2.6	1.3	0.4	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	26.3	4.7	39.9	4.4	24.7	3.4	8.3	2.1	0.8	0.5	0.0	0.0	458
Rural	31.1	3.6	30.6	3.2	25.3	2.2	10.7	2.0	2.2	0.7	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	21.3	3.2	29.7	5.9	24.3	4.3	19.4	4.3	5.4	1.7	0.0	0.0	137
Q2	23.0	3.6	31.7	5.2	33.1	4.1	9.9	2.6	2.2	1.3	0.0	0.0	208
Q3	33.1	3.8	31.8	5.3	23.4	4.9	10.6	3.2	1.1	0.8	0.0	0.0	206
Q4	34.9	5.4	30.3	4.4	27.9	4.0	6.0	1.8	0.9	0.9	0.0	0.0	223
Q5(richest)	29.8	6.1	45.7	4.5	17.8	3.4	6.7	2.8	0.0	0.0	0.0	0.0	252
M	0.0	0.0	80.0	20.6	20.0	20.6	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	52.2	4.9	37.8	5.4	9.3	3.2	0.5	0.4	0.0	0.0	0.0	0.0	235
30-44	35.8	3.9	37.5	3.7	22.0	2.9	4.6	1.6	0.1	0.1	0.0	0.0	329
45-59	20.3	5.5	36.2	4.2	32.2	3.8	8.7	2.4	2.5	1.1	0.0	0.0	262
60-69	3.7	1.9	29.5	6.1	38.2	5.6	22.5	3.6	6.0	2.7	0.0	0.0	116
70-79	0.5	0.4	21.2	8.0	40.4	9.9	35.0	8.1	2.9	1.7	0.0	0.0	70
80+	0.0	0.0	5.2	5.4	38.9	17.6	55.9	18.7	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	29.0	2.9	34.7	2.7	25.1	1.9	9.7	1.5	1.6	0.4			1,028

6.02 Difficulty with 'work and household activities' in last 30 days - q2001

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	46.7	4.1	26.6	3.7	18.3	2.2	5.6	1.1	2.7	1.2	0.0	0.0	502
Female	43.6	3.1	20.8	2.0	24.5	1.9	7.9	2.0	3.1	1.3	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	47.5	3.9	21.4	2.6	23.0	2.7	7.3	2.0	0.8	0.4	0.0	0.0	458
Rural	43.2	3.9	25.5	2.9	20.3	1.6	6.4	1.2	4.7	2.0	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	32.7	3.3	20.2	4.7	29.1	3.7	11.5	2.9	6.5	1.5	0.0	0.0	137
Q2	39.1	3.8	20.5	3.2	28.1	3.0	8.5	2.5	3.9	2.6	0.0	0.0	208
Q3	50.0	5.5	18.1	3.3	21.5	4.2	6.3	1.7	4.2	2.9	0.0	0.0	206
Q4	46.6	4.2	26.1	3.8	19.9	3.7	7.2	2.2	0.2	0.2	0.0	0.0	223
Q5(richest)	51.4	5.7	30.8	5.3	13.2	3.5	2.9	1.8	1.7	1.3	0.0	0.0	252
M	62.4	28.8	0.0	0.0	37.6	28.8	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	79.6	4.6	13.3	4.4	4.7	1.2	0.0	0.0	2.3	2.2	0.0	0.0	235
30-44	50.6	4.2	29.6	3.5	17.9	2.5	1.8	0.7	0.1	0.1	0.0	0.0	329
45-59	36.1	5.6	27.2	4.4	26.9	4.1	7.3	2.8	2.4	1.0	0.0	0.0	262
60-69	11.2	2.9	20.3	3.4	44.7	4.2	19.6	3.7	4.2	1.4	0.0	0.0	116
70-79	3.7	2.8	25.4	7.1	35.1	10.2	25.3	6.2	10.6	4.7	0.0	0.0	70
80+	5.2	5.4	11.9	8.0	24.9	13.6	24.6	15.8	33.5	23.6	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	45.2	2.8	23.6	2.0	21.5	1.5	6.8	1.1	2.9	1.1			1,028

6.03 Difficulty with 'moving around' in last 30 days - q2010

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	61.2	3.0	18.8	2.2	12.5	1.9	5.2	1.2	2.3	1.2	0.0	0.0	502
Female	55.0	3.1	19.9	1.7	14.4	2.3	9.9	2.4	0.9	0.3	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	54.8	3.2	20.3	2.5	17.2	3.3	7.0	2.1	0.6	0.4	0.0	0.0	458
Rural	60.6	3.4	18.6	1.9	10.4	1.2	8.1	1.6	2.4	1.0	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	42.9	5.8	23.7	4.4	15.9	3.7	13.4	3.2	4.1	1.3	0.0	0.0	137
Q2	53.2	3.4	21.5	3.4	13.7	2.1	8.0	2.4	3.6	2.5	0.0	0.0	208
Q3	60.9	5.1	13.9	2.5	15.4	4.0	8.6	3.3	1.1	0.6	0.0	0.0	206
Q4	57.5	4.6	22.1	3.5	15.7	3.8	4.7	2.3	0.0	0.0	0.0	0.0	223
Q5(richest)	68.0	3.9	17.4	3.0	8.3	1.9	5.9	2.3	0.4	0.4	0.0	0.0	252
M	80.0	20.6	0.0	0.0	20.0	20.6	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	85.8	4.1	7.3	2.4	4.6	1.7	0.0	0.0	2.3	2.2	0.0	0.0	235
30-44	71.5	3.5	15.9	2.6	11.8	3.0	0.5	0.3	0.3	0.2	0.0	0.0	329
45-59	50.3	5.0	29.0	3.8	12.7	2.5	6.8	2.7	1.2	0.5	0.0	0.0	262
60-69	20.8	4.2	23.0	3.9	24.7	4.4	28.0	5.2	3.5	1.5	0.0	0.0	116
70-79	4.7	2.4	34.5	9.1	33.6	9.9	23.2	8.0	4.0	1.8	0.0	0.0	70
80+	5.2	5.4	15.7	9.5	19.8	10.1	59.3	17.7	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	58.0	2.3	19.3	1.5	13.5	1.5	7.6	1.3	1.6	0.6			1,028

6.04 Difficulty with 'vigorous activities' in last 30 days - q2011

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	57.0	4.0	16.0	2.5	9.7	1.8	8.4	1.6	8.9	1.7	0.0	0.0	502
Female	51.7	3.4	14.5	1.7	11.5	2.0	9.5	1.6	12.8	2.4	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	54.5	4.4	13.6	2.1	13.6	3.0	8.5	1.8	9.8	2.1	0.0	0.0	458
Rural	54.1	3.9	16.6	1.8	8.2	1.6	9.3	1.7	11.7	2.2	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	42.0	5.3	16.4	3.5	12.6	3.9	10.3	3.4	18.8	3.2	0.0	0.0	137
Q2	47.3	4.6	18.6	3.0	11.8	3.7	8.3	2.8	14.1	3.2	0.0	0.0	208
Q3	56.0	5.5	9.4	2.2	11.0	3.3	13.6	3.3	10.0	3.3	0.0	0.0	206
Q4	55.4	5.1	17.6	4.2	9.0	2.5	9.0	2.4	9.0	2.8	0.0	0.0	223
Q5(richest)	64.1	5.0	14.6	4.4	9.8	2.7	4.9	1.8	6.5	2.3	0.0	0.0	252
M	80.0	20.6	0.0	0.0	0.0	0.0	20.0	20.6	0.0	0.0	0.0	0.0	3
Age													
18-29	85.0	3.7	6.4	1.8	5.1	1.9	1.2	0.4	2.3	2.2	0.0	0.0	235
30-44	59.2	4.2	21.0	4.2	10.8	3.3	8.3	2.5	0.7	0.2	0.0	0.0	329
45-59	48.5	4.6	19.0	3.0	9.2	1.8	13.2	2.8	10.2	3.0	0.0	0.0	262
60-69	27.5	6.6	11.4	2.9	13.4	3.8	15.7	3.5	32.1	4.7	0.0	0.0	116
70-79	6.1	3.4	13.6	4.4	27.5	9.9	11.7	3.7	41.1	8.4	0.0	0.0	70
80+	5.2	5.4	0.0	0.0	17.1	9.6	8.6	5.5	69.1	14.8	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	54.3	2.9	15.2	1.4	10.6	1.6	9.0	1.2	10.9	1.5			1,028

6.05 Difficulty with 'self-care' in last 30 days - q2020

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	81.2	2.7	10.4	2.0	5.0	1.5	2.5	1.2	0.9	0.4	0.0	0.0	502
Female	76.6	3.8	13.4	2.2	8.0	2.3	1.9	0.5	0.2	0.2	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	83.9	4.5	8.1	1.9	7.3	3.0	0.4	0.2	0.3	0.3	0.0	0.0	458
Rural	74.7	2.5	15.0	1.8	5.9	1.3	3.6	1.0	0.8	0.2	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	60.5	5.6	21.4	4.4	8.4	2.2	6.4	2.1	3.2	1.0	0.0	0.0	137
Q2	72.8	5.1	13.7	3.3	7.9	2.9	5.5	2.6	0.1	0.1	0.0	0.0	208
Q3	75.0	5.2	13.8	2.8	10.2	4.1	0.5	0.4	0.5	0.5	0.0	0.0	206
Q4	83.6	3.8	11.2	3.1	5.2	1.6	0.0	0.0	0.0	0.0	0.0	0.0	223
Q5(richest)	92.3	2.1	4.6	1.8	2.6	1.7	0.5	0.4	0.0	0.0	0.0	0.0	252
M	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	93.2	2.7	3.1	1.2	1.3	1.3	2.3	2.2	0.0	0.0	0.0	0.0	235
30-44	88.3	2.3	9.8	2.3	1.5	0.8	0.3	0.2	0.1	0.1	0.0	0.0	329
45-59	79.2	3.3	12.7	2.3	5.5	2.6	1.9	1.0	0.8	0.3	0.0	0.0	262
60-69	54.1	6.2	20.6	4.4	17.6	4.2	5.2	1.8	2.5	1.3	0.0	0.0	116
70-79	39.8	9.3	27.4	8.0	25.7	9.8	6.8	1.4	0.2	0.2	0.0	0.0	70
80+	19.8	10.6	38.6	17.7	38.1	22.2	3.5	3.6	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	78.8	2.5	11.9	1.4	6.5	1.5	2.2	0.6	0.6	0.2			1,028

6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	81.2	2.7	11.2	2.1	4.3	1.4	2.3	1.2	1.1	0.4	0.0	0.0	502
Female	76.9	3.5	13.8	2.2	7.2	2.3	1.8	0.5	0.2	0.2	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	84.3	4.2	8.6	2.0	6.5	3.0	0.1	0.1	0.5	0.4	0.0	0.0	458
Rural	74.8	2.5	15.7	1.8	5.2	1.3	3.6	1.0	0.8	0.2	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	61.0	5.6	21.1	4.4	8.4	2.2	6.2	2.1	3.2	1.0	0.0	0.0	137
Q2	73.4	5.0	13.7	3.3	7.3	2.9	5.5	2.6	0.1	0.1	0.0	0.0	208
Q3	74.9	5.2	14.7	2.9	9.5	4.1	0.5	0.4	0.5	0.5	0.0	0.0	206
Q4	83.9	3.4	13.1	2.9	3.0	1.4	0.0	0.0	0.0	0.0	0.0	0.0	223
Q5(richest)	92.3	2.1	4.7	1.8	2.6	1.7	0.0	0.0	0.4	0.4	0.0	0.0	252
M	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	93.3	2.7	4.4	1.7	0.0	0.0	2.3	2.2	0.0	0.0	0.0	0.0	235
30-44	89.5	2.2	9.0	2.1	1.2	0.8	0.3	0.2	0.1	0.1	0.0	0.0	329
45-59	79.1	3.3	13.2	2.3	5.1	2.6	1.3	0.9	1.2	0.5	0.0	0.0	262
60-69	54.6	6.2	22.4	4.5	15.6	4.4	5.0	1.8	2.5	1.3	0.0	0.0	116
70-79	35.3	8.8	32.4	8.5	25.2	9.8	6.8	1.4	0.2	0.2	0.0	0.0	70
80+	25.4	12.7	33.0	16.7	38.1	22.2	3.5	3.6	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	79.0	2.4	12.5	1.4	5.8	1.5	2.0	0.6	0.7	0.2			1,028

6.07 Amount of 'bodily aches or pains' in last 30 days - q2030

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	61.0	3.7	17.4	2.4	16.1	3.9	4.3	1.0	1.3	0.5	0.0	0.0	502
Female	55.6	3.0	24.5	1.8	9.9	1.9	8.3	2.1	1.7	0.9	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	55.4	3.9	23.3	2.7	14.0	3.2	5.1	1.2	2.3	1.1	0.0	0.0	458
Rural	60.5	3.3	19.2	1.9	12.0	3.2	7.4	1.4	0.8	0.4	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	48.3	6.1	23.0	4.0	12.5	2.9	13.8	2.9	2.5	1.3	0.0	0.0	137
Q2	57.7	3.8	23.8	4.2	11.6	2.9	6.2	2.1	0.7	0.6	0.0	0.0	208
Q3	56.5	4.8	17.7	3.9	16.2	4.2	8.7	3.2	0.9	0.6	0.0	0.0	206
Q4	61.3	4.0	22.3	3.0	12.0	2.8	3.8	1.4	0.6	0.4	0.0	0.0	223
Q5(richest)	63.2	6.5	18.8	4.1	12.2	6.0	2.9	1.5	2.8	1.9	0.0	0.0	252
M	17.6	18.5	62.4	28.8	20.0	20.6	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	88.9	3.3	7.5	2.0	3.3	1.7	0.0	0.0	0.3	0.2	0.0	0.0	235
30-44	68.5	3.8	15.3	2.9	14.6	4.1	1.3	0.6	0.3	0.3	0.0	0.0	329
45-59	51.7	5.5	29.7	3.9	10.5	3.6	5.0	1.4	3.1	2.0	0.0	0.0	262
60-69	17.3	3.8	38.6	6.1	18.4	3.1	23.2	4.9	2.6	1.3	0.0	0.0	116
70-79	9.5	4.5	29.6	7.3	37.6	8.7	19.7	7.5	3.5	2.0	0.0	0.0	70
80+	15.6	9.9	31.5	16.6	8.6	5.5	44.3	21.4	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	58.2	2.5	21.0	1.6	12.9	2.3	6.4	1.0	1.5	0.6			1,028

6.08 Amount of 'bodily discomfort' in last 30 days - q2031

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	56.5	3.8	23.9	3.1	14.4	2.4	4.4	1.0	0.7	0.4	0.0	0.0	502
Female	52.4	2.7	27.3	2.0	10.5	1.8	7.8	1.9	2.0	1.0	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	52.9	3.8	26.3	2.9	13.5	2.5	5.3	1.4	2.1	1.1	0.0	0.0	458
Rural	55.7	3.7	25.1	1.9	11.5	2.2	6.9	1.3	0.9	0.4	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	45.3	5.9	25.1	4.0	14.9	3.5	12.0	2.2	2.6	1.3	0.0	0.0	137
Q2	50.6	4.0	27.2	4.0	15.4	4.0	6.1	2.1	0.7	0.6	0.0	0.0	208
Q3	53.1	4.1	23.6	4.7	14.5	3.3	7.3	3.2	1.5	0.8	0.0	0.0	206
Q4	56.0	4.5	26.2	3.6	11.8	2.9	5.6	1.7	0.3	0.3	0.0	0.0	223
Q5(richest)	62.1	6.9	25.9	4.7	7.1	2.8	2.7	1.5	2.1	1.8	0.0	0.0	252
M	62.4	28.8	17.6	18.5	20.0	20.6	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	81.1	4.0	13.1	2.8	5.5	3.3	0.0	0.0	0.3	0.2	0.0	0.0	235
30-44	63.9	4.2	23.7	3.9	10.8	3.1	1.4	0.7	0.1	0.1	0.0	0.0	329
45-59	50.1	5.4	33.2	4.0	8.7	2.0	5.1	1.5	2.9	1.9	0.0	0.0	262
60-69	17.3	3.6	34.6	4.3	21.8	3.6	23.0	5.4	3.4	1.4	0.0	0.0	116
70-79	7.3	3.7	32.2	7.9	40.6	8.2	17.1	4.8	2.8	1.9	0.0	0.0	70
80+	15.6	9.9	31.5	16.6	13.2	7.7	39.6	22.3	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	54.4	2.6	25.6	1.7	12.4	1.6	6.2	0.9	1.4	0.5			1,028

6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	77.6	2.5	12.1	1.6	7.9	1.6	2.0	0.7	0.3	0.3	0.0	0.0	502
Female	63.1	3.4	22.8	2.2	9.3	1.8	4.7	1.4	0.2	0.1	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	66.3	3.9	19.7	2.0	11.1	2.1	2.8	0.8	0.1	0.1	0.0	0.0	458
Rural	73.4	2.4	15.9	2.0	6.6	1.2	3.8	1.1	0.4	0.3	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	62.4	4.2	21.5	3.4	9.5	2.3	5.1	1.2	1.5	1.2	0.0	0.0	137
Q2	66.7	4.3	18.7	2.8	11.0	2.7	3.4	1.6	0.2	0.2	0.0	0.0	208
Q3	74.4	4.9	14.3	2.8	6.4	2.4	4.9	2.9	0.0	0.0	0.0	0.0	206
Q4	65.8	4.0	22.1	3.2	9.1	2.0	3.0	1.3	0.0	0.0	0.0	0.0	223
Q5(richest)	77.8	4.0	13.3	3.0	7.5	2.5	1.4	1.2	0.0	0.0	0.0	0.0	252
M	80.0	20.6	0.0	0.0	20.0	20.6	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	89.6	3.3	8.4	3.2	1.3	1.0	0.7	0.7	0.0	0.0	0.0	0.0	235
30-44	86.4	2.5	9.6	2.2	2.8	1.0	1.2	0.6	0.0	0.0	0.0	0.0	329
45-59	68.6	4.3	21.3	3.5	7.5	2.5	2.1	1.3	0.5	0.6	0.0	0.0	262
60-69	29.8	4.2	32.6	3.9	30.0	5.6	7.7	2.2	0.0	0.0	0.0	0.0	116
70-79	18.8	6.1	42.5	8.8	27.1	7.8	10.3	2.9	1.3	0.8	0.0	0.0	70
80+	2.1	2.2	37.6	17.3	17.7	9.6	42.6	21.7	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	70.2	2.2	17.6	1.5	8.6	1.2	3.4	0.7	0.2	0.2			1,028

6.10 Difficulty with 'learning a new task' in last 30 days - q2051

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	78.5	2.6	11.2	1.8	7.4	1.6	2.5	0.9	0.3	0.3	0.0	0.0	502
Female	65.9	2.5	20.3	1.8	8.4	2.0	4.1	1.2	1.3	1.0	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	69.5	3.2	17.8	2.6	8.9	1.9	3.7	1.0	0.1	0.1	0.0	0.0	458
Rural	74.2	2.0	14.2	1.7	7.2	1.5	3.0	0.7	1.5	1.0	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	64.3	4.5	17.7	3.5	11.1	2.6	5.0	1.1	1.9	1.2	0.0	0.0	137
Q2	60.7	5.8	23.1	4.4	11.6	3.0	4.4	1.7	0.2	0.2	0.0	0.0	208
Q3	77.9	5.0	11.7	2.5	5.8	2.3	1.9	1.1	2.7	2.7	0.0	0.0	206
Q4	71.5	4.1	18.0	3.6	5.4	1.5	5.2	2.1	0.0	0.0	0.0	0.0	223
Q5(richest)	81.3	3.5	10.5	3.2	7.1	2.3	1.1	1.1	0.0	0.0	0.0	0.0	252
M	80.0	20.6	0.0	0.0	20.0	20.6	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	94.1	2.3	3.1	1.4	0.8	0.8	2.0	1.4	0.0	0.0	0.0	0.0	235
30-44	88.2	2.3	9.6	2.1	1.8	0.9	0.4	0.4	0.0	0.0	0.0	0.0	329
45-59	71.4	4.9	19.7	4.5	6.4	2.2	2.0	1.3	0.5	0.6	0.0	0.0	262
60-69	30.8	4.3	34.3	4.3	25.8	6.2	9.1	2.7	0.0	0.0	0.0	0.0	116
70-79	10.6	3.9	37.4	8.4	34.6	9.0	16.1	5.3	1.3	0.8	0.0	0.0	70
80+	2.1	2.2	37.6	17.3	15.9	9.2	7.5	6.3	37.0	22.8	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	72.1	1.8	15.8	1.5	7.9	1.2	3.3	0.6	0.8	0.6			1,028

6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	86.4	2.1	8.6	1.8	3.8	1.5	0.8	0.5	0.4	0.4	0.0	0.0	502
Female	80.1	1.8	12.2	1.7	5.5	1.7	2.2	0.8	0.0	0.0	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	84.0	2.3	10.1	2.2	4.1	1.6	1.8	0.8	0.0	0.0	0.0	0.0	458
Rural	82.6	1.7	10.7	1.8	5.1	1.3	1.3	0.5	0.4	0.3	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	71.7	4.5	14.6	4.8	10.4	2.6	1.7	1.2	1.5	1.4	0.0	0.0	137
Q2	77.9	3.9	14.0	2.6	5.3	2.3	2.8	1.5	0.0	0.0	0.0	0.0	208
Q3	80.3	3.6	12.6	3.2	6.5	3.4	0.6	0.4	0.0	0.0	0.0	0.0	206
Q4	87.0	2.9	8.0	2.4	2.2	1.4	2.8	1.4	0.0	0.0	0.0	0.0	223
Q5(richest)	92.6	2.5	5.7	2.0	1.7	1.1	0.0	0.0	0.0	0.0	0.0	0.0	252
M	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	90.3	3.2	7.5	2.9	1.5	1.5	0.7	0.7	0.0	0.0	0.0	0.0	235
30-44	91.4	2.3	5.1	1.5	2.3	1.2	1.2	0.9	0.0	0.0	0.0	0.0	329
45-59	84.5	2.6	9.5	1.9	4.4	1.7	0.9	0.7	0.7	0.7	0.0	0.0	262
60-69	66.0	4.6	24.6	5.3	6.9	3.0	2.4	1.6	0.0	0.0	0.0	0.0	116
70-79	56.1	9.9	21.7	6.3	15.6	7.5	6.5	3.1	0.1	0.1	0.0	0.0	70
80+	36.6	17.7	26.4	13.9	37.0	22.8	0.0	0.0	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	83.2	1.4	10.4	1.4	4.6	1.0	1.5	0.5	0.2	0.2			1,028

6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	84.6	2.0	11.2	1.8	3.8	1.5	0.4	0.3	0.0	0.0	0.0	0.0	502
Female	78.2	3.2	16.3	2.6	3.3	0.8	2.2	0.9	0.0	0.0	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	81.2	3.8	14.1	2.3	3.2	1.4	1.5	0.8	0.0	0.0	0.0	0.0	458
Rural	81.4	1.9	13.6	2.0	3.8	0.9	1.1	0.5	0.0	0.0	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	74.3	5.2	14.2	4.7	9.7	2.6	1.7	1.2	0.1	0.2	0.0	0.0	137
Q2	79.1	3.9	15.0	2.8	4.1	2.1	1.8	1.1	0.0	0.0	0.0	0.0	208
Q3	82.6	3.8	13.1	3.0	4.1	2.2	0.2	0.2	0.0	0.0	0.0	0.0	206
Q4	82.5	4.8	13.4	3.7	1.0	0.7	3.1	1.6	0.0	0.0	0.0	0.0	223
Q5(richest)	85.2	2.6	13.1	2.3	1.7	1.1	0.0	0.0	0.0	0.0	0.0	0.0	252
M	37.6	28.8	62.4	28.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	89.6	3.0	8.2	2.6	1.5	1.5	0.7	0.7	0.0	0.0	0.0	0.0	235
30-44	87.4	2.3	11.1	2.2	1.5	0.8	0.0	0.0	0.0	0.0	0.0	0.0	329
45-59	81.8	2.9	12.8	2.8	3.4	1.3	2.0	1.3	0.0	0.0	0.0	0.0	262
60-69	66.7	4.7	23.9	4.6	7.5	3.0	1.8	1.5	0.0	0.0	0.0	0.0	116
70-79	56.9	9.8	21.7	6.7	14.8	7.6	6.4	3.1	0.1	0.1	0.0	0.0	70
80+	40.1	18.3	59.9	18.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	81.3	2.0	13.8	1.5	3.5	0.8	1.3	0.5	0.0	0.0			1,028

6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	35.2	4.4	5.7	1.0	4.5	1.5	1.4	0.6	0.2	0.2	53.2	4.7	502
Female	29.0	3.2	13.0	2.4	3.9	1.0	4.4	1.4	0.6	0.4	49.0	3.7	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	41.7	4.9	12.0	2.6	3.6	0.9	2.4	0.7	0.6	0.5	39.8	6.7	458
Rural	24.2	3.9	7.4	1.2	4.7	1.5	3.4	1.4	0.2	0.2	60.1	4.1	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	23.1	6.3	11.7	3.1	5.1	2.9	4.3	2.3	0.2	0.2	55.6	5.6	137
Q2	21.8	4.1	17.1	4.1	6.1	2.6	2.4	1.2	0.5	0.5	52.0	5.7	208
Q3	34.7	4.4	4.4	1.4	2.2	0.8	6.7	3.0	0.0	0.0	52.1	5.2	206
Q4	34.2	5.7	6.5	1.9	5.9	1.8	1.3	0.9	0.0	0.0	52.1	6.4	223
Q5(richest)	40.7	6.0	8.5	2.4	2.2	1.2	1.1	0.6	1.2	1.0	46.4	7.0	252
M	62.4	28.8	20.0	20.6	0.0	0.0	0.0	0.0	0.0	0.0	17.6	18.5	3
Age													
18-29	37.2	5.6	2.0	1.0	1.2	1.2	0.0	0.0	0.0	0.0	59.6	5.5	235
30-44	29.8	4.5	4.4	1.8	2.1	1.4	0.0	0.0	0.0	0.0	63.6	5.0	329
45-59	34.1	4.9	12.8	3.0	6.1	1.6	2.8	1.1	0.1	0.1	44.2	5.5	262
60-69	34.1	6.7	19.6	4.3	5.6	2.0	11.9	4.6	2.5	2.0	26.3	4.5	116
70-79	12.5	4.6	28.3	7.2	15.4	5.7	2.5	1.3	0.1	0.1	41.1	8.7	70
80+	37.7	17.3	10.2	7.6	0.0	0.0	43.1	21.2	5.6	5.8	3.5	3.6	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	32.0	3.1	9.4	1.4	4.2	1.0	2.9	0.8	0.4	0.2	51.1	3.8	1,028

6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	33.2	4.4	6.6	1.4	5.7	1.5	1.3	0.5	0.0	0.0	53.2	4.7	502
Female	28.3	2.9	11.8	1.9	5.8	1.9	4.8	1.3	0.2	0.2	49.0	3.7	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	40.5	5.2	10.0	2.1	7.0	2.0	2.6	0.8	0.0	0.0	39.8	6.7	458
Rural	22.8	3.5	8.7	1.5	4.8	1.4	3.4	1.0	0.2	0.2	60.1	4.1	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	23.6	4.4	10.1	3.2	5.9	1.9	4.7	1.8	0.2	0.2	55.6	5.6	137
Q2	23.7	4.0	13.0	3.5	7.8	2.8	3.0	1.5	0.5	0.5	52.0	5.7	208
Q3	33.6	4.4	7.1	2.0	3.0	1.0	4.2	2.8	0.0	0.0	52.1	5.2	206
Q4	31.2	5.4	7.6	1.9	6.7	1.9	2.3	0.5	0.0	0.0	52.1	6.4	223
Q5(richest)	37.2	6.2	8.9	2.2	5.6	2.0	2.0	1.1	0.0	0.0	46.4	7.0	252
M	62.4	28.8	20.0	20.6	0.0	0.0	0.0	0.0	0.0	0.0	17.6	18.5	3
Age													
18-29	36.3	5.2	2.9	1.4	1.2	1.2	0.0	0.0	0.0	0.0	59.6	5.5	235
30-44	31.2	4.4	3.1	1.3	2.0	1.4	0.1	0.1	0.0	0.0	63.6	5.0	329
45-59	32.8	5.1	11.4	2.9	9.8	2.9	1.7	0.8	0.1	0.1	44.2	5.5	262
60-69	26.1	5.1	20.9	4.3	11.9	3.2	14.8	3.7	0.0	0.0	26.3	4.5	116
70-79	9.2	3.7	31.5	7.4	15.2	5.4	2.8	1.4	0.1	0.1	41.1	8.7	70
80+	33.7	16.8	12.0	8.1	0.0	0.0	45.1	20.8	5.6	5.8	3.5	3.6	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	30.7	3.1	9.3	1.3	5.8	1.2	3.1	0.7	0.1	0.1	51.1	3.8	1,028

6.15 Problem with 'sleeping' in last 30 days - q2080

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	58.8	3.2	23.2	3.0	10.0	1.9	7.6	1.5	0.4	0.2	0.0	0.0	502
Female	47.5	2.0	31.2	3.1	11.8	1.9	7.1	1.2	2.4	1.2	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	47.5	4.1	30.3	3.9	13.1	2.1	8.7	1.2	0.4	0.2	0.0	0.0	458
Rural	57.5	2.3	24.8	2.3	9.2	1.5	6.3	1.2	2.3	1.0	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	46.6	5.2	26.1	5.1	16.4	3.5	6.3	2.4	4.6	1.9	0.0	0.0	137
Q2	43.9	4.9	32.4	4.2	13.5	2.5	10.0	3.0	0.2	0.2	0.0	0.0	208
Q3	55.1	4.3	24.5	4.1	12.1	3.6	4.6	1.8	3.8	2.8	0.0	0.0	206
Q4	59.7	4.4	22.7	3.6	8.0	1.8	9.6	2.2	0.0	0.0	0.0	0.0	223
Q5(richest)	57.1	5.8	29.6	5.6	7.4	2.5	5.9	2.7	0.0	0.0	0.0	0.0	252
M	0.0	0.0	62.4	28.8	17.6	18.5	20.0	20.6	0.0	0.0	0.0	0.0	3
Age													
18-29	78.7	3.3	16.5	3.9	3.0	1.3	1.8	0.9	0.0	0.0	0.0	0.0	235
30-44	63.6	4.2	25.9	3.7	5.7	1.6	4.8	2.1	0.1	0.1	0.0	0.0	329
45-59	46.7	5.4	36.2	5.3	8.1	2.2	8.4	2.0	0.7	0.4	0.0	0.0	262
60-69	19.0	4.0	26.2	3.9	38.8	4.3	13.5	4.7	2.5	1.0	0.0	0.0	116
70-79	9.9	4.7	39.5	7.0	25.1	8.1	19.9	5.3	5.6	3.4	0.0	0.0	70
80+	2.1	2.2	23.0	11.5	16.3	8.8	25.2	16.0	33.5	23.6	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	53.0	2.3	27.3	2.2	10.9	1.3	7.4	0.8	1.4	0.6			1,028

6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	54.2	3.3	24.4	2.8	15.0	2.7	6.2	1.2	0.2	0.2	0.0	0.0	502
Female	43.8	3.0	31.6	2.6	15.5	2.4	7.4	1.4	1.7	1.1	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	43.4	4.6	30.0	3.2	19.3	2.4	7.2	1.6	0.1	0.1	0.0	0.0	458
Rural	53.4	2.4	26.6	2.1	12.0	1.7	6.4	1.1	1.7	1.0	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	38.0	3.8	32.4	4.2	19.0	4.3	8.6	3.1	2.1	0.9	0.0	0.0	137
Q2	43.8	4.1	30.3	3.3	18.8	2.6	6.8	1.9	0.2	0.2	0.0	0.0	208
Q3	56.2	4.2	22.3	3.5	13.1	3.1	5.3	1.8	3.1	2.7	0.0	0.0	206
Q4	52.3	4.6	27.4	3.6	12.6	2.6	7.7	1.9	0.0	0.0	0.0	0.0	223
Q5(richest)	50.4	5.5	28.9	4.8	14.5	2.6	6.2	3.3	0.0	0.0	0.0	0.0	252
M	17.6	18.5	82.4	18.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	73.8	3.6	19.7	3.3	4.7	1.8	1.8	0.9	0.0	0.0	0.0	0.0	235
30-44	56.0	3.9	30.7	3.6	9.0	2.5	4.3	2.1	0.1	0.1	0.0	0.0	329
45-59	47.3	5.3	31.5	4.5	14.3	2.9	6.6	2.2	0.3	0.3	0.0	0.0	262
60-69	14.0	3.1	29.0	4.4	41.8	6.1	13.1	3.6	2.2	1.0	0.0	0.0	116
70-79	6.4	2.7	33.0	7.3	38.7	8.1	21.1	5.2	0.8	0.6	0.0	0.0	70
80+	7.0	5.8	17.8	9.7	16.4	8.9	25.2	16.0	33.5	23.6	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	48.9	2.4	28.1	1.9	15.2	1.5	6.8	0.9	1.0	0.5			1,028

6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	58.7	3.4	26.4	2.9	11.8	2.2	2.8	0.9	0.3	0.2	0.0	0.0	502
Female	44.6	3.3	30.0	3.0	13.1	2.0	10.9	2.2	1.4	0.6	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	44.8	3.9	29.9	2.4	15.8	2.6	9.2	2.0	0.4	0.3	0.0	0.0	458
Rural	56.9	2.6	26.9	2.4	9.9	1.8	5.1	1.4	1.2	0.5	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	36.1	4.5	28.8	5.3	22.5	4.7	8.5	3.1	4.2	2.1	0.0	0.0	137
Q2	48.9	4.4	31.4	3.0	13.2	3.6	6.1	1.6	0.3	0.2	0.0	0.0	208
Q3	55.9	5.0	22.3	4.3	14.2	2.3	7.1	3.1	0.5	0.4	0.0	0.0	206
Q4	56.2	4.3	25.2	3.0	9.9	2.7	8.3	2.2	0.3	0.3	0.0	0.0	223
Q5(richest)	54.6	5.8	33.1	4.4	6.8	2.2	5.3	2.1	0.2	0.2	0.0	0.0	252
M	20.0	20.6	0.0	0.0	62.4	28.8	17.6	18.5	0.0	0.0	0.0	0.0	3
Age													
18-29	74.9	4.7	17.4	3.8	5.0	2.2	2.5	1.1	0.3	0.2	0.0	0.0	235
30-44	56.3	3.5	29.6	2.9	8.5	2.1	5.5	1.8	0.1	0.1	0.0	0.0	329
45-59	49.0	5.6	30.9	5.1	11.5	3.5	8.4	3.0	0.3	0.3	0.0	0.0	262
60-69	26.6	4.2	33.0	5.0	29.5	3.8	7.4	3.5	3.4	1.3	0.0	0.0	116
70-79	13.6	4.6	39.2	7.4	28.8	7.1	14.1	3.7	4.3	3.2	0.0	0.0	70
80+	1.2	1.2	32.5	16.6	25.4	12.5	41.0	22.0	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	51.5	2.2	28.2	1.7	12.5	1.6	6.9	1.2	0.8	0.3			1,028

6.18 Problem with 'worry or anxiety' in last 30 days - q2091

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	54.3	3.6	25.3	2.3	12.6	1.8	6.8	2.0	0.9	0.7	0.0	0.0	502
Female	37.8	2.6	33.4	2.4	12.8	1.5	14.4	2.0	1.5	0.6	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	38.7	3.6	32.0	1.9	14.4	2.0	13.4	2.2	1.4	0.8	0.0	0.0	458
Rural	51.6	2.9	27.5	2.1	11.3	1.1	8.5	1.9	1.1	0.5	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	35.1	4.8	26.6	4.3	17.1	3.7	17.1	4.3	4.2	2.1	0.0	0.0	137
Q2	42.7	4.6	29.9	2.5	13.6	3.2	11.8	4.2	2.1	1.6	0.0	0.0	208
Q3	47.8	4.5	28.4	4.6	12.9	2.3	10.5	3.1	0.3	0.2	0.0	0.0	206
Q4	50.0	5.3	31.0	4.4	10.2	2.6	8.2	2.1	0.6	0.4	0.0	0.0	223
Q5(richest)	49.4	5.5	30.6	3.6	11.1	2.7	8.6	3.5	0.2	0.2	0.0	0.0	252
M	20.0	20.6	0.0	0.0	62.4	28.8	17.6	18.5	0.0	0.0	0.0	0.0	3
Age													
18-29	69.9	4.1	23.1	3.9	4.0	1.6	2.7	1.1	0.3	0.2	0.0	0.0	235
30-44	52.2	3.4	28.9	2.9	10.3	2.3	8.3	2.3	0.3	0.2	0.0	0.0	329
45-59	39.0	5.8	33.1	4.0	12.4	2.2	13.9	3.8	1.5	1.3	0.0	0.0	262
60-69	22.3	4.1	26.4	4.7	27.6	4.0	20.6	4.9	3.0	1.2	0.0	0.0	116
70-79	11.2	4.3	44.5	8.6	26.2	7.2	13.2	4.2	5.0	3.2	0.0	0.0	70
80+	1.2	1.2	32.5	16.6	25.4	12.5	41.0	22.0	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	45.9	2.3	29.5	1.4	12.7	1.0	10.7	1.4	1.2	0.4			1,028

7.1 Self-reported utilization of health services and perceived unmet need

	<u>Ambulatory Adult</u>		<u>Ambulatory Child</u>		<u>Inpatient Adult</u>		<u>Inpatient Child</u>		<u>Informal Care</u>		<u>Unmet Care</u>		<u>Non use</u>	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female	34.6	2.8	14.3	2.2	4.6	1.2	0.8	0.6	0.5	0.3	0.5	0.4	44.7	3.4
Male	32.7	4.0	17.4	2.8	2.0	0.7	1.7	0.7	0.0	0.0	0.0	0.0	46.1	4.3
Area of Residence														
Urban	43.6	4.9	15.4	2.0	1.5	0.7	0.1	0.1	0.4	0.3	0.0	0.0	38.9	5.4
Rural	26.4	3.2	16.1	2.8	4.7	1.2	2.1	0.8	0.1	0.1	0.4	0.4	50.2	4.2
Age Group														
18-29y	33.7	4.8	16.3	3.4	2.4	1.2	0.9	0.9	0.2	0.2	0.0	0.0	46.5	5.0
30-44y	22.5	3.6	27.4	4.0	1.6	0.8	2.4	1.1	0.0	0.0	0.0	0.0	46.1	4.9
45-59y	34.9	4.6	10.0	3.0	4.7	1.7	1.0	0.6	0.6	0.5	0.0	0.0	48.9	5.5
60-69y	54.1	5.3	2.1	1.7	5.9	2.3	0.0	0.0	0.5	0.4	2.0	1.7	35.5	4.8
70+y	49.2	7.5	0.0	0.0	6.2	2.5	0.0	0.0	0.0	0.0	0.1	0.1	44.6	7.9
Education Years														
0	38.5	7.2	3.4	2.6	12.1	4.2	0.0	0.0	0.6	0.7	0.0	0.0	45.3	6.0
1-5	46.3	7.5	3.0	2.1	5.5	2.3	0.0	0.0	0.1	0.1	0.0	0.0	45.2	8.0
6-11	30.4	3.4	12.6	3.1	3.3	1.2	1.3	0.8	0.0	0.0	0.1	0.1	52.3	4.3
12+	33.0	3.8	21.9	2.5	1.6	0.7	1.7	0.9	0.3	0.2	0.4	0.4	41.2	4.0
Income Quintiles														
Q1	36.1	5.2	11.2	4.2	3.7	1.6	2.7	1.9	0.0	0.0	0.3	0.3	45.9	6.2
Q2	35.7	5.7	10.3	3.4	3.9	1.4	0.0	0.0	0.1	0.1	0.0	0.0	50.0	6.1
Q3	28.6	4.0	18.6	3.2	2.6	1.3	0.2	0.2	0.3	0.2	0.0	0.0	49.7	5.2
Q4	37.6	4.9	16.1	4.1	2.7	1.5	1.7	1.2	0.5	0.5	0.0	0.0	41.4	4.6
Q5	31.6	5.7	19.7	2.8	3.9	1.6	2.0	1.5	0.2	0.2	0.9	0.9	41.6	5.7
Ownership														
Government	47.0	3.1	21.7	2.5	5.2	1.2	2.0	0.8	0.3	0.2	0.1	0.1	23.7	2.8
Private	94.2	6.0	5.8	6.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
NGO														
Other	16.5	16.4	0.0	0.0	0.0	0.0	0.0	0.0	22.5	20.7	0.0	0.0	61.0	25.2
TOTAL	33.7	2.8	15.8	1.8	3.3	0.8	1.3	0.5	0.2	0.1	0.3	0.2	45.4	3.3

This table presents the percent of people reporting utilization of health services when they last needed care. It combines information from Q7000, Q7001 and Q7016. Utilization was recorded in the WHS in three separate categories: ambulatory care, hospital inpatient care, and informal care. Time cut-offs for utilization recall used here are 1 year for ambulatory care and informal care, and 3 years for hospital inpatient care. Utilization of ambulatory and inpatient care services

have been further sub-divided into "adult" and "child" categories. In addition to utilization, two other related aspects are being reported here - unmet need and non-use (regardless of expressed need) of health services. Unmet need was estimated on the basis of responses to Q7000 and Q7004. Non-use of health services is simply the percent of respondents who reported no need in Q7000 or did not utilize services in the time cut-offs mentioned above.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, and ownership (of health facility). Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership was constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services.

7.2 Patient assessed characteristics of ambulatory health services

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
Sex																
Female	83.8	4.1	99.6	0.3	93.2	2.8	85.0	4.4	24.1	2.4	64.4	4.4	68.1	21.8	2.1	1.4
Male	88.7	3.8	99.0	0.8	92.7	2.8	70.9	5.9	21.3	1.8	73.4	4.2	37.0	4.6	3.2	2.0
Area of Residence																
Urban	85.5	2.8	99.3	0.4	96.9	1.6	76.4	6.2	19.2	2.7	58.8	5.3	37.7	4.9	1.1	0.7
Rural	87.1	4.8	99.3	0.7	89.1	3.2	79.2	5.3	26.1	2.3	78.9	4.4	63.3	20.6	4.2	2.2
Age Group																
18-29y	76.9	6.4	98.4	1.4	96.8	1.7	92.3	3.4	24.5	3.2	73.4	4.8	26.8	3.4	3.7	3.7
30-44y	84.4	5.3	99.5	0.5	94.0	3.0	67.7	7.2	19.3	2.2	74.9	4.6	42.4	13.8	1.1	0.8
45-59y	90.1	4.1	99.5	0.5	92.7	3.8	74.3	6.8	25.4	4.3	59.1	7.7	91.9	49.9	0.4	0.4
60-69y	96.4	2.3	100.0	0.0	88.3	4.3	75.1	7.5	23.0	3.5	66.8	7.8	48.7	7.9	2.0	1.8
70+y	99.9	0.1	100.0	0.0	83.1	9.5	87.0	5.4	23.6	2.9	56.8	10.9	43.2	13.7	13.1	8.7
Education Years																
0	96.8	2.6	100.0	0.0	81.9	11.6	89.5	8.5	23.9	3.3	75.3	10.0	39.2	15.7	9.7	7.8
1-5	78.7	16.1	100.0	0.0	96.6	2.0	81.6	10.1	20.6	2.8	79.7	7.6	51.2	9.0	2.1	1.4
6-11	88.6	4.5	99.4	0.6	96.1	2.2	87.9	3.0	29.2	3.6	46.8	7.6	38.0	6.2	4.6	3.7
12+	84.8	4.4	99.1	0.7	92.5	2.5	71.0	5.9	19.6	2.5	77.6	3.8	62.5	21.8	0.8	0.5
Income Quintiles																
Q1	85.6	8.2	100.0	0.0	82.4	7.2	79.8	6.7	30.2	3.2	76.2	6.6	34.5	9.0	8.6	5.0
Q2	80.4	7.2	100.0	0.0	92.9	3.3	81.8	6.8	24.2	3.3	63.9	10.9	40.9	9.4	0.8	0.8
Q3	91.8	4.6	99.6	0.4	85.4	6.0	73.7	8.1	26.4	3.8	70.6	7.3	40.2	8.3	4.8	4.6
Q4	86.4	4.4	97.5	1.9	99.2	0.8	86.5	5.6	18.0	2.3	71.0	6.8	27.4	5.5	0.4	0.4
Q5	86.3	5.7	99.6	0.4	99.6	0.4	70.3	7.1	18.4	3.2	67.2	6.6	99.2	51.0	1.1	1.1
Condition																
Child-Visit	91.1	5.2	100.0	0.0	96.4	3.6	73.7	9.0	17.7	2.5	86.4	5.3	22.6	0.5	0.0	0.0
Preventive	100.0	0.0	100.0	0.0	87.8	10.6	100.0	0.0	25.4	6.2	77.4	12.0	40.5	0.5	0.0	0.0
Childbirth	94.4	3.2	100.0	0.0	99.7	0.2	73.5	7.1	18.5	2.6	78.9	5.0	27.3	4.6	0.0	0.0
Acute	89.0	10.6	100.0	0.0	90.0	9.6	68.9	16.5	16.3	4.2	93.0	5.4	13.3	3.6	0.0	0.0
Injury	100.0	0.0	100.0	0.0	89.8	7.9	91.4	5.0	26.2	4.4	70.2	8.9	51.6	10.6	8.1	7.7
Chronic	78.9	5.8	98.6	0.9	95.1	1.9	82.8	4.4	23.7	2.8	65.1	5.6	77.2	23.9	1.3	0.8

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
Other																
Ownership																
Government	88.0	2.5	99.4	0.4	92.7	2.0	77.1	4.3	22.6	1.9	68.9	3.7	35.1	3.6	2.5	1.2
Private	42.6	16.2	97.4	2.7	97.4	2.7	93.0	6.9	24.6	4.7	75.6	11.4	215.0	104.7	7.0	6.9
NGO																
Other	77.5	20.7	100.0	0.0	100.0	0.0	77.5	20.7	22.4	14.7	43.9	28.4	75.1	18.7	0.0	0.0
Self-Assessed Health																
Very-Good	87.0	5.3	99.6	0.4	96.4	2.4	85.3	7.0	23.0	2.9	65.5	6.3	46.9	22.3	3.9	3.7
Good	84.4	3.4	99.1	0.9	93.5	2.5	75.3	7.0	23.6	3.8	76.4	6.4	33.7	5.4	1.0	0.7
Moderate	82.5	6.8	99.6	0.4	91.9	3.3	79.8	5.8	19.5	1.8	63.8	6.6	88.7	49.1	1.4	1.1
Bad	98.2	1.8	98.5	1.5	84.1	5.6	71.6	9.5	26.2	3.6	62.0	11.4	50.2	8.8	8.4	5.1
Very-Bad	100.0	0.0	100.0	0.0	100.0	0.0	9.8	11.8	0.5	0.5	74.9	23.6	100.5	0.5	0.0	0.0
TOTAL	86.3	2.8	99.3	0.4	92.9	1.9	77.8	4.1	22.7	1.9	69.0	3.6	52.6	12.6	2.6	1.2

This table looks at various characteristics of a visit to an ambulatory care facility, for those who utilized such services in the previous 12 months (Q7204 and Q7205). Characteristics of a visit reported here include:

- (1) The percent of patients visiting a medical doctor (of those visiting any ambulatory provider) (Q7302).
- (2) Patients' rating of *skills*, *equipment*, and *drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7304, Q7305 and Q7306, respectively.
- (3) Barriers or facilitators to care access have been reported here for those who utilized ambulatory care services in the last 12 months. They include:
 - (a) *Mean time taken to reach facility*: patient reported times from Q7307 were used to calculate averages across sample sub-groups.
 - (b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7308.
 - (c) *Mean cost of visit*: patient reported costs in Q7309-Q7313 were summed up for an individual and then averaged across sample sub-groups.
 - (d) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination listed in Q7328-Q7334.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include: sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents could select one of the

following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex												
Female	19.3	5.0	21.2	4.4	10.3	3.3	12.0	3.6	18.1	4.4	27.5	4.0
Male	17.8	3.9	26.1	4.4	22.1	4.2	15.1	3.9	17.3	3.6	22.1	4.4
Area of Residence												
Urban	10.4	3.5	17.1	4.2	11.4	3.9	7.7	3.2	11.2	3.2	22.0	4.1
Rural	26.4	6.3	30.0	6.0	21.2	5.3	19.3	5.0	23.9	4.9	27.2	5.2
Age Group												
18-29y	12.1	4.2	18.4	5.9	12.9	5.0	9.8	4.1	12.4	4.7	22.3	5.5
30-44y	17.9	5.0	24.3	4.4	14.1	4.0	12.7	4.2	18.3	4.6	25.0	5.6
45-59y	24.4	6.2	32.1	7.2	22.9	6.5	17.2	5.7	18.0	5.4	24.6	5.8
60-69y	24.2	7.2	17.6	4.9	15.3	5.4	12.8	4.5	20.3	6.1	23.0	6.2
70+y	18.6	9.2	30.2	10.0	24.6	9.8	24.1	9.8	28.6	10.0	36.1	11.1
Education Years												
0	22.4	9.3	21.6	8.5	16.7	8.9	16.1	8.9	17.6	8.8	24.3	11.1
1-5	23.4	9.8	28.8	10.1	18.0	8.7	17.8	8.1	19.6	8.8	25.9	8.6
6-11	23.9	6.2	23.7	5.6	18.0	4.7	13.2	4.0	17.1	3.6	20.9	4.6
12+	14.6	3.6	23.2	4.2	15.3	3.7	12.9	3.5	17.7	3.8	26.5	4.4
Income Quintiles												
Q1	41.8	10.2	51.2	8.6	26.5	8.2	28.3	8.7	24.6	7.4	33.7	9.7
Q2	26.2	8.2	23.5	8.2	17.7	7.0	11.0	5.4	19.6	7.0	22.2	6.8
Q3	14.7	5.2	17.6	5.0	14.5	4.9	15.4	5.0	18.4	6.0	30.1	6.5
Q4	17.8	5.3	23.5	7.0	15.3	5.5	13.4	5.0	19.3	5.9	33.0	7.4
Q5	4.8	3.3	15.0	5.0	12.7	4.9	7.0	4.0	10.9	3.8	11.0	3.5
Condition												
Child-Visit	12.5	6.6	7.2	4.2	8.9	5.1	8.9	5.1	9.0	4.2	17.2	9.2
Preventive	29.0	17.2	47.6	19.7	29.5	16.2	29.5	16.2	29.5	16.2	29.5	16.2
Childbirth	19.5	6.0	22.5	5.2	10.3	3.9	6.9	3.6	15.4	5.2	17.8	6.0
Acute	11.8	10.6	19.3	12.3	25.2	15.7	10.0	9.6	0.0	0.0	16.9	12.1
Injury	29.3	10.8	26.5	9.4	20.7	9.2	17.7	9.2	15.6	8.2	31.3	10.5
Chronic	13.7	4.0	20.2	5.1	16.9	5.1	15.4	4.7	20.6	4.6	28.6	5.6
Other												
Ownership												
Government	18.0	3.5	23.6	3.8	16.7	3.4	14.0	3.1	17.2	3.2	24.6	3.3
Private	22.3	16.5	19.5	16.6	0.0	0.0	0.0	0.0	20.4	13.0	17.8	12.8
NGO												
Other	80.1	19.9	43.9	28.4	43.9	28.4	43.9	28.4	61.0	25.2	61.0	25.2
Self-Assessed Health												
Very-Good	15.7	5.1	16.6	5.3	7.9	3.6	8.0	3.8	17.8	5.2	22.2	6.1
Good	13.9	4.7	21.6	5.2	17.5	4.7	14.5	4.6	16.3	4.6	23.5	4.6
Moderate	20.3	5.6	27.0	6.2	18.5	5.4	14.5	4.6	16.0	4.2	23.6	5.3
Bad	39.3	12.0	42.9	10.8	29.4	11.7	23.0	7.4	27.7	7.8	39.3	8.9
Very-Bad	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TOTAL	18.5	3.6	23.7	3.8	16.4	3.3	13.6	3.0	17.7	3.0	24.7	3.3

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Autonomy				Confidentiality				Choice of Healthcare Provider		Basic Amenities			
	Treatment Information		Involvement		Talk Privately		Confidentiality of Records		%	SE	Cleanliness	SE	Space	
	%	SE	%	SE	%	SE	%	SE					%	SE
Sex														
Female	26.3	4.7	28.1	5.5	19.3	4.8	13.3	4.1	31.3	5.7	30.8	6.7	38.3	6.4
Male	28.5	4.5	24.0	4.4	18.2	4.4	10.7	3.2	35.1	6.1	22.6	4.6	34.7	5.2
Area of Residence														
Urban	18.5	4.6	17.5	5.2	12.7	4.6	6.9	3.7	30.6	7.2	25.6	7.1	33.7	7.4
Rural	36.1	4.9	34.3	5.5	24.6	5.7	16.8	3.8	35.8	5.4	27.6	3.8	39.1	5.7
Age Group														
18-29y	16.2	5.4	17.9	5.8	13.1	5.1	10.3	4.5	20.8	5.9	29.6	8.6	32.1	8.4
30-44y	36.5	5.3	29.6	5.9	21.7	6.2	9.9	3.3	35.8	8.7	22.1	5.2	40.9	6.8
45-59y	25.3	5.7	27.7	7.1	18.8	5.8	9.7	4.2	36.3	5.9	27.5	7.2	38.3	8.0
60-69y	24.8	6.7	24.8	6.9	15.1	5.4	13.1	5.2	43.1	8.1	20.8	6.0	25.9	6.6
70+y	40.3	11.0	38.0	11.0	33.9	11.2	32.4	11.2	41.2	10.5	45.5	11.1	48.7	11.0
Education Years														
0	29.0	11.1	25.4	11.1	24.6	11.1	25.0	11.0	39.7	10.2	39.7	12.0	44.6	11.8
1-5	27.3	8.6	25.1	9.5	19.6	8.8	13.2	7.3	35.2	12.5	20.3	7.0	25.9	8.1
6-11	24.8	5.4	26.9	5.7	14.1	3.9	7.9	2.9	29.6	6.0	20.2	4.5	25.5	4.4
12+	28.6	4.2	25.8	4.7	20.2	4.9	12.0	3.4	33.8	6.0	28.9	5.9	42.2	6.7
Income Quintiles														
Q1	38.4	9.5	38.8	9.9	28.5	8.9	19.7	7.8	43.8	9.4	23.9	8.0	39.1	7.9
Q2	23.4	6.1	20.8	7.7	16.3	6.4	10.2	4.4	31.7	8.8	31.6	6.0	38.0	6.2
Q3	29.8	7.2	28.0	7.1	19.3	6.6	14.1	4.9	37.3	7.8	34.0	10.2	52.7	9.9
Q4	34.6	7.1	34.0	7.7	29.1	7.3	14.5	5.5	36.1	7.1	32.0	7.4	45.7	9.2
Q5	17.3	3.9	15.3	5.2	7.3	3.1	5.7	3.9	23.4	7.9	13.6	4.9	12.9	4.7
Condition														
Child-Visit	22.9	9.3	7.9	4.0	14.6	9.1	6.0	4.4	21.8	11.0	33.4	9.8	36.7	11.2
Preventive	29.5	16.2	29.5	16.2	29.5	16.2	29.5	16.2	29.5	16.2	57.0	16.3	57.0	16.3
Childbirth	28.2	7.2	18.5	6.1	16.7	6.8	6.5	3.6	36.9	11.2	24.1	7.6	35.7	8.2
Acute	16.9	12.1	16.9	12.1	16.9	12.1	21.0	13.8	37.5	17.3	19.0	11.9	43.4	17.9
Injury	32.0	10.5	30.6	10.4	22.1	9.8	19.2	10.0	34.5	10.5	20.4	10.0	25.4	10.2
Chronic	29.0	4.6	29.5	5.8	19.4	5.7	11.4	4.2	25.4	5.0	26.0	5.2	29.7	6.2
Other														
Ownership														
Government	27.1	3.7	25.6	4.0	18.7	3.8	12.5	2.9	34.3	4.7	27.5	4.1	37.7	4.7
Private	25.3	15.6	24.3	14.0	13.1	12.2	0.0	0.0	0.0	0.0	0.0	0.0	2.8	2.9
NGO														
Other	83.5	16.4	83.5	16.4	66.4	22.6	0.0	0.0	66.4	22.6	52.9	30.6	52.9	30.6
Self-Assessed Health														
Very-Good	31.6	6.6	25.5	6.5	23.0	6.4	10.5	4.6	30.1	7.4	32.2	8.0	42.6	9.1
Good	21.5	4.7	23.4	5.8	16.4	4.8	8.8	3.2	30.4	6.1	28.6	4.9	38.2	5.6
Moderate	26.4	5.5	24.5	5.9	14.9	4.9	13.2	4.7	33.6	6.0	20.8	4.6	31.3	5.7
Bad	43.9	11.7	42.3	8.8	27.3	8.0	25.1	8.2	46.7	8.6	21.0	7.4	30.0	7.5
Very-Bad	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	74.9	23.6	0.0	0.0	0.0	0.0
TOTAL	27.5	3.6	26.0	4.0	18.7	3.8	12.0	2.7	33.3	4.6	26.6	4.0	36.4	4.6

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.4 Patient assessed characteristics of hospital inpatient health services

	Patients staying for more than 6 days on last hospital visit		Patients rating characteristics of health care provider as adequate / more than adequate				Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Patients waiting more than a month to be admitted		Mean number of patients sleeping in the same room		Mean cost for a hospital visit		Percent of population perceiving discrimination			
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	%	SE	No	SE	mean cost	SE	%	SE
Sex																				
Female	62.1	6.5	98.2	1.2	93.0	3.0	87.3	3.4	48.1	5.9	88.3	3.2	0.0	0.0	4.0	0.7	284.3	0.5	2.0	1.5
Male	71.7	5.6	95.9	2.6	91.9	3.8	83.2	5.3	48.0	5.4	73.8	4.6	0.0	0.0	3.8	0.7	235.6	0.5	3.5	1.9
Area of Residence																				
Urban	72.4	7.0	98.0	1.5	91.7	4.0	83.2	5.8	50.7	7.8	71.0	5.8	0.0	0.0	4.0	0.8	146.6	0.5	1.8	1.9
Rural	61.7	6.1	96.6	1.9	93.1	3.8	87.2	3.1	46.1	3.6	90.2	2.3	0.0	0.0	3.9	0.6	322.8	0.5	3.2	1.5
Age Group																				
18-29y	34.7	9.4	99.3	0.7	96.7	1.9	92.3	5.1	38.1	6.9	76.0	9.0	0.0	0.0	4.3	1.2	162.0	0.5	0.0	0.0
30-44y	61.4	7.4	95.9	2.1	88.6	4.2	88.7	4.1	44.7	7.0	87.4	4.1	0.0	0.0	3.5	0.7	476.4	0.5	3.2	2.1
45-59y	83.5	6.3	95.9	4.0	95.8	4.0	78.6	7.2	51.6	6.7	89.7	4.0	0.0	0.0	4.0	0.7	168.9	0.5	1.9	1.9
60-69y	75.0	8.1	100.0	0.0	92.7	4.9	83.5	7.6	53.3	10.0	72.9	8.5	0.0	0.0	4.8	0.9	153.1	0.5	6.4	4.4
70+y	94.5	4.0	100.0	0.0	95.2	4.8	78.6	10.8	73.0	25.4	49.1	14.5	0.0	0.0	3.4	0.7	307.0	0.5	0.3	0.3
Education Years																				
0	87.6	8.3	100.0	0.0	99.3	0.8	76.3	7.4	50.7	7.9	97.1	1.7	0.0	0.0	4.2	0.8	194.6	0.5	10.1	7.6
1-5	77.2	9.5	100.0	0.0	92.4	7.2	83.4	10.0	72.6	22.4	58.5	11.2	0.0	0.0	4.8	1.1	210.1	0.5	0.0	0.0
6-11	67.9	6.8	93.3	3.8	92.6	3.9	89.0	4.5	45.8	6.0	83.3	4.9	0.0	0.0	3.7	0.7	417.0	0.5	1.6	1.6
12+	59.2	5.6	98.8	0.8	91.0	3.6	85.6	4.9	46.1	6.6	80.7	3.7	0.0	0.0	3.9	0.7	175.9	0.5	2.0	1.2
Income Quintiles																				
Q1	72.9	10.6	100.0	0.0	99.9	0.2	85.7	8.0	52.9	7.8	86.0	7.3	0.0	0.0	4.6	0.9	266.4	0.5	11.4	6.2
Q2	79.8	7.7	98.2	1.7	93.9	3.5	83.2	5.9	65.9	14.0	84.3	4.8	0.0	0.0	4.0	0.7	261.8	0.5	5.5	4.0
Q3	62.9	10.1	100.0	0.0	90.5	6.0	90.3	5.0	36.3	3.5	88.4	4.6	0.0	0.0	3.7	0.8	69.3	0.5	0.0	0.0
Q4	60.9	8.2	94.0	3.7	89.3	5.3	84.2	6.5	44.7	7.3	74.8	6.2	0.0	0.0	4.1	1.0	435.4	0.5	0.8	0.8
Q5	57.8	9.7	95.3	4.3	92.3	4.8	83.9	7.1	44.8	7.1	78.6	5.7	0.0	0.0	3.7	0.9	95.3	0.5	0.0	0.0
Condition																				
Child-Visit	75.0	11.8	95.4	4.3	94.2	4.8	85.5	8.0	32.0	4.4	85.8	8.2	0.0	0.0	3.3	0.9	259.3	0.5	9.4	7.2
Fever	54.2	12.2	95.0	3.9	94.9	3.9	92.5	5.3	51.7	16.2	93.8	4.2	0.0	0.0	3.2	0.8	289.0	30.0	0.0	0.0
Childbirth	13.2	6.3	95.8	4.0	85.9	8.1	87.5	8.4	35.3	5.3	86.0	7.7	0.0	0.0	3.7	0.8	169.9	127.2	4.2	4.0
Arthritis	95.5	3.8	100.0	0.0	100.0	0.0	100.0	0.0	49.5	7.1	89.1	11.1	0.0	0.0	4.1	0.9	72.8	2.7	0.0	0.0
Asthma	100.0	0.0	100.0	0.0	62.8	28.1	41.4	25.7	99.9	37.6	41.4	25.7	0.0	0.0	3.0	1.0	70.5	0.5	0.0	0.0
Heart-Disease	81.0	11.8	100.0	0.0	84.9	11.5	80.5	8.6	43.4	5.3	72.5	9.8	0.0	0.0	4.2	0.9	209.7	66.8	5.1	5.0
Injury	57.5	14.7	100.0	0.0	93.4	6.5	93.4	6.5	47.8	14.6	56.5	15.6	0.0	0.0	4.1	0.9	157.9	70.4	0.0	0.0
Minor-Surgery	72.5	7.8	100.0	0.0	98.0	2.0	79.2	6.8	53.9	7.4	81.4	7.1	0.0	0.0	4.6	0.9	596.2	256.6	2.7	2.1
Other	75.6	7.9	93.8	4.1	92.6	4.2	86.9	5.3	42.3	6.7	90.4	3.7	0.0	0.0	3.7	0.7	152.4	33.0	4.3	3.1
Ownership																				
Government	66.8	4.4	97.2	1.3	92.5	2.8	85.4	3.0	48.1	4.0	81.9	2.8	0.0	0.0	3.9	0.6	188.3	0.5	2.7	1.2
Private	0.0	0.0	100.0	0.0	100.0	0.0	100.0	0.0	40.5	0.5	0.0	0.0	0.0	0.0	4.5	0.5	1,430.5	0.5	0.0	0.0
NGO																				
Other																				
Self-Assessed Health																				
Very-Good	43.2	12.1	99.2	0.8	99.2	0.8	96.2	2.8	40.4	3.2	89.4	5.0	0.0	0.0	3.7	0.7	316.0	51.0	2.4	2.4
Good	58.1	7.0	96.8	2.2	90.9	3.8	80.9	6.2	42.1	5.9	80.6	5.2	0.0	0.0	4.2	0.8	138.6	37.4	2.7	2.3
Moderate	76.8	6.2	94.7	3.4	86.4	7.6	83.8	5.4	57.2	10.9	79.5	5.8	0.0	0.0	3.4	0.7	478.3	185.9	4.2	2.4
Bad	90.6	4.4	100.0	0.0	98.4	1.7	84.8	6.8	52.1	11.6	80.2	7.0	0.0	0.0	4.3	0.7	159.4	36.9	0.1	0.1
Very-Bad	87.0	12.1	100.0	0.0	100.0	0.0	90.2	8.5	57.9	16.6	86.6	10.2	0.0	0.0	5.1	1.1	340.2	55.7	0.0	0.0
TOTAL	66.2	4.6	97.2	1.3	92.5	2.7	85.5	3.0	48.0	4.0	82.1	2.8	0.0	0.0	3.9	0.6	266.0	0.5	2.6	1.2

This table looks at various characteristics of a visit to a hospital inpatient care facility, for those who utilized such services in the past 3 years (Q7200, Q7201 and Q7402). Characteristics of a visit reported here include:

(1) The percent of patients who stayed 6 days or more at an inpatient facility during their last visit (Q7404).

(2) Patients' rating of *skills, equipment, and drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7405, Q7406 and Q7407, respectively.

(3) Barriers or facilitators to care access have been reported here for those who utilized inpatient care services in the last 3 years. They include:

(a) *Mean time taken to reach facility*: patient reported times from Q7408 were used to calculate averages across sample sub-groups.

(b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7409.

(c) *Patients waiting more than a month to be admitted*: percent of patients who reported waiting more than a month to be admitted in Q7410.

(d) *Mean number of patients sleeping in same room*: averages across sample sub-groups have been constructed for numbers reported in Q7417.

(e) *Mean cost of visit*: patient reported costs in Q7411-Q7415 were summed up for an individual and then averaged across sample sub-groups.

(f) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination asked in Q7433-Q7439.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication				Autonomy			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions		Treatment Information		Involvement	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex																
Female	29.2	6.8	17.5	5.9	11.9	4.9	12.7	5.1	14.2	3.7	21.1	5.2	27.2	5.9	23.5	6.0
Male	22.1	6.0	20.0	5.2	12.7	4.7	10.9	4.3	16.8	4.3	24.7	6.1	23.9	4.6	23.3	5.2
Area of Residence																
Urban	25.7	7.5	22.2	8.3	7.9	3.7	9.5	4.1	15.6	3.7	21.7	5.3	22.7	5.0	27.5	8.5
Rural	26.5	6.1	16.0	4.3	15.4	6.2	13.8	5.1	15.2	3.9	23.3	6.5	28.0	5.7	20.5	4.4
Age Group																
18-29y	14.7	7.6	22.4	8.4	3.4	3.3	0.0	0.0	11.0	5.9	12.3	6.0	25.8	7.6	25.6	8.5
30-44y	26.8	8.4	20.0	8.2	17.3	7.5	15.0	6.1	14.2	4.5	27.7	8.0	30.0	7.1	29.4	7.9
45-59y	30.3	9.6	18.8	6.8	9.4	4.8	14.2	6.5	16.8	6.6	18.6	6.7	20.9	6.3	15.6	6.1
60-69y	32.3	7.1	15.0	5.0	15.0	7.3	15.6	7.3	25.0	6.8	30.9	8.2	23.4	8.6	20.5	7.3
70+y	20.9	10.5	5.1	4.8	5.1	4.8	5.1	4.8	6.5	4.9	11.4	7.4	20.8	7.0	13.6	7.9
Education Years																
0	16.9	6.0	7.1	7.3	7.1	7.3	7.1	7.3	17.5	10.4	16.1	10.4	10.0	7.9	9.5	7.9
1-5	26.8	9.5	8.2	5.3	7.2	5.2	8.6	5.4	9.4	5.4	8.7	5.4	15.6	8.1	20.3	7.6
6-11	38.0	9.3	27.3	9.9	14.3	4.9	10.1	4.3	15.1	5.4	23.2	7.2	26.7	5.8	35.6	8.8
12+	20.7	4.8	16.8	4.5	12.6	5.5	14.5	5.6	15.7	3.3	25.3	6.3	29.9	7.0	19.2	4.6
Income Quintiles																
Q1	50.8	10.4	16.0	8.4	9.8	5.6	5.3	4.2	15.2	8.6	17.5	8.5	19.2	8.2	20.9	8.3
Q2	21.1	7.1	11.9	6.1	15.0	10.3	15.4	10.3	11.3	4.4	21.7	10.3	26.1	10.1	10.9	4.2
Q3	27.4	12.8	29.5	12.7	14.0	6.9	8.7	5.7	12.3	6.1	21.6	8.0	27.0	8.6	43.3	12.0
Q4	27.1	6.4	20.5	6.9	7.5	4.6	9.1	4.9	14.2	5.8	16.6	5.8	25.4	6.9	16.9	6.2
Q5	16.5	7.0	14.2	6.8	14.3	6.8	18.5	7.1	23.5	7.0	30.6	9.2	28.9	8.8	23.9	9.2
Condition																
Child-Visit	41.5	18.3	35.9	19.0	8.0	6.0	8.0	6.0	10.4	6.5	20.8	9.9	20.4	10.5	37.6	18.2
Fever	29.3	7.6	13.3	7.3	4.0	3.7	4.0	3.7	7.0	4.4	20.3	8.9	21.0	8.9	7.2	4.5
Childbirth	8.0	4.8	8.0	4.8	8.4	5.1	8.4	5.1	12.0	5.7	6.4	3.8	22.6	9.9	24.9	9.7
Arthritis	46.8	22.6	9.8	9.0	1.0	1.0	38.0	24.6	38.0	24.6	8.8	8.9	8.8	8.9	8.8	8.9
Asthma	37.2	28.1	37.2	28.1	53.9	26.7	53.9	26.7	53.9	26.7	53.9	26.7	53.9	26.7	53.9	26.7
Heart-Disease	45.9	13.0	29.3	11.9	20.2	12.0	20.2	12.0	38.2	11.7	33.8	11.9	31.7	12.5	29.8	12.5
Injury	11.6	7.2	2.6	2.6	31.6	17.8	23.5	18.8	2.7	2.8	36.0	17.9	42.6	17.0	15.1	9.0
Minor-Surgery	19.1	7.7	16.8	7.2	4.3	2.7	3.4	2.5	11.1	5.8	14.4	6.4	24.2	8.8	16.9	6.5
Other	30.7	10.3	28.3	10.5	12.8	4.8	10.1	4.3	14.6	4.8	26.5	6.1	22.5	6.4	37.8	9.5
Ownership																
Government	25.5	4.8	18.8	4.3	12.4	4.1	12.1	3.5	15.5	2.8	22.8	4.4	26.0	4.0	23.6	4.4
Private	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
NGO																
Other																
Self-Assessed Health																
Very-Good	3.4	2.6	3.4	2.2	12.6	11.8	12.6	11.8	4.7	3.4	26.2	12.8	24.7	12.8	6.2	3.0
Good	13.5	4.5	13.9	5.1	11.3	4.3	8.3	3.9	14.0	4.7	20.2	6.0	26.1	7.4	24.4	6.8
Moderate	42.7	10.0	35.3	10.5	16.7	7.8	15.5	7.7	20.7	8.0	23.0	8.3	26.9	8.1	37.9	10.2
Bad	46.4	10.2	14.7	6.1	5.2	3.6	13.1	9.9	20.8	9.7	18.8	6.4	23.6	6.3	18.2	6.4
Very-Bad	56.9	17.7	24.0	14.5	10.8	9.2	10.8	9.2	20.8	12.5	37.6	17.0	27.6	16.2	9.9	9.2
TOTAL	26.2	4.7	18.6	4.3	12.2	4.0	12.0	3.5	15.4	2.7	22.6	4.4	25.8	4.0	23.4	4.4

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Confidentiality		Wait Time		Choice of health care provider		Clear Explanation		Basic Amenities Time for Questions		Treatment Information		Social Support Involvement	
	Travel Time %	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female	20.8	5.1	9.7	3.5	22.9	4.1	38.4	6.9	30.0	6.7	16.2	4.6	25.8	6.0
Male	21.4	5.5	12.9	4.9	31.8	6.3	31.2	6.7	31.3	7.0	24.4	7.0	27.4	7.0
Area of Residence														
Urban	16.5	4.9	7.2	3.2	24.6	4.0	41.7	8.6	34.4	6.7	16.6	4.3	29.7	7.0
Rural	24.4	6.0	13.9	3.7	28.2	4.9	30.7	6.1	27.8	5.0	22.0	5.8	24.1	5.6
Age Group														
18-29y	13.2	6.2	0.8	0.9	41.1	8.4	32.3	8.6	29.2	8.2	16.3	6.8	26.0	8.1
30-44y	28.6	8.1	13.1	4.7	20.1	5.4	33.6	8.2	29.4	8.2	17.7	6.4	28.2	8.4
45-59y	11.3	5.2	11.0	6.4	24.0	7.4	42.6	9.2	29.1	8.1	26.1	8.2	30.2	7.4
60-69y	26.1	8.7	17.4	7.3	30.1	7.4	38.1	8.4	33.3	7.8	19.6	5.6	20.5	5.7
70+y	16.4	8.4	10.0	7.3	38.3	15.0	18.9	10.6	42.8	13.5	17.3	6.3	13.7	8.2
Education Years														
0	12.7	8.4	9.5	7.9	23.5	14.5	49.0	17.0	30.5	15.4	15.5	9.2	24.3	8.5
1-5	22.2	8.9	15.2	8.4	40.4	11.6	32.2	10.8	39.3	11.3	25.8	10.2	18.9	9.0
6-11	16.5	5.5	11.0	4.6	25.4	6.1	32.0	10.2	27.4	9.3	17.4	6.1	32.0	10.1
12+	25.6	6.2	10.9	4.3	26.6	5.0	34.8	5.2	31.5	4.8	21.5	5.3	24.3	5.1
Income Quintiles														
Q1	11.7	5.3	4.9	4.2	23.5	5.9	27.3	9.8	19.3	8.7	20.3	8.6	22.4	8.6
Q2	23.9	10.1	7.8	3.4	31.4	7.4	33.6	10.5	27.3	6.8	18.6	5.8	16.7	5.8
Q3	26.1	8.8	12.9	6.8	22.0	8.3	41.9	11.6	36.4	12.0	13.9	6.7	39.4	12.4
Q4	11.7	5.6	3.3	2.6	27.8	7.0	29.0	9.5	26.7	9.0	15.4	5.8	22.7	8.6
Q5	28.0	8.6	23.1	6.1	28.2	6.5	41.8	6.9	38.2	7.2	30.5	6.2	29.3	6.4
Condition														
Child-Visit	7.5	5.9	5.8	4.8	17.6	9.4	43.2	16.6	39.7	18.2	13.6	8.6	39.2	18.3
Fever	13.2	8.8	13.2	8.8	25.7	11.1	35.0	10.6	26.8	8.2	35.3	8.8	34.9	8.7
Childbirth	21.3	8.4	9.9	5.3	31.3	10.1	18.5	7.3	29.9	10.3	20.0	8.6	25.9	9.6
Arthritis	28.7	15.4	48.0	21.2	56.8	18.6	75.1	14.7	37.1	24.9	37.1	24.9	37.1	24.9
Asthma	53.9	26.7	0.0	0.0	58.6	25.7	58.6	25.7	58.6	25.7	37.2	28.1	37.2	28.1
Heart-Disease	30.9	12.5	20.7	12.0	25.8	11.6	36.9	11.7	32.5	11.9	27.4	11.6	30.2	11.6
Injury	34.1	18.1	0.0	0.0	11.0	7.4	8.4	5.4	19.6	8.9	11.1	6.7	7.1	5.2
Minor-Surgery	17.4	6.8	2.6	1.6	21.6	7.0	51.1	10.1	34.4	9.8	6.9	4.1	16.5	5.6
Other	16.0	5.2	13.7	4.9	29.2	7.2	29.3	10.2	29.7	10.1	18.5	5.4	33.9	9.9
Ownership														
Government	21.2	4.2	11.2	2.6	26.9	3.3	35.6	5.0	30.8	4.1	19.9	3.8	26.7	4.4
Private	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
NGO														
Other														
Self-Assessed Health														
Very-Good	22.8	12.8	8.1	7.2	10.1	7.2	20.3	8.6	12.4	7.4	9.2	7.2	12.0	7.6
Good	20.9	5.6	7.1	3.1	22.4	6.2	36.5	7.6	31.0	6.8	16.6	4.4	21.4	4.8
Moderate	24.6	8.2	16.5	7.8	39.8	8.9	50.9	9.6	40.7	10.2	22.0	8.1	38.1	10.6
Bad	14.7	5.7	16.1	10.0	37.7	10.2	27.4	9.7	33.8	10.0	36.5	10.6	35.6	10.9
Very-Bad	9.9	9.2	2.8	2.6	10.0	9.6	10.0	9.6	27.8	14.9	24.9	15.3	24.9	15.3
TOTAL	21.1	4.2	11.1	2.6	26.7	3.4	35.3	5.1	30.6	4.2	19.7	3.9	26.5	4.4

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"

	Prompt Attention		Dignity		Communication		Autonomy		Confidentiality		Choice		Basic Amenities		Social Support	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex																
Female	63.2	4.2	63.8	4.1	63.3	5.0	58.4	4.3	64.0	4.3	59.5	4.9	61.1	5.0	58.6	4.8
Male	63.0	3.7	61.3	3.2	59.2	4.0	57.6	3.8	59.8	3.6	53.2	3.6	56.8	4.0	53.4	4.0
Area of Residence																
Urban	61.4	5.0	62.8	4.1	59.3	6.5	57.3	5.6	63.3	4.9	54.9	5.7	58.4	7.0	53.0	6.2
Rural	64.3	4.6	62.4	4.5	62.8	5.0	58.5	4.6	60.9	4.6	57.5	4.6	59.5	4.9	58.3	4.9
Age Group																
18-29y	63.1	5.0	63.7	4.6	59.4	6.3	54.5	4.7	62.0	5.0	52.8	6.0	56.0	5.9	54.9	6.1
30-44y	71.7	3.9	70.8	4.6	69.8	4.6	66.8	4.3	70.3	4.0	66.4	4.6	65.7	4.5	62.5	4.7
45-59y	58.0	5.6	58.5	4.8	60.0	5.9	57.4	6.2	57.9	5.9	54.2	5.1	60.5	6.3	53.8	5.7
60-69y	51.8	6.0	48.5	6.3	51.0	5.7	48.1	5.3	50.1	6.3	45.7	5.4	47.7	5.2	50.4	5.4
70+y	57.4	7.5	57.0	8.0	47.3	6.6	45.0	6.6	54.3	8.2	44.2	6.7	50.9	6.7	45.4	6.5
Education Years																
0	47.4	10.2	46.3	9.6	50.3	9.9	43.2	9.5	48.7	10.4	44.8	9.8	46.0	9.7	45.9	8.6
1-5	50.6	8.3	43.7	8.2	45.5	8.6	41.0	8.3	40.9	8.4	42.9	8.4	46.8	8.6	45.3	8.4
6-11	62.4	4.7	65.3	4.2	61.8	4.8	55.5	4.7	61.1	4.4	56.2	4.2	61.6	5.0	61.9	4.9
12+	68.2	3.6	66.6	3.6	65.4	4.1	64.9	4.0	68.0	3.4	60.7	3.7	61.7	4.2	55.9	4.3
Income Quintiles																
Q1	52.8	6.9	51.6	5.7	55.3	6.6	50.6	5.8	49.6	5.8	46.4	4.7	51.1	6.0	49.6	6.2
Q2	58.1	8.3	56.5	5.7	51.3	8.3	51.8	7.8	53.2	8.2	49.9	8.1	50.8	8.3	51.9	8.1
Q3	63.0	6.2	63.3	6.2	56.2	7.2	53.4	6.1	61.1	6.1	53.9	6.8	53.6	7.1	54.3	7.0
Q4	69.3	4.1	66.9	3.8	69.7	4.7	64.5	3.8	72.6	3.8	67.0	4.0	69.9	5.0	64.9	5.0
Q5	67.5	3.8	69.1	5.2	69.6	3.9	65.7	5.4	67.3	4.2	60.9	5.0	65.3	4.8	56.7	6.3
Ownership																
Government	63.1	4.0	62.9	3.5	61.5	5.2	59.8	4.1	62.4	4.2	56.3	4.6	56.7	5.0	55.5	4.8
Private	60.4	17.9	32.9	15.2	46.7	20.3	57.7	18.4	48.1	11.8	38.6	11.6	46.0	20.4	41.5	20.9
NGO																
Other	83.5	16.4	83.5	16.4	66.4	22.6	83.5	16.4	83.5	16.4	66.4	22.6	83.5	16.4	66.4	22.6
Self-Assessed Health																
Very-Good	76.6	4.0	75.5	4.0	69.5	4.7	67.0	5.2	70.9	4.1	66.4	4.4	68.8	5.2	66.5	4.9
Good	56.2	4.9	58.9	5.2	60.0	6.0	55.5	5.3	60.6	5.4	54.5	5.4	55.7	5.6	50.9	5.4
Moderate	57.6	5.3	58.1	5.7	59.6	5.6	53.3	5.3	55.1	5.4	52.4	5.1	57.3	5.4	54.7	5.6
Bad	62.3	5.4	51.8	6.1	48.6	5.3	53.6	6.5	61.7	4.7	47.6	5.4	49.0	5.6	46.1	5.3
Very-Bad	53.2	15.6	38.8	12.9	37.9	14.3	46.3	15.7	35.1	14.0	28.8	12.6	37.9	14.3	55.3	15.5
Type of Care																
Ambulatory-Adult	57.6	5.1	53.5	4.8	55.0	5.7	55.6	5.1	60.0	5.2	49.5	5.0	50.8	5.3	49.7	5.5
Ambulatory-Child	66.4	4.8	71.5	5.3	68.8	6.0	66.6	4.2	64.5	4.8	62.8	4.9	65.8	5.9	65.9	6.3
Inpatient-Adult	75.9	7.7	68.0	9.0	74.7	8.7	60.0	9.4	59.6	9.2	69.6	8.2	62.3	9.4	65.2	10.2
Inpatient-Child	78.7	14.5	90.2	9.8	55.7	20.2	75.5	15.2	87.0	10.6	67.2	18.8	51.0	20.6	55.7	20.2
Non-Users	64.6	4.4	65.0	4.3	62.6	3.9	56.2	4.8	61.9	4.4	58.0	4.2	62.8	4.5	56.8	4.2
TOTAL	63.1	3.4	62.6	3.1	61.3	4.0	58.0	3.6	61.9	3.4	56.4	3.6	59.0	4.1	56.1	3.9

This table shows the percent of people who rated a responsiveness domain as "Extremely Important". All survey respondents were queried on their perception of the importance of each of the eight responsiveness domains in section 7100 of the WHS where each domain was described in detail (Q7100-Q7107).

Respondents were asked to say whether the domain was "Extremely Important", "Very Important", "Moderately Important", "Slightly Important" or "Not

Important".

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, ownership (of health facility), self-assessed health, and type of care. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, self-assessed health and type of care sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad". Type of care is a constructed sample sub-group combining information from Q7000, Q7001 and Q7016. There are five categories of type of care: "Ambulatory adult", "Ambulatory child", "Inpatient adult", "Inpatient child" and "Non users".

8.0 Expenditure Background Indicators (2002)

GDP per capita (\$)	1,415
Total health expenditure per capita (\$)	130
Total population (000)	4,126
Life expectancy at birth (Both sexes)	72.8
Life expectancy at birth (male)	69.3
Life expectancy at birth (female)	76.4
Healthy life expectancy at birth (Both sexes)	64.3
Healthy life expectancy at birth (Male)	62.3
Healthy life expectancy at birth (Female)	66.4
Government funded health expenditure as a percentage of total health expenditure	49.8
Social security funded health expenditure as a percentage of total health expenditure	39.9
Private health insurance premiums as a percentage of total health expenditure	0
Out-of-pocket payment as a percentage of total health expenditure	50.2

8.1 Distribution of health payments across different population groups

	Mean Household expenditure (EXP) in local currency	% Poor	% Impoverished	% Catastrophic	OOP % EXP	OOP % non-subsistence spending	Mean OOP in local currency
Catastrophic							
No	523	24.5	1.4		3.9	6.7	26
Yes	433	39.5	30.9		47.5	63.2	198
Poor							
No	626		7.0	10.3	7.9	12.7	58
Yes	192		0.0	18.8	13.4	17.0	18
Insurance							
No	387	41.6	3.4	27.2	20.4	25.0	41
Yes	518	25.7	5.2	11.8	8.8	13.2	48
Residence							
rural	473	34.3	4.5	12.9	8.3	12.7	38
urban	553	17.8	5.8	12.2	10.5	15.0	58
Income_Quintile							
1st(poorest)	168	100.0	0.0	21.9	16.0	19.4	19
2nd	326	29.7	16.9	14.1	7.2	12.2	23
3rd	419	0.0	6.1	8.9	6.1	10.6	25
4th	613	0.0	1.5	9.5	8.0	13.6	49
5th(richest)	1,050	0.0	0.0	8.0	9.5	13.1	124
total	511	26.4	5.1	12.6	9.3	13.8	47

OOP: Out of pocket

Household consumption expenditure (EXP) comprises both monetary and in-kind payment on all goods and services and the money value of the consumption of home-made products.

Poor - a household is considered poor when its total household expenditure is smaller than its subsistence spending
Impoverishment - a non-poor household is impoverished by health payment when it becomes poor after paying for health services.

Catastrophic health expenditure occurs when a household's total out-of-pocket health payments equals or exceeds 40% of the households capacity to pay or non-subsistence spending.

Out of pocket health payments (OOP) refer to payments made by households at the point they receive health services. It includes doctor's consultation fees, purchases of medication and hospital bills. Spending on alternative and/or traditional medicine is included. Expenditure on health related transportation and special nutrition is excluded. Insurance reimbursements are also excluded.

Subsistence spending - household subsistence spending is the minimum requirement to maintain a basic life in a society. A poverty line is used in the analysis as subsistence spending.

Food expenditure is the amount spent on all foodstuffs by the household plus the value of the family's own food production consumed within the household. It excludes expenditure on alcoholic beverages, tobacco and food consumption outside the home (e.g hotel and restaurants).

Household capacity to pay - is defined as a household's non-subsistence spending.

8.2 Percent distribution of structure of out-of-pocket health payments

	Inpatient	Outpatient	Traditional	Drugs	Other	Total
Catastrophic						
No	3.1	0.4	4.2	48.1	44.2	100
Yes	11.0	1.9	0.5	29.3	57.3	100
Poor						
No	7.3	1.0	2.5	35.4	53.8	100
Yes	6.4	2.3	0.6	64.2	26.5	100
Insurance						
No	3.3	0.6	0.6	49.2	46.3	100
Yes	7.4	1.2	2.3	37.8	51.3	100
Residence						
rural	12.7	1.8	0.4	38.7	46.4	100
urban	3.3	0.7	3.6	38.0	54.4	100
Income_Quintile						
1st(poorest)	5.6	2.9	0.8	68.8	22.0	100
2nd	4.5	5.7	0.0	52.0	37.8	100
3rd	7.4	0.0	0.0	58.6	33.9	100
4th	6.5	1.7	9.7	38.2	43.9	100
5th(richest)	8.3	0.0	0.4	26.8	64.5	100
total	7.2	1.2	2.3	38.3	51.1	100

8.3 Financial sources used by households for paying for health services

	Savings	Sold Items	Borrow from relatives &	Borrow from others	Health insurance	Current Income	Other
OOP%non_subsistence_spending							
less than 10%	1.1	0.0	6.7	1.3	1.0	82.4	2.8
10-20%	12.5	1.3	12.8	2.9	1.6	93.0	1.2
20-40%	1.2	0.0	13.4	1.3	1.0	91.6	3.2
above 40%	4.3	2.5	13.2	9.8	1.4	87.3	7.1
Hospitalization							
No	2.5	0.3	7.7	1.9	0.6	85.3	2.4
Yes	2.8	1.6	19.1	7.4	4.9	84.5	10.6
Insurance							
No	4.6	3.4	24.9	6.6	0.0	61.8	10.7
Yes	2.4	0.3	8.2	2.3	1.2	86.4	2.9
Residence							
rural	1.5	0.0	6.2	1.1	1.3	87.9	2.9
urban	3.6	1.0	12.1	4.1	0.8	82.3	3.8
Income_Quintile							
1st(poorest)	0.8	1.4	12.8	3.1	1.2	86.2	8.2
2nd	1.3	0.0	3.1	0.3	0.0	84.2	2.1
3rd	0.0	0.0	9.2	6.6	1.2	78.2	1.4
4th	2.2	0.9	12.9	2.7	2.7	84.4	3.6
5th(richest)	8.2	0.0	7.5	0.3	0.5	92.9	1.0
total	2.5	0.5	9.0	2.5	1.1	85.2	3.3

8.4 Insurance Coverage

	Insured	SE	Mandatory Insurance	SE	Voluntary Insurance	SE	Insured as dependent	SE	Mandatory and Voluntary	SE
Q1(poorest)	12.2	1.8	12.3	1.8	0.0	0.0	12.0	2.1	0.0	0.0
Q2	17.8	2.5	17.9	2.4	50.7	26.6	17.9	3.0	61.0	29.9
Q3	20.2	1.7	20.1	1.7	0.0	0.0	20.9	2.2	0.0	0.0
Q4	22.7	2.1	22.9	2.1	0.0	0.0	22.4	2.4	0.0	0.0
Q5(richest)	27.2	3.0	26.7	2.8	49.3	26.6	26.7	3.8	39.0	29.9
Total	100		100		100		100		100	
Number	3,129.0		3,099.0		5.0		1,501.0		4.0	

10.1 Inability to control the important things in life - q8000

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
Gender													
Male	51.4	4.2	19.8	2.7	22.1	3.0	4.6	1.2	1.7	1.1	0.4	0.4	502
Female	34.2	2.9	22.5	2.4	34.3	3.3	7.9	1.7	0.9	0.4	0.2	0.1	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	37.0	4.6	22.2	3.2	32.5	4.0	6.6	1.5	1.0	0.5	0.6	0.5	458
Rural	47.0	4.1	20.3	2.4	25.0	3.0	6.0	0.9	1.6	1.0	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	25.8	5.9	19.4	3.4	40.7	6.3	12.2	3.2	1.9	1.2	0.0	0.0	137
Q2	41.4	5.0	18.0	3.7	32.1	5.5	4.8	1.4	3.6	2.6	0.0	0.0	208
Q3	38.9	5.4	30.1	4.1	23.7	3.8	6.2	2.7	1.2	0.7	0.0	0.0	206
Q4	46.4	5.7	17.9	3.4	26.6	4.2	7.6	2.0	0.4	0.4	1.2	1.0	223
Q5(richest)	52.7	5.9	19.9	3.8	24.2	3.8	3.2	1.7	0.0	0.0	0.0	0.0	252
M	0.0	0.0	62.4	28.8	20.0	20.6	17.6	18.5	0.0	0.0	0.0	0.0	3
Age													
18-29	56.6	5.1	15.0	3.2	23.5	4.1	2.6	1.7	2.3	2.2	0.0	0.0	235
30-44	51.3	4.4	20.9	3.0	20.7	4.2	5.5	1.6	0.8	0.5	0.8	0.6	329
45-59	38.0	5.3	24.5	3.5	30.8	5.3	5.3	1.8	1.3	0.8	0.0	0.0	262
60-69	14.3	3.5	23.7	4.1	54.0	5.3	7.2	2.3	0.8	0.5	0.0	0.0	116
70-79	24.2	7.2	27.1	6.3	29.8	7.4	17.3	5.0	1.6	0.9	0.0	0.0	70
80+	19.4	15.0	16.9	9.6	26.7	12.6	37.0	22.8	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	42.6	3.0	21.2	1.9	28.4	2.5	6.3	0.8	1.3	0.6	0.3	0.2	1,028

10.2 Inability to cope - q8001

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
Gender													
Male	43.9	4.2	23.8	3.0	26.8	2.7	3.0	0.9	2.2	1.2	0.4	0.4	502
Female	30.1	3.0	22.6	2.4	38.3	2.9	7.6	1.6	1.2	0.5	0.2	0.1	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	32.8	4.9	20.4	2.7	38.7	3.8	6.5	1.4	1.0	0.5	0.6	0.5	458
Rural	40.0	4.1	25.4	2.9	27.9	2.9	4.5	1.0	2.2	1.0	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	24.5	5.8	15.5	2.4	46.6	6.3	8.8	2.5	4.6	2.0	0.0	0.0	137
Q2	35.0	5.5	17.5	2.8	38.5	5.9	5.1	1.9	3.9	2.6	0.0	0.0	208
Q3	35.3	5.2	33.9	4.2	25.8	4.1	4.6	2.9	0.5	0.4	0.0	0.0	206
Q4	36.9	5.9	23.5	3.6	31.3	4.9	6.4	2.0	0.7	0.7	1.2	1.0	223
Q5(richest)	46.5	6.7	22.6	4.2	27.3	4.4	3.6	1.4	0.0	0.0	0.0	0.0	252
M	0.0	0.0	62.4	28.8	37.6	28.8	0.0	0.0	0.0	0.0	0.0	0.0	3
Age													
18-29	47.6	6.0	19.3	5.1	28.6	3.5	2.2	1.3	2.3	2.2	0.0	0.0	235
30-44	44.4	4.5	26.1	3.6	24.5	3.7	2.9	1.1	1.3	0.7	0.8	0.6	329
45-59	34.6	5.6	23.8	3.7	35.9	5.1	4.4	1.4	1.3	0.8	0.0	0.0	262
60-69	10.9	2.8	22.7	3.9	54.3	5.0	10.6	3.3	1.5	0.7	0.0	0.0	116
70-79	19.8	5.9	24.1	7.0	37.7	6.5	15.2	5.0	3.2	1.8	0.0	0.0	70
80+	19.4	15.0	11.4	7.4	32.2	14.4	37.0	22.8	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	36.8	3.1	23.2	2.0	32.7	2.4	5.4	0.8	1.7	0.6	0.3	0.2	1,028

10.3 Satisfaction with your health - q8002

	Very dissatisfied	SE	Dissatisfied	SE	Neither satisfied nor dissatisfied	SE	Satisfied	SE	Very Satisfied	SE	Missing	SE	N
Gender													
Male	5.7	1.4	31.8	3.0	29.0	3.2	13.4	2.5	19.7	3.8	0.4	0.4	502
Female	8.8	1.9	27.6	2.5	36.0	2.2	15.0	3.0	12.4	1.9	0.2	0.1	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	6.1	1.7	29.6	3.0	29.1	2.6	18.7	3.3	16.0	4.5	0.6	0.5	458
Rural	8.3	2.0	29.7	2.8	35.4	2.9	10.6	2.5	15.9	2.3	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	15.3	3.3	16.8	2.9	46.4	4.8	7.1	2.1	14.4	3.9	0.0	0.0	137
Q2	8.8	2.6	22.6	3.4	42.7	3.9	11.0	3.7	14.8	3.0	0.0	0.0	208
Q3	7.7	3.1	29.7	4.6	31.9	3.6	12.9	4.7	17.7	3.6	0.0	0.0	206
Q4	3.1	1.3	31.0	4.8	34.8	4.6	16.3	3.7	13.6	3.8	1.2	1.0	223
Q5(richest)	5.1	2.1	41.4	5.0	15.4	2.2	19.5	3.9	18.6	5.0	0.0	0.0	252
M	0.0	0.0	0.0	0.0	37.6	28.8	62.4	28.8	0.0	0.0	0.0	0.0	3
Age													
18-29	6.3	2.7	23.3	4.6	17.2	3.8	18.1	4.9	35.2	5.5	0.0	0.0	235
30-44	3.3	1.2	34.6	4.0	29.8	3.9	17.2	2.8	14.2	2.3	0.8	0.6	329
45-59	7.0	2.7	33.8	4.3	34.3	3.8	13.2	2.7	11.7	5.4	0.0	0.0	262
60-69	13.3	3.2	26.7	4.6	51.7	6.1	5.6	1.6	2.7	1.6	0.0	0.0	116
70-79	14.2	5.5	20.9	6.3	57.7	9.2	5.9	3.1	1.4	1.0	0.0	0.0	70
80+	33.5	23.6	14.4	8.4	39.9	18.6	12.2	8.2	0.0	0.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	7.3	1.3	29.7	2.1	32.6	2.0	14.2	2.0	16.0	2.4	0.3	0.2	1,028

10.4 Feeling safe when alone at home - q8011

	Completely safe	SE	Very safe	SE	Moderately safe	SE	Slightly safe	SE	Not safe at all	SE	Missing	SE	N
Gender													
Male	34.5	5.2	31.7	4.3	23.5	3.5	5.3	1.5	5.0	1.5	0.0	0.0	502
Female	16.4	2.9	24.9	2.5	33.4	2.5	17.6	2.6	7.6	1.4	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	15.2	3.6	29.8	3.2	38.5	4.1	10.4	1.8	6.1	1.2	0.0	0.0	458
Rural	33.3	5.3	27.0	3.8	20.6	2.4	12.6	2.3	6.5	2.2	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	31.7	7.5	23.0	5.3	27.8	5.0	9.2	3.1	8.3	3.1	0.0	0.0	137
Q2	27.2	5.7	31.0	5.2	24.9	3.5	10.4	2.7	6.5	2.2	0.0	0.0	208
Q3	29.8	5.6	25.9	3.7	25.7	5.4	11.9	3.1	6.6	2.2	0.0	0.0	206
Q4	21.8	4.3	28.0	4.4	29.7	6.0	14.4	3.4	6.1	2.3	0.0	0.0	223
Q5(richest)	19.7	4.8	31.0	5.4	32.9	4.5	11.2	2.2	5.2	2.0	0.0	0.0	252
M	0.0	0.0	20.0	20.6	62.4	28.8	17.6	18.5	0.0	0.0	0.0	0.0	3
Age													
18-29	25.1	4.7	32.6	5.3	30.4	4.7	7.1	2.4	4.7	1.6	0.0	0.0	235
30-44	29.0	4.6	23.3	3.8	32.7	4.8	13.1	2.7	1.8	0.8	0.0	0.0	329
45-59	23.3	4.5	31.0	5.4	24.1	4.7	11.6	2.8	9.9	3.2	0.0	0.0	262
60-69	21.1	4.3	23.7	6.8	30.3	5.2	12.7	2.8	12.3	3.7	0.0	0.0	116
70-79	16.8	4.8	36.3	7.4	23.1	5.5	15.0	5.1	8.7	4.4	0.0	0.0	70
80+	48.0	20.6	17.6	11.2	1.9	2.0	23.5	15.7	9.1	7.0	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	25.2	3.6	28.2	2.6	28.6	2.4	11.6	1.5	6.3	1.3			1,028

10.5 Feeling safe when walking down the street alone after dark - q8012

	Completely safe	SE	Very safe	SE	Moderately safe	SE	Slightly safe	SE	Not safe at all	SE	Missing	SE	N
Gender													
Male	20.5	3.2	39.8	4.2	27.9	4.0	6.8	1.7	4.9	1.4	0.0	0.0	502
Female	7.0	1.5	16.4	2.1	45.7	2.8	19.6	2.8	11.2	2.1	0.0	0.0	526
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	10.4	2.8	27.5	2.8	40.9	2.8	12.9	2.4	8.3	2.3	0.0	0.0	458
Rural	16.2	3.0	28.2	4.2	33.9	4.2	13.7	2.6	8.1	1.8	0.0	0.0	570
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	16.4	6.1	34.4	6.7	27.5	6.4	13.0	3.6	8.8	2.9	0.0	0.0	137
Q2	9.3	2.4	29.5	5.1	37.4	4.6	12.8	3.5	11.0	4.1	0.0	0.0	208
Q3	10.6	2.5	29.1	4.4	38.3	5.4	13.6	4.2	8.3	2.4	0.0	0.0	206
Q4	17.7	4.3	22.8	4.3	41.2	5.6	11.1	2.1	7.3	2.3	0.0	0.0	223
Q5(richest)	14.5	4.0	26.5	5.0	37.6	5.4	15.1	3.1	6.2	2.0	0.0	0.0	252
M	0.0	0.0	20.0	20.6	0.0	0.0	80.0	20.6	0.0	0.0	0.0	0.0	3
Age													
18-29	11.0	3.2	28.1	4.7	44.8	5.0	8.8	2.7	7.4	2.4	0.0	0.0	235
30-44	17.1	3.5	25.2	3.6	35.5	4.5	16.1	3.0	6.0	1.8	0.0	0.0	329
45-59	14.1	3.2	30.6	6.1	35.8	4.9	11.8	2.6	7.6	2.3	0.0	0.0	262
60-69	10.9	2.8	29.3	7.3	29.1	5.3	15.4	3.5	15.2	3.9	0.0	0.0	116
70-79	9.1	3.4	32.6	6.6	37.4	5.9	9.7	3.7	11.2	4.5	0.0	0.0	70
80+	10.4	7.8	3.2	2.6	29.4	17.0	48.6	20.6	8.3	6.7	0.0	0.0	17
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	13.6	2.1	27.9	2.6	37.0	2.7	13.4	1.8	8.2	1.4			1,028