

# **REPORT OF HUNGARY**

## TABLE OF CONTENTS

1	Introduction
2	Methodology
2.1	Sampling
2.2.1	Geographic Component of the Survey
2.2.2	The map
2.3	Survey Metrics
2.3.1	Sample deviation index
2.3.2	HH Sample deviation index
2.3.3	Response rate
2.3.4	Missing data
2.3.5	Reliability
3	Socio-Demographic Profile
3.1	Household composition by residence and age groups according to income quintile
3.2	Selected age distributions by income quintile
3.3	Characteristics of head of household by income quintile
3.4	Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex
3.5	Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education
4	Risk Factors
4.1	Prevalence of smoking and average daily tobacco consumption
4.2	Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers
4.3	Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

- 4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water
- 4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility
- 4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used
- 5 Coverage
  - 5.01 Tuberculosis
  - 5.02 HIV Prevention and Screening
  - 5.03 Antenatal Care and Mother to Child Transmission of HIV
  - 5.04 Birth Delivery Care
  - 5.05 Child's Health - Common Illness
  - 5.06 Child's Health - Danger Signs
  - 5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card
  - 5.08 Vaccination Status of Children 12-23 Months
  - 5.09 Women's Cancer Screening
  - 5.10 Oral Health and Injuries
  - 5.11 Self Reported Non-Communicable Conditions
  - 5.12 Mental Health Conditions
  - 5.13 Cataracts
- 6 Health State Descriptions
  - 6.01 In general, rate your health today - q2000
  - 6.02 Difficulty with 'work and household activities' in last 30 days - q2001
  - 6.03 Difficulty with 'moving around' in last 30 days - q2010
  - 6.04 Difficulty with 'vigorous activities' in last 30 days - q2011
  - 6.05 Difficulty with 'self-care' in last 30 days - q2020
  - 6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021

- 6.07 Amount of 'bodily aches or pains' in last 30 days - q2030
- 6.08 Amount of 'bodily discomfort' in last 30 days - q2031
- 6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050
- 6.10 Difficulty with 'learning a new task' in last 30 days - q2051
- 6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060
- 6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061
- 6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071
- 6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072
- 6.15 Problem with 'sleeping' in last 30 days - q2080
- 6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081
- 6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090
- 6.18 Problem with 'worry or anxiety' in last 30 days - q2091
- 6.19 Adjusted for vignette responses
- 7 Health System Responsiveness
  - 7.1 Self-reported utilization of health services and perceived unmet need
  - 7.2 Patient assessed characteristics of ambulatory health services
    - 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1
    - 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2
  - 7.4 Patient assessed characteristics of hospital inpatient health services
    - 7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1
    - 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2
  - 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"
- 8 Health Expenditure, Insurance and Human Resources for Health

- 8.0 Expenditure Background Indicators (2002)
- 8.1 Distribution of health payments across different population groups
- 8.2 Percent distribution of structure of out-of-pocket health payments
- 8.3 Financial sources used by households for paying for health services
- 8.4 Insurance Coverage
- 8.5 Human resources for health
- 9.1 Adult Mortality Chart 1
- 9.2 Adult Mortality Chart 2
- 9.3 Adult Mortality Chart 3
- 9.4 Adult Mortality Chart 4
- 10 Stress and Social Capital
- 10.1 Inability to control the important things in life - q8000
- 10.2 Inability to cope - q8001
- 10.3 Satisfaction with your health - q8002
- 10.4 Feeling safe when alone at home - q8011
- 10.5 Feeling safe when walking down the street alone after dark - q8012

# 1. Introduction

The World Health Organization initiated the World Health Survey (WHS) as a *data collection platform* to obtain information on the health of populations and on health systems. The WHS responds to the need of countries for a detailed and sustainable health information system and gathers data through surveys to measure essential population health parameters. The WHS uses standard survey procedures and instruments for general population surveys to gather comparable data across WHO Member States. These methods and instruments have been refined through scientific review of literature, extensive consultations with international experts and large-scale pilot tests. WHS is designed to evolve through its implementation by continuous input from collaborators, including policy makers, survey institutions, scientists and other interested parties. The data is jointly owned by the countries and the WHO, and there is a commitment for a long-term data collection, building local capacity and using this information for health policy.

The objectives of the World Health Survey were to:

- develop a **means** of providing:
  - **valid, relevant, reliable** and **comparable** information on **outcomes, functions and inputs** to health systems
  - to **supplement** the information provided by routine **HIS**
  
- build the **evidence base** necessary for policy-makers:
  - to **routinely monitor** if health systems are achieving the **desired goals**
  - and in some countries, to assess if the **increase in health resources** is having the desired effect

The introductory chapter of each country report will provide details on the current sources of data from routine health information systems and ad hoc surveys. It will describe the health situation in the country and identify gaps. It will also present a brief overview on the health system organization within the country.

## 2. Sampling

The WHO sampling guidelines emphasized scientific principles of survey sampling as explicit standards for quality, included examples of good sampling plans, and identified quality assurance standards for countries to adhere to. WHO and technical advisers provided technical support to countries when needed. The following principles were used for sampling in the WHS:

1. The sample should target the *de facto* population (i.e. all people living in that country including guest workers, immigrants and refugees) and not *de jure* population (i.e. the citizens of that country alone) with complete geographic coverage of the country, if possible, or a representative geographic segment of the country.
2. A *sample frame* with 90% coverage that included all key subgroups of interest. Enumeration or listing of households to update the frame was recommended.
3. The WHS sample identified all adult members of the general population aged 18 years or older as its sample. Households are selected using a random, stratified sampling procedure. One individual per household is then selected through a random selection procedure using the Kish table method.
4. The WHS used a sampling strategy which encompasses a known non-zero selection probability for any individual included in the study. This makes it possible to extrapolate the data to the whole population. Households and individuals are selected with a known probability in order to provide selection probabilities for each household and individual in the data set.
5. The WHS sampling strategy was without replacement and categories of non-respondents were clearly defined.
6. The WHS used GPS devices in certain countries and digitized maps to geocode the data within specified guidelines.

Each national report will present a detailed description of the sampling strategy and the allocation tables

---

## 2.2.1 Geographic Component of the Survey

---

Geographic Information refers to any information associated with a location on the Earth's surface. This location can be associated to a coordinate (latitude/longitude), a place's name, a postal address or any other geographic object.

Household Surveys represent an important source of information. If the geographic location of the surveyed clusters is collected at the same time that the interview is conducted users are then able to integrate the spatial dimension in their analysis

The geographic component of the WHO WHS is stored in two sections of the questionnaire:

- 0100 Sampling Information allowing the link between the location of each cluster and the sampling frame used for the survey. This section has also been used in order to collect the type of setting in which each household was located at the time of the survey. This section of the questionnaire has been used for all the countries part of the WHO WHS


- 0200 Geocoding Information for the collection of the latitude and longitude of each household. This section has only been used in 27 countries where GPS devices were available or have been provided. The choice of these countries has been done on the basis of the availability of digital maps delimiting the surveyed clusters, the field data collection planning and their distribution within the different WHO regions. Specific training and field material have been prepared and used by the interviewers or field supervisors in these countries in order to collect the latitude and longitude of each of the surveyed household.

As the sampling design may not always include the administrative structure of the country and as the codes entered in the data set for the sampling level needed to be labelled, a key correspondence table containing this information has been collected after the survey itself.

Once all the survey data and the key correspondence table available a specific protocol has been applied to the two sections mentioned earlier in order to clean eventual data collection or data entry mistakes as well as to fill the gaps in close collaboration with the survey institution.


The questionnaire as well as the training material and data cleaning protocols used can be downloaded from the WHO WHS web site (<http://www.who.int/healthinfo/survey/en/index.html>) in the "Survey Instrument and Related Documents" section.

### 2.3.1 Sample deviation index



SDI

### 2.3.2 HH Sample deviation index



HDI


---

### 2.3.3 Response rate


---

<b>Household</b>	
Selected	1,955
Interviewed	1,412
Percentages	72
<b>Individual</b>	
Selected	1,419
Interviewed	1,419
Percentages	100

### 2.3.4 Missing data



### 2.3.5 Reliability



### 3.1 Household composition by residence and age groups according to income quintile

	Percent Rural	SE	Mean Number of									
			Household members	SE	Children < 2 years	SE	Children < 5years	SE	Children 5 -17 years	SE	Persons 18+ years	SE
Q1(poorest)	44.2	4.26	1.6	0.05	0.0	0.00	0.0	0.01	0.1	0.02	1.5	0.04
Q2	32.9	3.35	2.0	0.06	0.0	0.01	0.0	0.01	0.1	0.02	1.9	0.05
Q3	28.9	3.40	2.4	0.09	0.0	0.01	0.1	0.01	0.2	0.03	2.1	0.07
Q4	29.5	3.23	2.9	0.10	0.0	0.01	0.1	0.02	0.4	0.05	2.5	0.08
Q5(richest)	26.4	3.76	3.4	0.10	0.0	0.01	0.1	0.02	0.5	0.06	2.8	0.07
All	33.0	2.04	2.3	0.04	0.0	0.00	0.1	0.01	0.2	0.01	2.0	0.03

Note: table based on de jure members of household i.e. usual residents

---

### 3.2 Selected age distributions by income quintile

---

	<b>Females Age 18–49</b>	<b>Persons Age 50+</b>	<b>Persons Age 60+</b>
Q1 (poorest)	56	450	400
Q2	94	390	285
Q3	116	291	178
Q4	171	200	83
Q5 (richest)	179	151	48
All	615	1,482	994

Note: table based on de jure members of household i.e. usual residents

---

### 3.3 Characteristics of head of household by income quintile

---

	<b>Mean Age of Head</b>	<b>SE</b>	<b>Median Education of Head</b>	<b>Percent Female Head</b>	<b>SE</b>
Q1 (poorest)	67	1.07	4	59.1	3.68
Q2	58	1.10	5	43.3	3.92
Q3	54	1.18	5	37.3	3.72
Q4	49	1.20	5	30.8	3.68
Q5 (richest)	46	1.01	5	21.7	2.81
All	57	0.59	5	41.9	1.78

#### Education

1=no formal education

2=less than primary school

3=primary school completed

4=secondary school completed

5=high school (or equivalent) completed

6=college/pre-university/university completed

7=post graduate degree completed

Note: table based on de jure members of household i.e. usual residents

### 3.4 Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex

	Male	SE	Female	SE	Missing	Total	SE
<b>Age</b>							
0-4	3.0	0.40	2.3	0.30		2.60	0.30
5-14	8.3	0.70	6.3	0.50		7.20	0.50
15-29	21.7	1.10	16.7	0.90		19.00	0.70
30-44	15.6	1.00	13.9	0.80		14.70	0.70
45-59	24.0	1.10	25.2	1.10		24.60	0.90
60-69	12.9	1.10	17.2	1.30		15.20	0.90
70-79	10.6	0.90	13.6	1.30		12.20	0.90
80+	4.0	0.70	4.7	0.70		4.40	0.50
Missing	0.0	0.00	0.1	0.10		0.10	0.00
<b>Residence</b>							
Urban	64.8	2.30	65.7	1.90		65.30	2.00
Rural	35.2	2.30	34.3	1.90		34.70	2.00
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Marital_Status</b>							
Never	36.5	1.40	24.9	1.20		30.20	0.90
Currently	48.0	1.30	41.7	1.20		44.60	1.10
Separated	0.7	0.30	1.1	0.40		0.90	0.20
Divorced	4.4	0.70	9.2	1.10		7.00	0.70
Widowed	5.7	0.90	19.3	1.40		13.00	0.80
CoHabiting	4.6	0.60	3.7	0.50		4.10	0.50
Missing	0.1	0.10	0.1	0.00		0.10	0.10
<b>Education</b>							
NoFormal	9.2	0.90	6.9	0.60		8.00	0.50
LessthanPrimary	1.1	0.30	1.4	0.30		1.30	0.20
Primary	5.4	0.70	9.1	0.90		7.40	0.60
Secondary	25.1	1.30	30.8	1.40		28.20	1.10
HighSchool	47.2	1.50	39.6	1.50		43.10	1.20
College	11.4	1.10	11.8	1.20		11.60	0.90
PostGraduate	0.5	0.20	0.2	0.10		0.40	0.10
Missing	0.1	0.10	0.1	0.10		0.10	0.10
<b>Total</b>	<b>100.0</b>		<b>100.0</b>			<b>100.00</b>	
Number	1,984.0		2,297.0			4,281.00	
Number of Households						1,412.00	

**3.5 Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education**

	Male	SE	Female	SE	Missing	Total	SE
<b>Age</b>							
18-29	24.7	2.10	20.9	1.80		22.70	1.30
30-44	27.4	2.20	23.6	1.60		25.40	1.30
45-59	27.2	2.00	26.3	1.70		26.70	1.30
60-69	11.2	1.30	13.5	1.30		12.40	0.90
70-79	7.3	1.00	11.1	1.10		9.30	0.70
80+	2.2	0.60	4.6	0.80		3.50	0.50
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Residence</b>							
Urban	65.3	2.60	64.6	2.40		64.90	2.00
Rural	34.7	2.60	35.4	2.40		35.10	2.00
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Marital_Status</b>							
Never	26.4	2.20	15.6	1.70		20.70	1.30
Currently	56.4	2.50	49.6	2.00		52.80	1.50
Separated	0.9	0.40	1.0	0.40		1.00	0.30
Divorced	5.0	1.00	10.2	1.40		7.80	0.90
Widowed	4.8	0.80	16.8	1.30		11.20	0.70
CoHabiting	6.4	1.20	6.9	1.10		6.70	0.90
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Education</b>							
NoFormal	0.3	0.20	0.2	0.10		0.20	0.10
LessthanPrimary	0.4	0.20	0.9	0.20		0.70	0.20
Primary	3.8	0.70	8.1	0.90		6.10	0.60
Secondary	18.3	1.70	31.4	1.90		25.30	1.40
HighSchool	63.9	2.30	45.5	2.10		54.10	1.60
College	12.2	1.50	13.2	1.40		12.70	1.10
PostGraduate	1.2	0.60	0.6	0.30		0.90	0.30
Missing	0.0	0.00	0.1	0.10		0.10	0.10
Total	100.0		100.0			100.00	
Number	663.0		756.0			1,419.00	

#### 4.1 Prevalence of smoking and average daily tobacco consumption

	Current daily smoker	SE	Smoker, not daily	SE	Non-smoker	SE	Missing	Average daily tobacco consumption*	SE	Number of respondents
<b>Sex</b>										
Male	38.6	2.20	3.9	0.90	57.1	2.10	0.5	18.5	0.80	663
Female	27.7	1.70	3.5	0.80	68.6	2.00	0.2	15.3	0.60	756
Missing										0
<b>Residence</b>										
Urban	33.0	1.80	4.0	0.80	63.0	1.80	0.0	16.8	0.70	921
Rural	32.3	2.60	3.0	0.70	63.7	2.50	1.0	17.7	0.50	498
Missing										0
<b>Income Quintile</b>										
Q1	33.7	3.30	2.8	1.20	62.8	3.50	0.8	19.7	1.20	238
Q2	38.9	3.40	3.1	1.20	58.0	3.70	0.0	18.2	0.90	259
Q3	34.7	3.50	3.1	1.00	61.9	3.30	0.3	17.5	1.00	264
Q4	30.9	3.00	2.4	1.10	66.6	3.00	0.0	16.3	0.80	293
Q5	25.1	3.10	6.5	1.90	67.7	3.10	0.7	14.1	1.10	311
Missing										54
<b>Age</b>										
18-29	39.0	3.50	7.1	1.70	53.9	3.60	0.0	15.2	0.80	322
30-44	42.8	3.20	3.7	1.10	53.0	3.10	0.6	18.0	0.80	360
45-59	36.3	2.70	2.7	0.80	60.5	2.70	0.4	18.7	0.90	379
60-69	17.3	2.60	1.9	0.90	80.2	2.80	0.6	15.1	1.60	177
70-79	13.2	3.00	1.8	1.30	85.1	3.20	0.0	13.0	2.70	132
80+	0.0	0.00	0.0	0.00	100.0	0.00	0.0			49
Missing										0
All	32.8	1.50	3.7	0.60	63.2	1.50	0.3	17.1	0.50	1,419

\* Denominator used is current daily smoker

#### 4.2 Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers

	Life time abstainers	SE	Non-heavy drinkers	SE	Infrequent Heavy Drinkers*	SE	Frequent Heavy drinkers**	SE	Missing	Number of respondents
<b>Sex</b>										
Male	3.6	0.90	78.5	1.90	13.2	1.40	4.8	0.90	0.0	663
Female	9.4	1.10	88.9	1.20	1.1	0.40	0.6	0.40	0.0	756
Missing										0
<b>Residence</b>										
Urban	6.0	1.10	84.2	1.40	7.4	0.90	2.5	0.60	0.0	921
Rural	7.9	1.40	83.7	2.10	5.6	1.20	2.8	0.80	0.0	498
Missing										0
<b>Income Quintile</b>										
Q1	10.9	2.20	80.0	3.00	3.9	1.40	5.2	1.60	0.0	238
Q2	7.0	1.90	83.5	2.50	6.1	1.70	3.4	1.00	0.0	259
Q3	7.3	1.90	84.5	2.60	5.7	1.60	2.5	0.90	0.0	264
Q4	5.4	1.50	85.7	2.40	7.2	1.80	1.7	0.90	0.0	293
Q5	3.4	1.40	85.9	2.50	9.4	1.90	1.3	0.70	0.0	311
Missing										54
<b>Age</b>										
18-29	5.8	1.60	80.9	2.70	11.5	2.00	1.8	1.10	0.0	322
30-44	6.0	1.50	86.3	2.20	5.7	1.30	1.9	0.80	0.0	360
45-59	3.7	1.00	86.3	2.00	8.3	1.70	1.7	0.60	0.0	379
60-69	7.4	2.20	84.2	2.80	1.5	0.90	6.9	1.90	0.0	177
70-79	14.5	3.10	78.4	3.50	3.3	1.30	3.9	1.60	0.0	132
80+	16.1	5.60	83.9	5.60	0.0	0.00	0.0	0.00	0.0	49
Missing										0
All	6.7	0.90	84.0	1.20	6.8	0.70	2.6	0.50	0.0	1,419

\*Infrequent heavy drinkers: 1-2 days with 5 or more standard drinks per week(last 7 days)

\*\*Frequent heavy drinkers: 3 or more days with 5 or more standard drinks per week(last 7 days)

### 4.3 Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

	Fruit and vegetable intake					Physical Activity					Number of respondents
	*Insufficient	SE	Sufficient	SE	Missing	**Insufficient	SE	Sufficient	SE	Missing	
<b>Sex</b>											
Male	83.3	2.00	16.7	2.00	0.0	8.7	1.70	91.3	1.70	0.0	663
Female	77.4	1.90	22.6	1.90	0.0	9.6	1.20	90.4	1.20	0.0	756
Missing											0
<b>Residence</b>											
Urban	79.6	2.10	20.4	2.10	0.0	8.9	1.20	91.1	1.20	0.0	921
Rural	81.0	2.30	19.0	2.30	0.0	9.7	1.80	90.3	1.80	0.0	498
Missing											0
<b>Income Quintile</b>											
Q1	86.5	2.00	13.5	2.00	0.0	21.1	2.80	78.9	2.80	0.0	238
Q2	83.4	2.30	16.6	2.30	0.0	11.0	2.30	89.0	2.30	0.0	259
Q3	80.5	2.80	19.5	2.80	0.0	4.9	1.50	95.1	1.50	0.0	264
Q4	74.5	3.40	25.5	3.40	0.0	4.5	1.40	95.5	1.40	0.0	293
Q5	78.8	3.00	21.2	3.00	0.0	6.4	2.00	93.6	2.00	0.0	311
Missing											54
<b>Age</b>											
18-29	74.8	3.30	25.2	3.30	0.0	5.3	1.60	94.7	1.60	0.0	322
30-44	82.6	2.70	17.4	2.70	0.0	6.5	1.90	93.5	1.90	0.0	360
45-59	80.1	2.10	19.9	2.10	0.0	5.6	1.20	94.4	1.20	0.0	379
60-69	80.7	3.00	19.3	3.00	0.0	10.7	2.50	89.3	2.50	0.0	177
70-79	85.1	3.10	14.9	3.10	0.0	24.7	3.80	75.3	3.80	0.0	132
80+	81.6	5.90	18.4	5.90	0.0	33.9	7.20	66.1	7.20	0.0	49
Missing											0
All	80.1	1.50	19.9	1.50	0.0	9.2	1.00	90.8	1.00	0.0	1,419

\*Insufficient intake of fruit/vegetables: less than 5 servings of fruit/vegetables per one typical day

\*\*Insufficient physical activity: less than 150 minutes per week(last 7 days) spent on walking/moderate activity/vigorous activity



#### 4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water

	Piped into households	SE	other improved sources	SE	Total improved sources	SE	No improved sources	SE	Missing	Number of households
<b>Residence</b>										
Urban	95.8	1.00	4.2	1.00	100.0	0.00	0.0	0.00	0.0	946
Rural	88.2	2.00	11.7	2.00	99.9	0.10	0.1	0.10	0.0	466
Missing										0
<b>Income_Quintile</b>										
Q1(poorest)	81.3	3.00	18.6	3.00	99.9	0.10	0.1	0.10	0.0	389
Q2	96.9	1.10	3.1	1.10	100.0	0.00	0.0	0.00	0.0	308
Q3	97.7	0.80	2.3	0.80	100.0	0.00	0.0	0.00	0.0	251
Q4	98.3	0.80	1.7	0.80	100.0	0.00	0.0	0.00	0.0	216
Q5(richest)	99.0	0.60	1.0	0.60	100.0	0.00	0.0	0.00	0.0	185
Missing										63
Total	93.3	0.90	6.7	0.90	100.0	0.00	0.0	0.00	0.0	1,412

other improved sources: public standpipe/protected tube well/borehole/protected dug well/protected spring/rainwater collection

no improved sources: unprotected well/unprotected spring/water taken directly from a pond or stream/tanker truck water/vendor provided water

#### 4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility

	Flush toilet to sewage system	SE	Other improved toilet facilities	SE	Total improved sanitation	SE	No improved sanitation	SE	Missing	Number of households
<b>Residence</b>										
Urban	95.4	1.00	4.6	1.00	100.0	0.00	0.0	0.00	0.0	946
Rural	80.8	2.00	19.2	2.00	100.0	0.00	0.0	0.00	0.0	466
Missing										0
<b>Income_Quintile</b>										
Q1(poorest)	70.5	3.10	29.5	3.10	100.0	0.00	0.0	0.00	0.0	389
Q2	97.0	1.00	3.0	1.00	100.0	0.00	0.0	0.00	0.0	308
Q3	98.3	0.80	1.7	0.80	100.0	0.00	0.0	0.00	0.0	251
Q4	100.0	0.00	0.0	0.00	100.0	0.00	0.0	0.00	0.0	216
Q5(richest)	100.0	0.00	0.0	0.00	100.0	0.00	0.0	0.00	0.0	185
Missing										63
Total	90.6	1.00	9.4	1.00	100.0	0.00	0.0	0.00	0.0	1,412

other improved sanitation facilities: connection to septic system/pour-flush latrine/covered dry latrine(with privacy) [provided facilities not shared]  
no improved sanitation: uncovered dry latrine(without privacy)/bucket latrine/no facilities(open defecation)



## 5.01 Tuberculosis

	Total	Tuberculosis			
		Need	Coverage	SE	M
<b>Sex</b>					
Female	756	15.6	76.6	4.3	0.0
Male	663	13.9	83.5	3.9	0.0
Missing	0				
<b>Residence</b>					
Urban	921	14.0	80.5	4.2	0.0
Rural	498	16.3	78.3	4.1	0.0
Missing	0				
<b>Income_Quintile</b>					
Q1(poorest)	238	18.4	73.8	6.5	0.0
Q2	259	16.9	80.8	6.2	0.0
Q3	264	13.8	78.4	6.0	0.0
Q4	293	14.8	87.9	5.0	0.0
Q5(richest)	311	11.9	82.3	6.2	0.0
Missing	54				
<b>Age</b>					
18-29	322	10.3	64.9	10.5	0.0
30-44	360	12.5	85.6	4.9	0.0
45-59	379	19.8	86.6	3.8	0.0
60-69	177	19.7	79.0	6.8	0.0
70-79	132	13.8	70.4	9.2	0.0
80+	49	8.3	53.0	22.1	0.0
Missing	0				
Total	1,419	14.8	79.6	3.0	0.0

**M = % Missing of Covered      SE= Standard Error of % Covered**

**Tuberculosis Indicators:**

**"Need" = % of all respondents with self-reported continuous cough lasting 3 weeks, OR had blood in phlegm, OR had coughed blood in the last 12 months**

**"Coverage" = % of respondents with self reported symptoms who indicated receiving a TB screening test in the last 12 months.**

## 5.02 HIV Prevention and Screening

	Total (persons of reproductive age 18-49)	Voluntary Counselling and Testing for HIV/AIDS				Condom Usage			
		Need	Coverage	SE	M	Need	Coverage	SE	M
<b>Sex</b>									
Females	407	53.3	2.6	1.0	0.0	13.8	40.6	7.3	0.0
Males	406	62.2	1.7	0.9	40.6	24.1	39.4	6.0	0.0
Missing									
<b>Residence</b>									
Urban	518	57.9	2.0	0.9	19.3	20.5	38.1	6.1	0.0
Rural	295	57.4	2.3	1.0	26.4	16.3	43.6	6.6	0.0
Missing									
<b>Income_Quintile</b>									
Q1(poorest)	74	48.3	2.1	2.1	28.1	14.7	8.6	8.3	0.0
Q2	117	46.2	5.1	3.0	25.5	12.8	43.4	15.4	0.0
Q3	145	59.6	1.4	1.2	17.6	21.5	36.7	9.8	0.0
Q4	206	59.2	1.8	1.1	21.0	25.6	34.0	8.5	0.0
Q5(richest)	242	62.6	2.0	1.4	19.2	17.4	55.5	9.9	0.0
Missing	28								
Total	813	57.7	2.1	0.7	21.9	18.9	39.8	4.7	0.0

**M = % Missing of Covered      SE= Standard Error of % Covered**

**HIV Prevention and Screening Indicators:**

**"Total" = number of respondents of reproductive age.**

**"Voluntary Counselling and Testing for HIV/AIDS Need" = % of respondents of reproductive age who are sexually active, i.e. have had sexual intercourse in the last 12 months.**

**"Voluntary Counselling and Testing for HIV/AIDS Coverage" = % of respondents of reproductive age who are sexually active who indicated having taken an HIV test and receiving the results of test.**

**"Condom Usage Need" = % of respondents of reproductive age whose last sexual encounter during the last 12 months is classified as risky sexual behavior, i.e. having had sexual intercourse with a non-cohabitating partner .**

**"Condom Usage Coverage" = % of respondents of reproductive age who used a condom last time had risky sex during the last 12 months (with non-cohabitating partner).**

### 5.03 Antenatal Care and Mother to Child Transmission of HIV

	Antenatal Care											Maternal to Child HIV Trans.			
	Women 18-49	Birth in last 5 years	Cov	SE	Visit 1x	Visit 2x	Visit 3x	Blood pressure checked	Blood sample taken	Told Signs of Pregnancy Complications	M	Birth in last 2 years	Cov	SE	M
<b>Residence</b>															
Urban	257	18.5	69.1	7.7	98.5	98.5	96.2	100.0	98.4	71.5	5.0	10.0	11.7	6.3	0.0
Rural	151	18.5	74.4	7.3	98.3	98.3	91.7	98.3	98.3	81.1	1.7	14.4	0.0	0.0	2.5
Missing															
<b>Income_Quintile</b>															
Q1(poorest)	40	49.1	80.8	8.5	92.8	92.8	92.8	96.3	96.3	80.8	3.5	21.8	12.3	11.9	0.0
Q2	61	26.8	58.5	12.8	100.0	100.0	100.0	100.0	95.3	58.5	0.0	23.2	8.4	8.0	0.0
Q3	72	23.1	65.1	13.3	100.0	100.0	93.2	100.0	100.0	71.8	4.4	11.0	0.0	0.0	6.9
Q4	113	16.5	68.9	13.4	100.0	100.0	85.0	100.0	100.0	83.9	0.0	6.9	0.0	0.0	0.0
Q5(richest)	107	17.0	81.8	10.3	100.0	100.0	100.0	100.0	100.0	81.8	9.3	7.8	8.7	8.6	0.0
Missing	14														
Total	407	22.1	71.6	5.3	98.4	98.4	94.1	99.2	98.3	76.0	3.5	11.6	6.3	3.7	1.2

M = % Missing of Covered SE= Standard Error of %Covered

#### Antenatal Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least 3 times or more by health care professional, received blood pressure, blood sample taken, and told signs about complications for their last pregnancy (aggregate antenatal care indicator).

"Visit 1x" "Visit 2x" "Visit 3x" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least once, twice or three times or more respectively by a health care professional during their last pregnancy.

"Blood pressure checked" = % of female respondents of reproductive age with live birth in the last 5 years who had blood pressure taken when they saw health care professional during their last pregnancy.

"Blood sample taken" = % of female respondents of reproductive age with live birth in the last 5 years who had a blood sample taken when they saw health care professional during their last pregnancy.

"Told Signs of Pregnancy Complications" = % of female respondents of reproductive age with live birth in the last 5 years who had been told signs of pregnancy complications when they saw health care professional during their last pregnancy.

#### Maternal to Child HIV Transmission Indicators:

"Birth in last 2 years" = % of female respondents of reproductive age indicating having had a live birth in the last 2 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 2 years who indicated having been counseled and tested and given results for HIV during their last pregnancy.

## 5.04 Birth Delivery Care

	Women 18-49	Delivery Care				Place of Delivery						
		Need	Coverage	SE	M	Public Health Facility	SE	Private Health Facility	SE	Home	SE	M
<b>Residence</b>												
Urban	257	18.5	88.1	5.5	0.0	97.3	2.7	0.0	0.0	0.0	0.0	0.0
Rural	151	18.5	88.4	4.9	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Missing												
<b>Income_Quintile</b>												
Q1(poorest)	40	49.1	94.5	5.3	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Q2	61	26.8	80.1	9.5	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Q3	72	23.1	83.1	9.3	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Q4	113	16.5	90.2	6.8	0.0	93.1	6.7	0.0	0.0	0.0	0.0	0.0
Q5(richest)	107	17.0	91.2	8.4	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Missing	14											
Total	407	22.1	88.2	3.7	0.0	98.6	1.4	0.0	0.0	0.0	0.0	0.0

M = % Missing of Covered SE= Standard Error of % Covered Birth Delivery Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Coverage" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having received assistance with birth delivery from a health care professional (doctor, nurse or midwife) during their last birth.

"Public Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a public health facility.

"Private Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a private health facility.

"Home" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered home or in the field (etc.)

## 5.05 Child's Health - Common Illness

	Total Children under Age of 5	Fever						Cough						Diarrhea					
		More liquid	SE	Same or more food	SE	M	More liquid	SE	Same or more food	SE	M	More liquid	SE	Same or more food	SE	M			
<b>Sex</b>																			
Females	83	40.3	82.8	6.5	58.2	11.0	0.0	21.6	66.4	11.7	43.8	40.3	0.0	11.7	82.5	15.7	70.1	17.7	0.0
Males	81	41.0	71.2	7.2	52.1	10.2	0.0	42.9	71.0	7.4	43.3	41.0	0.0	8.6	83.1	15.1	72.5	16.4	0.0
Missing	0																		
<b>Residence</b>																			
Urban	82	44.9	75.0	6.3	56.4	9.5	0.0	34.5	68.5	8.9	42.5	44.9	0.0	12.2	83.2	15.2	60.6	17.9	0.0
Rural	82	36.4	79.5	7.7	53.6	11.5	0.0	29.8	70.5	7.2	44.6	36.4	0.0	8.1	82.1	15.9	87.0	12.2	0.0
Missing																			
<b>Income_Quintile</b>																			
Q1(poorest)	26	34.2	90.8	9.0	32.3	18.0	0.0	35.7	59.5	14.5	18.2	34.2	0.0	10.5	100.0	0.0	60.0	31.2	0.0
Q2	30	50.2	58.8	14.5	44.5	14.4	0.0	38.6	65.2	16.4	35.2	50.2	0.0	14.7	61.2	27.6	33.8	22.6	0.0
Q3	31	28.9	58.2	18.8	61.6	17.8	0.0	25.7	49.8	19.7	68.9	28.9	0.0	3.7	100.0	0.0	100.0	0.0	0.0
Q4	36	40.4	79.5	10.6	54.0	15.2	0.0	28.1	72.7	13.2	45.0	40.4	0.0	12.3	72.7	22.3	80.3	17.9	0.0
Q5(richest)	41	45.9	96.2	3.8	69.7	10.9	0.0	32.4	94.6	5.4	47.2	45.9	0.0	10.1	100.0	0.0	100.0	0.0	0.0
Missing	1																		
Total	164	40.7	77.0	4.9	55.1	7.4	0.0	32.2	69.4	5.8	43.5	40.7	0.0	10.2	82.7	11.1	71.1	12.4	0.0

**M = % Missing of Covered      SE= Standard Error of % Covered**

### **Child's Common Illness Indicators:**

**Total = Number of children under the age of 5 years at the time of the interview.**

**"Fever"/"Cough"/"Diarrhea" = % of children under the age of 5 with reported episode of (cough, fever, or diarrhea) in the last 3 months.**

**"More Liquid" = % of children under 5 with either fever, cough, or diarrhea in the last 3 months, that were given more liquids to drink during their illness.**

**"Same or More Food" = % of children under 5 with either fever, cough, or diarrhea in the last 3 months, that were given the same amount or more food to eat during their illness.**

## 5.06 Child's Health - Danger Signs

	Children under 5	Fast Breath	Seeking Treatment						Vomit	Seeking Treatment						No Drink or Eat	Seeking Treatment					
			Yes	SE	Public	Private	Trad	M		Yes	SE	Public	Private	Trad	M		Yes	SE	Public	Private	Trad	M
<b>Sex</b>																						
Females	83	0.1	100.0	0.0	100.0	0.0	0.0	0.0	8.3	0.0	0.0	100.0	0.0	0.0	0.0	16.1	100.0	0.0	91.4	0.0	8.6	0.0
Males	81	0.2	91.4	5.9	100.0	0.0	0.0	0.0	14.8	0.0	0.0	100.0	0.0	0.0	0.0	14.9	97.4	2.7	100.0	0.0	0.0	0.0
Missing	0																					
<b>Residence</b>																						
Urban	82	0.1	90.4	6.5	100.0	0.0	0.0	0.0	9.7	0.0	0.0	100.0	0.0	0.0	0.0	15.3	97.5	2.6	90.7	0.0	9.3	0.0
Rural	82	0.1	100.0	0.0	100.0	0.0	0.0	0.0	13.4	0.0	0.0	100.0	0.0	0.0	0.0	15.7	100.0	0.0	100.0	0.0	0.0	0.0
Missing																						
<b>Income_Quintile</b>																						
Q1(poorest)	26	0.1	87.4	13.5	100.0	0.0	0.0	0.0	24.0	0.0	0.0	100.0	0.0	0.0	0.0	20.1	93.9	6.7	100.0	0.0	0.0	0.0
Q2	30	0.2	85.2	13.9	100.0	0.0	0.0	0.0	21.0	0.0	0.0	100.0	0.0	0.0	0.0	17.2	100.0	0.0	100.0	0.0	0.0	0.0
Q3	31	0.1	100.0	0.0	100.0	0.0	0.0	0.0	5.6	0.0	0.0	100.0	0.0	0.0	0.0	15.8	100.0	0.0	76.7	0.0	23.3	0.0
Q4	36	0.1	100.0	0.0	100.0	0.0	0.0	0.0	8.7	0.0	0.0	100.0	0.0	0.0	0.0	10.7	100.0	0.0	100.0	0.0	0.0	0.0
Q5(richest)	41	0.1	100.0	0.0	100.0	0.0	0.0	0.0	4.3	0.0	0.0	100.0	0.0	0.0	0.0	16.0	100.0	0.0	100.0	0.0	0.0	0.0
Missing	1																					
Total	164	0.1	94.6	3.9	100.0	0.0	0.0	0.0	11.5	0.0	0.0	100.0	0.0	0.0	0.0	15.5	98.8	1.3	95.4	0.0	4.6	0.0

**M = % Missing of those Seeking Treatment      SE= Standard Error of % Seeking Treatment**

**Child Danger Signs Indicators:**

**Total = Number of children under the age of 5 years at the time of the interview.**

**"Fast Breathing"/"Vomiting"/"No Drink or Eat" = % of children under the age of 5 with reported episodes of (fast breathing, vomiting, or no eating or drinking) in the last 3 months.**

**"Seeking Treatment" = % of children under 5 with episodes of fast breathing, vomiting, or no eating or drinking in the last 3 months that were taken for care/treatment at either a private health facility, public health facility, or traditional healer (other).**

**Note: large number of missings are usually due to not having information type of facility (either a no answer or a don't know answer).**





## 5.09 Women's Cancer Screening

	Total Females	Percent Females 18+	PAP Smear	SE	M	Percent Females 40+	Mammography / Breast Exam	SE	M
<b>Residence</b>									
Urban	488	84.8	62.4	2.4	0.5	50.2	57.8	3.1	0.6
Rural	267	83.5	60.1	3.0	2.8	45.4	56.1	4.2	1.0
Missing	0								
<b>Income_Quintile</b>									
Q1(poorest)	155	56.2	44.3	5.0	3.2	36.2	27.7	5.1	2.4
Q2	141	84.6	52.5	5.3	2.3	60.8	47.7	5.1	0.0
Q3	136	89.2	63.8	3.9	0.9	51.9	69.4	5.0	0.0
Q4	158	94.0	65.0	4.3	0.6	43.2	69.1	5.6	1.8
Q5(richest)	138	98.9	72.9	4.0	0.7	50.5	64.1	6.3	0.0
Missing	28								
<b>Age</b>									
18-29	158	20.9	63.4	4.4	0.6				
30-39	113	14.9	76.3	4.2	2.7				
40-49	137	18.1	70.0	3.5	0.6	18.1	48.4	4.3	0.9
50-59	127	16.8	58.4	4.0	0.8	16.8	72.7	3.7	0.0
60-69	102	13.5	35.4	4.2	2.4	13.5	49.8	3.9	1.3
70+	119	15.7	0.0	0.0		15.7	0.0	0.0	
Missing	0								
Total	756	84.3	61.6	1.9	1.3	48.5	57.2	2.4	0.7

M = % Missing of those with Pap Smear/Mammography SE= Standard Error of % with Pap Smear/Mammography  
 \*\*\*Note: Values in the age row under the "Females 18 plus" and "Females 40 plus" columns are column percent values.

### Women's Cancer Screening Indicators:

"Papsmear need" = % of all female respondents aged 18-69

"Papsmear" coverage = % of those female respondents aged 18-69 who self-reported receiving a pap smear test during pelvic examination in the last 3 years.

"Mammography need" = % of all female respondents aged 40-69

"Mammography" coverage = % of those female respondents aged 40-69 who self-reported receiving a mammography test or breast examination in the last 3 years.

\*\*\*\*In several countries women above age 69 were asked the cancer screening questions, and for this reason they are included in the table. For countries that did not ask this questions of women above age 69, these values are expected to be missing.

## 5.10 Oral Health and Injuries

	Total	Oral Health Problems				Road Traffic Accidents				Injury			
		Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
<b>Sex</b>													
Female	756	36.2	80.8	2.7	0.0	2.9	10.8	7.0	17.4	7.6	15.0	4.9	4.0
Male	663	36.8	73.7	2.9	0.0	2.2	37.8	12.5	0.0	8.5	18.9	7.1	9.3
Missing	0												
<b>Residence</b>													
Urban	921	38.4	77.2	2.7	0.0	2.8	9.0	5.6	14.7	7.8	8.3	3.2	6.5
Rural	498	33.0	78.1	3.1	0.0	2.1	52.7	15.8	0.0	8.4	31.7	10.7	6.7
Missing	0												
<b>Income_Quintile</b>													
Q1(poorest)	238	28.4	61.1	6.3	0.0	2.6	40.1	21.4	21.6	7.5	26.1	11.1	5.7
Q2	259	26.3	69.2	6.0	0.0	2.1	0.0	0.0	15.5	9.8	11.5	5.8	4.7
Q3	264	34.6	76.1	5.5	0.0	1.4	19.8	14.1	0.0	7.8	17.4	8.2	1.3
Q4	293	44.8	81.1	3.9	0.0	5.1	19.9	11.2	0.0	9.2	7.5	4.2	7.3
Q5(richest)	311	42.7	85.4	3.6	0.0	1.9	26.9	17.3	26.9	5.8	33.7	21.9	0.0
Missing	54												
<b>Age</b>													
18-29	322	48.1	84.2	3.8	0.0	2.7	0.0	16.3	0.0	8.2	3.6	0.0	3.6
30-44	360	41.9	78.7	3.8	0.0	2.6	17.4	16.3	17.4	7.9	0.0	11.5	0.0
45-59	379	36.9	75.9	4.4	0.0	2.1	0.0	14.9	0.0	8.9	14.9	7.8	14.9
60-69	177	24.9	74.0	6.0	0.0	4.7	26.5	3.5	26.5	6.2	9.3	8.4	9.3
70-79	132	16.5	53.9	11.0	0.0	1.2	0.0	0.0	0.0	7.3	5.3	11.8	5.3
80+	49	12.5	21.7	13.1	0.0	0.5	0.0	0.0	0.0	9.7	0.0	0.0	0.0
Missing	0												
Total	1,419	36.5	77.5	2.1	0.0	2.6	21.6	6.9	10.4	8.0	16.9	5.0	6.6

M = % Missing of % Covered    SE= Standard Error of % Covered

### Indicators:

#### Oral Health:

"Need" = % of all respondents indicating any problems in the last 12 months with their mouth and/or teeth.

"Coverage" = % of respondents reporting problems with their mouth and teeth who indicated receiving dental or medical care in last 12 months.

#### Road Traffic Accident:

"Need" = % of all respondents indicating having suffered bodily injury from a road traffic accident in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury from a road traffic accident in last 12 months who indicating having received treatment for their injuries on-site or at health facility within 1 hour of accident .

#### Injuries from Non-traffic Accidents:

"Need" = % of all respondents indicating having suffered bodily injury due to non-traffic accidents in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury due to non-traffic accidents in last 12 months who indicating having received treatment for injuries on-site or at health facility within 1 hour of accident

## 5.11 Self Reported Non-Communicable Conditions

	Total	Angina				Osteo-Arthritis				Asthma				Diabetes			
		Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
<b>Sex</b>																	
Female	756	20.4	63.0	4.1	1.1	28.7	42.4	3.7	0.4	6.1	38.7	7.8	0.0	10.5	71.5	5.6	0.0
Male	663	12.3	73.5	6.7	0.0	21.5	34.1	4.5	0.0	7.7	29.3	7.1	0.0	6.8	79.2	8.3	0.0
Missing	0																
<b>Residence</b>																	
Urban	921	15.6	64.4	4.4	0.5	23.7	36.9	3.5	0.4	7.5	23.9	5.9	0.0	9.1	73.8	6.5	0.0
Rural	498	18.4	70.3	6.1	1.0	28.4	42.6	4.1	0.0	5.4	59.1	8.8	0.0	8.2	75.4	7.4	0.0
Missing	0																
<b>Income_Quintile</b>																	
Q1(poorest)	238	31.9	66.5	4.7	1.0	35.0	59.9	4.8	0.0	4.7	73.5	12.4	0.0	11.0	68.6	9.4	0.0
Q2	259	16.3	68.4	7.2	2.1	24.7	35.2	5.3	0.0	7.0	31.2	11.3	0.0	11.3	78.4	6.8	0.0
Q3	264	14.6	73.7	6.7	0.0	28.0	36.0	5.4	0.0	6.7	42.3	13.4	0.0	11.8	68.4	11.4	0.0
Q4	293	15.1	58.1	8.8	0.0	21.1	33.6	7.1	1.2	8.0	24.3	10.6	0.0	6.5	88.1	8.0	0.0
Q5(richest)	311	8.2	70.4	9.2	0.0	20.1	23.7	5.1	0.0	7.0	25.7	9.7	0.0	4.5	72.0	13.3	0.0
Missing	54																
<b>Age</b>																	
18-29	322	3.0	32.3	18.8	0.0	4.6	29.0	12.7	0.0	9.9	22.3	10.7	0.0	0.9	0.0	0.0	0.0
30-44	360	7.8	45.7	12.7	0.0	18.0	25.4	5.8	0.0	5.7	45.4	8.8	0.0	4.2	74.9	12.1	0.0
45-59	379	22.3	69.8	5.4	0.0	35.0	38.3	5.1	0.0	6.5	60.5	13.6	0.0	10.1	75.6	8.9	0.0
60-69	177	29.8	74.6	5.8	0.0	40.8	42.3	5.7	0.0	7.1	34.0	14.2		19.2	75.8	6.9	0.0
70-79	132	32.0	61.6	7.5	3.9	45.1	47.6	6.5	1.3	4.9	0.0	0.0		19.4	83.8	6.8	0.0
80+	49	36.3	92.1	5.3	0.0	32.2	66.0	12.0	0.0	0.0	0.0	0.0		17.6	57.6	18.7	0.0
Missing	0																
Total	1,419	16.6	66.7	3.6	0.7	25.3	39.1	2.7	0.2	6.8	33.7	5.3	0.0	8.8	74.3	5.0	0.0

**M = % Missing for Covered**

**SE= Standard Error of %Covered**

**Self Reported Non-Communicable Conditions Indicators:**

"Need" = % of total respondents indicating having received a diagnosis of the condition (angina, arthritis, asthma, diabetes) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (angina, arthritis, asthma, diabetes) indicating they have received treatment or screening for the condition in the last 2 weeks.

## 5.12 Mental Health Conditions

	Total	Depression				Schizophrenia				
		Need	Coverage	SE	M	Need	Coverage	SE	M	
<b>Sex</b>										
Female	756	11.7	54.4	5.5	0.6	2.6	35.4	11.2	0.0	
Male	663	4.6	43.3	10.6	0.0	2.3	43.7	14.5	0.0	
Missing	0									
<b>Residence</b>										
Urban	921	7.9	52.9	7.1	0.0	2.0	43.2	14.7	0.0	
Rural	498	9.2	49.4	7.7	1.1	3.2	34.2	10.7	0.0	
Missing	0									
<b>Income Quintile</b>										
Q1(poorest)	238	12.5	48.2	9.8	1.6	5.7	52.2	15.8	0.0	
Q2	259	9.1	83.7	6.6	0.0	2.3	84.1	14.9	0.0	
Q3	264	6.4	37.5	10.2	0.0	1.8	9.1	9.1	0.0	
Q4	293	8.7	28.3	9.8	0.0	1.6	22.7	14.6	0.0	
Q5(richest)	311	5.2	55.8	14.1	0.0	1.2	0.0	0.0	0.0	
Missing	54									
<b>Age</b>										
18-29	322	2.5	14.4	13.3	0.0	2.7	0.0	0.0	0.0	
30-44	360	8.5	36.7	10.5	0.0	2.1	33.9	21.2	0.0	
45-59	379	13.6	57.5	7.9	0.0	2.2	48.2	17.8	0.0	
60-69	177	9.0	77.9	9.1	3.1	5.3	68.7	14.3	0.0	
70-79	132	8.5	54.4	13.8	0.0	0.3	100.0	0.0	0.0	
80+	49	3.2	48.1	35.3	0.0	0.0	0.0	0.0		
Missing	0									
Total	1,419	8.4	51.5	5.4	0.4	2.4	39.0		0.0	

**M = % Missing for Covered**

**SE= Standard Error of %Covered**

**Mental Health Conditions Indicators:**

"Need" = % of total respondents indicating having received a diagnosis of the condition (depression, schizophrenia) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (depression, schizophrenia) indicating they have received treatment or screening for the condition in the last 2 weeks.

## 5.13 Cataracts

	Total 60+	Cataracts			M
		Need	Coverage	SE	
<b>Sex</b>					
Female	221	14.3	30.4	8.1	0.0
Male	137	13.2	50.2	11.1	0.0
Missing	0				
<b>Residence</b>					
Urban	230	15.8	34.1	7.5	0.0
Rural	128	10.3	47.3	11.5	0.0
Missing	0				
<b>Income_Quintile</b>					
Q1(poorest)	138	13.9	30.3	9.2	0.0
Q2	98	9.0	59.7	13.3	0.0
Q3	61	10.9	15.9	14.8	0.0
Q4	33	17.9	37.3	15.5	0.0
Q5(richest)	18	21.5	61.1	27.5	0.0
Missing	10				
<b>Age</b>					
60-69	177	9.2	33.5	10.7	0.0
70-79	132	19.3	38.0	8.7	0.0
80+	49	15.9	45.1	19.0	0.0
Missing	0				
Total	358	13.9	37.6	6.4	0.0

**M = % Missing of Covered SE= Standard Error of %Covered**

**Cataracts Indicators:**

**Total = Number of respondents 60 years of age or more.**

**"Need" = % in need (respondents 60 years of age or more with self-reported diagnosis of cataracts in the last 5 years).**

**"Coverage" = % respondents 60 years of age or more with self-reported diagnosis of cataracts having indicated receiving eye surgery to remove cataracts in the last 5 years)**

**6.01 In general, rate your health today - q2000**

	Very Good	SE	Good	SE	Moderate	SE	Bad	SE	Very Bad	SE	Missing	SE	N
<b>Gender</b>													
Male	14.4	2.0	44.3	2.2	30.3	1.8	9.2	1.1	1.6	0.5	0.2	0.2	663
Female	9.7	1.1	42.0	1.6	35.5	1.6	9.4	1.0	2.1	0.5	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	12.5	1.5	44.3	1.7	32.6	1.5	8.4	1.0	1.2	0.4	0.9	0.5	921
Rural	10.7	1.5	40.7	2.1	34.1	1.9	10.9	1.3	3.1	0.6	0.5	0.3	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	3.9	1.3	30.2	3.0	41.7	3.2	19.0	2.5	5.3	1.1	0.0	0.0	238
Q2	14.2	2.4	33.1	3.1	38.1	3.5	11.3	1.9	2.7	1.0	0.6	0.4	259
Q3	10.6	2.2	45.0	3.3	34.1	2.7	9.1	2.0	1.1	0.6	0.0	0.0	264
Q4	12.6	2.5	51.0	3.4	26.5	2.8	7.2	1.6	1.0	0.7	1.6	0.9	293
Q5(richest)	15.0	2.7	52.8	3.3	28.6	3.1	2.6	1.0	0.1	0.1	0.9	0.7	311
M	20.8	8.0	39.3	7.8	27.2	9.4	7.3	3.9	1.8	1.7	3.6	3.6	54
<b>Age</b>													
18-29	30.4	3.1	50.8	3.2	15.0	2.5	1.3	0.7	0.3	0.3	2.2	1.4	322
30-44	12.9	3.0	58.9	3.0	25.1	2.7	2.8	1.0	0.4	0.3	0.0	0.0	360
45-59	2.9	0.8	39.3	2.7	40.9	3.0	14.3	2.0	2.5	0.7	0.0	0.0	379
60-69	2.5	1.0	24.2	2.9	51.6	3.5	15.6	2.4	4.5	1.6	1.7	0.9	177
70-79	4.8	1.9	25.7	3.9	45.9	3.8	18.4	3.3	5.3	1.8	0.0	0.0	132
80+	6.1	3.3	20.2	5.3	48.4	7.5	23.3	6.2	0.0	0.0	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	11.9	1.1	43.1	1.3	33.1	1.1	9.3	0.8	1.9	0.3	0.8	0.3	1,419

**6.02 Difficulty with 'work and household activities' in last 30 days - q2001**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	66.1	2.1	14.7	1.6	12.1	1.5	5.0	0.9	1.8	0.6	0.2	0.2	663
Female	56.5	1.8	20.3	1.4	13.6	1.3	6.8	0.9	1.6	0.4	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	63.2	1.9	18.6	1.4	11.2	1.3	4.8	0.8	1.3	0.4	0.9	0.5	921
Rural	56.9	2.3	15.9	1.8	16.2	1.9	8.1	1.2	2.4	0.7	0.5	0.3	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	44.5	3.6	17.8	2.4	18.5	2.6	14.3	2.2	4.9	1.2	0.0	0.0	238
Q2	56.6	3.1	18.6	2.1	14.9	2.2	8.0	1.7	1.3	0.8	0.6	0.4	259
Q3	61.8	3.2	18.8	2.6	13.6	2.4	3.7	1.3	2.0	0.8	0.0	0.0	264
Q4	66.8	3.2	15.9	2.4	11.1	2.3	4.0	1.2	0.6	0.4	1.6	0.9	293
Q5(richest)	69.8	2.9	17.6	2.6	9.6	1.9	1.8	0.8	0.3	0.3	0.9	0.7	311
M	68.0	7.2	16.5	5.8	5.1	3.0	4.6	2.5	2.3	2.2	3.6	3.6	54
<b>Age</b>													
18-29	80.7	2.6	9.6	1.8	6.2	1.9	1.2	0.6	0.0	0.0	2.2	1.4	322
30-44	72.9	3.2	15.2	2.4	8.3	1.6	3.3	1.2	0.3	0.3	0.0	0.0	360
45-59	50.8	3.1	23.5	2.8	17.4	2.5	7.1	1.3	1.2	0.6	0.0	0.0	379
60-69	45.3	4.1	20.1	2.9	20.5	3.4	7.9	2.4	4.4	1.6	1.7	0.9	177
70-79	42.5	3.9	21.4	2.9	15.8	2.8	14.5	2.5	5.8	1.9	0.0	0.0	132
80+	29.2	6.8	24.1	6.4	21.5	6.8	17.1	5.3	6.1	2.8	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	61.0	1.4	17.7	1.1	12.9	1.1	6.0	0.7	1.7	0.3	0.8	0.3	1,419

### 6.03 Difficulty with 'moving around' in last 30 days - q2010

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	77.0	1.6	7.8	1.1	8.0	1.3	5.0	0.8	1.6	0.6	0.5	0.4	663
Female	64.3	1.5	12.9	1.2	10.7	1.0	9.3	1.0	1.6	0.4	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	72.5	1.4	10.5	1.0	8.1	1.0	6.6	0.8	1.4	0.4	0.9	0.5	921
Rural	66.1	1.7	10.6	1.5	11.8	1.5	8.5	1.1	2.1	0.8	0.9	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	46.8	3.5	12.9	2.1	17.5	2.5	17.2	2.3	5.7	1.4	0.0	0.0	238
Q2	62.8	3.3	12.5	2.0	13.3	2.0	9.1	1.6	1.7	0.9	0.6	0.4	259
Q3	70.3	2.5	13.4	2.1	8.9	1.7	6.1	1.6	1.3	0.7	0.0	0.0	264
Q4	78.7	2.9	8.1	1.9	5.7	1.6	4.6	1.1	0.6	0.4	2.3	1.1	293
Q5(richest)	85.1	2.2	6.7	1.5	5.3	1.7	2.1	0.8	0.0	0.0	0.9	0.7	311
M	78.1	6.4	11.5	6.7	2.3	1.7	4.6	2.5	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	93.2	1.7	2.8	0.9	1.3	0.6	0.5	0.4	0.0	0.0	2.2	1.4	322
30-44	88.3	2.1	4.5	1.2	5.1	1.7	1.6	0.7	0.0	0.0	0.6	0.6	360
45-59	62.0	2.3	17.7	2.2	11.4	1.7	8.0	1.4	0.9	0.5	0.0	0.0	379
60-69	49.0	3.8	14.4	2.6	20.6	2.8	10.2	2.0	4.1	1.7	1.7	0.9	177
70-79	37.4	3.8	18.6	3.0	11.9	2.7	24.7	3.3	7.4	2.6	0.0	0.0	132
80+	16.8	5.4	13.9	4.7	33.5	7.5	28.9	6.7	4.9	3.3	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	70.3	1.1	10.5	0.8	9.4	0.8	7.2	0.6	1.6	0.4	0.9	0.4	1,419

## 6.04 Difficulty with 'vigorous activities' in last 30 days - q2011

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	51.0	2.4	6.4	1.0	8.1	1.2	6.9	1.0	10.5	1.3	17.1	2.1	663
Female	33.1	1.9	11.8	1.1	8.4	1.2	7.9	1.0	15.1	1.4	23.8	2.0	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	42.1	2.2	9.6	1.1	7.5	1.0	6.8	0.9	11.8	1.3	22.2	2.2	921
Rural	40.2	2.5	8.6	1.2	9.6	1.4	8.6	1.0	15.1	1.7	17.9	3.1	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	19.6	2.6	10.8	2.4	6.8	1.4	9.0	1.8	27.2	3.3	26.5	3.5	238
Q2	34.5	3.2	7.8	1.7	6.1	1.5	10.0	1.7	15.4	2.4	26.1	3.7	259
Q3	45.7	3.6	7.1	1.7	10.3	1.9	7.5	1.7	11.9	1.8	17.5	2.8	264
Q4	46.8	3.6	11.6	2.1	7.9	1.5	7.1	1.6	10.0	2.0	16.6	2.5	293
Q5(richest)	55.0	3.7	9.3	2.1	9.4	2.1	4.0	1.3	5.0	1.4	17.2	2.5	311
M	43.5	9.3	6.6	3.3	10.2	4.4	8.9	4.3	4.6	2.7	26.1	8.4	54
<b>Age</b>													
18-29	64.3	3.4	10.7	1.8	6.4	1.6	1.5	0.8	0.0	0.0	17.1	2.9	322
30-44	58.3	3.4	8.5	1.5	6.7	1.7	6.9	1.7	3.2	1.0	16.5	2.4	360
45-59	32.0	2.9	9.4	1.5	13.0	1.9	9.9	1.7	15.0	2.1	20.7	2.6	379
60-69	20.7	3.0	10.4	2.1	8.3	1.7	12.6	2.9	25.9	3.2	22.2	3.5	177
70-79	9.0	2.1	5.8	1.8	6.0	1.7	10.0	2.1	38.3	4.3	30.9	4.2	132
80+	3.4	2.4	9.6	5.0	1.4	1.0	5.6	3.2	38.6	7.5	41.4	7.4	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	41.5	1.6	9.3	0.8	8.2	0.8	7.4	0.7	12.9	1.0	20.7	1.7	1,419

**6.05 Difficulty with 'self-care' in last 30 days - q2020**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	91.0	1.2	4.9	0.9	1.8	0.4	1.1	0.4	0.7	0.3	0.5	0.4	663
Female	86.0	1.3	6.3	0.8	3.9	0.7	1.8	0.6	0.7	0.2	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	89.4	1.3	4.9	0.7	2.7	0.5	1.7	0.5	0.4	0.2	0.9	0.5	921
Rural	86.2	1.6	7.1	1.1	3.4	0.8	1.1	0.5	1.3	0.4	0.9	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	71.2	3.0	13.5	2.2	8.0	1.6	5.1	2.0	2.2	0.8	0.0	0.0	238
Q2	87.3	2.0	6.3	1.6	3.8	0.9	1.3	0.7	0.7	0.7	0.6	0.4	259
Q3	90.2	1.8	7.2	1.6	1.3	0.7	1.1	0.6	0.2	0.2	0.0	0.0	264
Q4	92.4	1.7	3.3	1.1	0.9	0.5	0.7	0.5	0.4	0.2	2.3	1.1	293
Q5(richest)	96.2	1.1	0.5	0.3	1.8	0.7	0.2	0.2	0.4	0.4	0.9	0.7	311
M	91.8	4.2	2.7	1.9	1.8	1.7	0.0	0.0	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	96.2	1.5	1.3	0.6	0.3	0.3	0.0	0.0	0.0	0.0	2.2	1.4	322
30-44	95.8	1.1	2.0	0.8	1.1	0.5	0.3	0.3	0.3	0.3	0.6	0.6	360
45-59	90.3	1.5	5.7	1.1	2.7	0.8	1.2	0.7	0.1	0.1	0.0	0.0	379
60-69	77.2	3.6	11.4	2.6	5.1	1.5	3.1	1.4	1.6	1.1	1.7	0.9	177
70-79	71.9	3.6	11.1	2.5	9.3	2.2	4.9	1.8	2.8	1.4	0.0	0.0	132
80+	50.8	7.3	25.7	6.5	10.7	4.0	7.6	3.8	3.2	3.2	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	88.3	1.0	5.7	0.6	2.9	0.4	1.5	0.4	0.7	0.2	0.9	0.4	1,419

**6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	93.0	1.1	3.7	0.8	1.3	0.4	1.0	0.4	0.5	0.3	0.5	0.4	663
Female	86.8	1.2	6.1	0.8	3.1	0.6	2.2	0.5	0.8	0.2	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	90.6	1.1	3.9	0.6	2.5	0.6	1.6	0.4	0.4	0.2	0.9	0.5	921
Rural	88.1	1.5	6.9	1.2	1.7	0.6	1.6	0.6	1.0	0.3	0.8	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	72.0	2.8	13.3	2.4	7.2	1.7	5.9	1.6	1.5	0.5	0.0	0.0	238
Q2	89.4	1.9	5.4	1.4	2.3	0.8	0.9	0.5	1.3	0.8	0.6	0.4	259
Q3	92.5	1.6	5.7	1.4	0.7	0.5	0.9	0.5	0.3	0.3	0.0	0.0	264
Q4	93.9	1.6	1.5	0.8	1.6	0.8	0.6	0.4	0.3	0.2	2.1	1.1	293
Q5(richest)	97.2	0.9	0.7	0.4	0.7	0.4	0.5	0.3	0.0	0.0	0.9	0.7	311
M	89.0	4.7	5.6	3.1	0.0	0.0	1.8	1.7	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	96.5	1.5	1.3	0.7	0.0	0.0	0.0	0.0	0.0	0.0	2.2	1.4	322
30-44	96.2	1.2	2.2	1.0	0.8	0.5	0.2	0.2	0.0	0.0	0.6	0.6	360
45-59	92.2	1.1	5.3	1.0	1.5	0.6	1.0	0.4	0.0	0.0	0.0	0.0	379
60-69	80.3	3.5	7.7	2.2	4.8	1.8	3.8	1.5	2.0	1.2	1.4	0.8	177
70-79	73.5	3.6	11.1	2.3	7.6	2.2	5.6	1.9	2.3	1.2	0.0	0.0	132
80+	55.3	8.1	20.2	7.0	8.9	3.7	8.8	3.3	4.8	3.4	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	89.7	0.9	5.0	0.5	2.2	0.4	1.6	0.3	0.6	0.2	0.9	0.4	1,419

**6.07 Amount of 'bodily aches or pains' in last 30 days - q2030**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	60.7	2.3	21.2	1.9	9.7	1.1	6.4	1.0	1.3	0.4	0.7	0.4	663
Female	48.4	1.9	23.7	1.7	16.6	1.2	8.6	1.1	1.5	0.4	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	57.4	2.0	22.3	1.6	11.0	1.0	7.0	1.0	1.2	0.3	1.0	0.5	921
Rural	48.2	2.2	23.0	2.4	17.6	1.5	8.5	1.1	1.8	0.5	0.9	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	38.5	3.3	21.5	2.7	19.2	2.4	17.1	2.2	3.7	1.1	0.0	0.0	238
Q2	48.4	3.2	21.5	2.4	19.2	2.5	7.9	1.6	2.4	1.0	0.6	0.4	259
Q3	54.2	3.4	21.9	2.8	16.2	2.4	6.5	1.5	0.9	0.5	0.3	0.3	264
Q4	58.9	3.6	22.6	3.2	10.3	1.7	5.2	1.4	0.6	0.4	2.3	1.1	293
Q5(richest)	64.9	3.5	25.0	3.2	5.9	1.1	3.4	1.1	0.0	0.0	0.9	0.7	311
M	64.0	6.6	20.2	6.6	5.4	3.0	5.0	3.9	1.8	1.7	3.6	3.6	54
<b>Age</b>													
18-29	71.5	3.1	20.0	2.6	4.3	1.2	2.0	1.0	0.0	0.0	2.2	1.4	322
30-44	68.0	3.2	19.9	2.7	9.3	1.5	1.7	0.7	0.6	0.6	0.6	0.6	360
45-59	44.7	2.9	24.7	2.4	19.2	2.0	10.3	1.7	1.0	0.5	0.0	0.0	379
60-69	37.9	3.6	26.1	3.3	19.5	2.7	11.8	2.3	3.0	1.2	1.7	0.9	177
70-79	35.6	4.2	20.1	3.0	16.3	2.7	20.9	2.9	6.6	1.9	0.6	0.6	132
80+	20.8	6.7	35.5	7.5	26.8	6.4	14.1	4.7	0.7	0.7	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	54.2	1.5	22.5	1.3	13.3	0.8	7.5	0.8	1.4	0.3	1.0	0.4	1,419

**6.08 Amount of 'bodily discomfort' in last 30 days - q2031**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	65.0	2.2	19.6	1.8	10.1	1.2	4.1	0.8	0.4	0.2	0.7	0.4	663
Female	53.9	1.7	22.1	1.7	15.0	1.2	6.6	0.9	1.2	0.4	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	60.7	1.9	21.6	1.5	11.0	1.0	4.7	0.8	0.9	0.3	1.1	0.5	921
Rural	56.2	1.8	19.6	1.6	15.7	1.6	6.8	1.2	0.7	0.2	0.9	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	39.0	3.5	22.5	2.9	19.7	2.4	16.6	2.5	2.2	0.9	0.0	0.0	238
Q2	54.5	2.9	22.7	2.8	16.4	2.2	4.3	1.1	1.5	0.8	0.6	0.4	259
Q3	60.9	3.0	19.9	2.6	14.2	2.2	4.6	1.3	0.3	0.4	0.0	0.0	264
Q4	62.9	3.2	21.1	2.5	9.9	1.8	3.4	1.2	0.5	0.3	2.3	1.1	293
Q5(richest)	73.6	3.3	17.0	2.5	7.1	2.0	1.4	0.7	0.0	0.0	0.9	0.7	311
M	56.7	7.7	31.7	9.1	3.9	2.3	0.0	0.0	1.8	1.7	5.8	4.1	54
<b>Age</b>													
18-29	73.3	2.9	19.0	2.3	5.5	1.5	0.0	0.0	0.0	0.0	2.2	1.4	322
30-44	73.2	2.8	16.9	2.1	7.9	1.7	1.5	0.8	0.0	0.0	0.6	0.6	360
45-59	52.1	2.5	23.1	2.2	17.0	1.8	7.3	1.3	0.2	0.2	0.3	0.3	379
60-69	40.7	3.5	26.3	3.2	20.1	3.1	8.5	2.1	2.7	1.3	1.7	0.9	177
70-79	39.1	3.7	20.4	3.4	19.2	2.9	17.2	3.1	4.1	1.7	0.0	0.0	132
80+	36.2	7.5	27.3	6.7	18.2	6.3	13.6	4.6	2.7	1.9	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	59.1	1.4	20.9	1.1	12.7	0.8	5.5	0.7	0.9	0.2	1.0	0.4	1,419

**6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	74.8	2.1	15.4	1.7	6.4	1.2	2.0	0.6	0.5	0.3	0.9	0.5	663
Female	69.1	1.9	17.4	1.5	8.2	0.9	3.3	0.6	0.8	0.4	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	72.1	1.9	16.7	1.4	7.0	1.0	2.5	0.5	0.5	0.3	1.1	0.5	921
Rural	71.1	2.1	16.0	1.7	7.9	1.3	2.9	0.8	1.0	0.5	1.0	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	53.5	3.5	21.5	2.6	15.0	2.4	8.0	2.0	1.7	0.9	0.3	0.3	238
Q2	73.5	2.9	17.5	2.5	6.0	1.6	1.8	0.7	0.3	0.3	0.8	0.6	259
Q3	75.3	2.6	16.2	2.5	7.1	1.4	1.4	0.7	0.0	0.0	0.0	0.0	264
Q4	74.1	3.4	15.6	2.4	4.2	1.2	1.9	0.8	1.5	1.0	2.7	1.2	293
Q5(richest)	79.5	2.6	13.7	2.2	4.7	1.5	1.2	0.6	0.0	0.0	0.9	0.7	311
M	69.9	6.0	11.3	4.4	12.5	5.3	1.8	1.7	0.9	0.9	3.6	3.6	54
<b>Age</b>													
18-29	78.7	2.9	14.0	2.1	3.0	1.0	0.9	0.5	1.2	0.9	2.2	1.4	322
30-44	80.6	2.7	12.8	2.2	4.5	1.2	0.9	0.5	0.4	0.4	0.9	0.6	360
45-59	72.4	2.3	15.3	1.8	7.9	1.4	4.1	1.1	0.1	0.1	0.2	0.2	379
60-69	58.1	4.0	26.2	3.6	11.0	2.4	2.0	1.1	1.0	0.8	1.7	0.9	177
70-79	56.7	3.5	21.5	2.5	13.1	2.8	7.1	2.2	1.2	0.8	0.3	0.4	132
80+	47.4	7.1	19.9	6.0	22.8	6.9	6.6	2.9	1.4	1.4	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	71.8	1.4	16.5	1.1	7.3	0.8	2.7	0.4	0.7	0.3	1.1	0.4	1,419

## 6.10 Difficulty with 'learning a new task' in last 30 days - q2051

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	81.1	1.7	10.3	1.3	4.0	0.8	1.6	0.5	1.1	0.4	1.8	0.6	663
Female	74.8	1.8	9.6	0.9	6.7	0.8	2.4	0.6	1.7	0.6	4.9	1.0	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	78.4	1.7	10.4	1.1	4.8	0.7	2.0	0.5	1.2	0.4	3.2	0.8	921
Rural	76.4	1.9	9.1	1.2	6.6	1.1	2.1	0.7	1.7	0.6	4.0	1.0	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	53.8	3.3	12.7	1.9	15.2	2.0	7.4	2.0	2.8	1.2	8.1	2.0	238
Q2	76.3	2.9	12.1	2.0	4.4	1.2	1.3	0.6	2.5	1.0	3.4	1.2	259
Q3	82.5	2.1	8.4	1.7	5.0	1.3	2.1	0.8	0.3	0.3	1.7	0.7	264
Q4	83.8	2.4	9.7	1.8	0.8	0.5	0.7	0.5	1.9	1.1	3.0	1.2	293
Q5(richest)	87.9	2.2	7.0	1.7	3.5	1.3	0.0	0.0	0.0	0.0	1.6	0.8	311
M	74.7	5.6	13.3	4.3	6.0	3.7	0.0	0.0	0.9	0.9	5.2	3.8	54
<b>Age</b>													
18-29	87.6	2.7	6.7	1.8	2.3	0.8	0.0	0.0	1.2	0.9	2.2	1.4	322
30-44	90.0	2.0	5.7	1.6	2.5	0.9	1.0	0.5	0.0	0.0	0.8	0.6	360
45-59	80.4	2.2	9.7	1.7	4.2	0.9	2.6	0.9	0.5	0.3	2.6	1.0	379
60-69	62.0	3.6	18.6	2.6	7.5	1.8	1.4	0.8	3.4	1.6	7.1	1.7	177
70-79	53.2	4.1	16.9	2.8	14.2	2.9	5.0	1.7	3.0	1.6	7.7	2.2	132
80+	24.8	6.6	14.0	5.0	27.0	6.4	11.8	5.7	8.8	3.7	13.6	5.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	77.7	1.3	9.9	0.8	5.5	0.6	2.0	0.4	1.4	0.4	3.5	0.6	1,419

**6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	90.0	1.4	6.6	1.3	1.4	0.4	1.0	0.4	0.4	0.2	0.5	0.4	663
Female	84.8	1.6	6.9	1.1	5.0	0.8	1.4	0.5	0.2	0.2	1.6	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	86.7	1.6	7.6	1.3	2.9	0.6	1.5	0.5	0.4	0.1	1.1	0.5	921
Rural	88.2	1.4	5.4	1.1	4.2	0.7	0.8	0.4	0.2	0.2	1.3	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	75.6	3.2	10.0	2.2	10.2	2.1	2.4	1.1	0.7	0.3	1.1	0.6	238
Q2	90.1	1.9	4.3	1.4	2.0	0.8	2.8	1.1	0.2	0.2	0.6	0.4	259
Q3	91.0	1.8	5.8	1.4	2.2	0.9	0.6	0.6	0.4	0.4	0.0	0.0	264
Q4	89.3	2.3	5.7	1.9	2.2	0.8	0.2	0.2	0.3	0.3	2.3	1.1	293
Q5(richest)	88.9	2.3	8.4	1.9	1.0	0.5	0.8	0.5	0.0	0.0	0.9	0.7	311
M	85.4	6.4	5.8	3.2	4.3	2.9	0.0	0.0	0.0	0.0	4.5	3.8	54
<b>Age</b>													
18-29	92.8	1.8	2.7	1.0	1.7	0.7	0.3	0.3	0.3	0.3	2.2	1.4	322
30-44	88.9	2.0	7.7	1.8	1.8	0.7	0.8	0.5	0.3	0.3	0.6	0.6	360
45-59	86.3	2.0	8.9	1.7	3.5	1.0	0.8	0.4	0.0	0.0	0.4	0.3	379
60-69	84.3	3.1	8.3	2.4	2.8	1.4	2.0	1.4	0.3	0.3	2.2	1.0	177
70-79	82.3	3.4	5.4	1.8	7.5	2.4	4.4	2.3	0.0	0.0	0.4	0.4	132
80+	68.9	7.0	9.0	4.1	13.8	6.0	2.9	1.9	3.4	2.0	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	87.2	1.2	6.8	0.9	3.3	0.5	1.2	0.3	0.3	0.1	1.1	0.4	1,419

**6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	83.9	1.7	11.8	1.6	3.0	0.7	0.1	0.1	0.2	0.2	1.0	0.5	663
Female	79.0	1.8	11.7	1.2	5.6	1.0	1.3	0.4	0.1	0.1	2.3	0.7	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	79.7	1.9	12.8	1.4	4.8	0.8	0.9	0.3	0.1	0.1	1.7	0.7	921
Rural	84.4	1.9	9.7	1.5	3.6	0.9	0.3	0.2	0.3	0.2	1.7	0.6	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	73.4	3.5	14.1	2.6	7.9	1.9	1.5	0.9	0.0	0.0	3.0	1.3	238
Q2	82.5	2.9	9.0	2.3	5.7	1.5	1.1	0.8	0.2	0.2	1.5	0.9	259
Q3	86.9	2.1	10.6	2.0	2.5	1.0	0.0	0.0	0.0	0.0	0.0	0.0	264
Q4	79.3	2.8	14.9	2.7	2.4	1.0	0.5	0.3	0.7	0.4	2.3	1.1	293
Q5(richest)	82.8	2.7	11.5	1.8	4.3	1.4	0.0	0.0	0.0	0.0	1.4	0.8	311
M	85.4	5.7	4.4	2.4	2.9	2.7	3.8	2.8	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	85.2	2.4	9.0	1.7	2.6	0.9	0.8	0.6	0.3	0.3	2.2	1.4	322
30-44	80.3	2.7	12.5	2.2	5.5	1.5	0.9	0.2	0.0	0.0	0.8	0.6	360
45-59	80.0	2.2	14.8	2.2	4.6	0.9	0.0	0.0	0.3	0.2	0.2	0.2	379
60-69	82.6	3.4	8.8	2.5	3.6	1.4	2.1	1.2	0.3	0.3	2.7	1.3	177
70-79	81.3	3.6	10.2	2.4	5.2	2.3	0.4	0.4	0.0	0.0	2.9	1.8	132
80+	69.2	7.2	15.2	5.0	6.8	3.4	0.0	0.0	0.0	0.0	8.8	4.4	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	81.3	1.4	11.7	1.1	4.4	0.6	0.7	0.2	0.2	0.1	1.7	0.5	1,419

**6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	90.9	1.1	4.4	0.8	1.4	0.5	2.1	0.5	0.7	0.3	0.5	0.4	663
Female	82.7	1.5	7.8	0.9	3.2	0.6	2.6	0.6	2.2	0.5	1.6	0.8	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	86.7	1.4	7.1	0.8	1.8	0.4	2.0	0.4	1.1	0.4	1.2	0.7	921
Rural	86.2	1.6	4.5	0.9	3.2	0.8	3.0	0.8	2.1	0.6	0.9	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	72.0	2.7	9.3	1.9	5.8	1.4	7.0	1.5	6.0	1.5	0.0	0.0	238
Q2	82.9	2.2	9.2	1.9	3.0	0.9	3.3	1.0	0.7	0.5	0.9	0.5	259
Q3	92.7	1.4	3.8	1.1	2.1	0.8	1.2	0.6	0.2	0.2	0.0	0.0	264
Q4	90.1	2.0	4.7	1.1	0.4	0.3	0.9	0.6	1.1	0.6	2.9	1.3	293
Q5(richest)	92.2	2.0	5.3	1.4	0.8	0.4	0.3	0.3	0.5	0.4	0.9	0.7	311
M	86.3	6.3	2.9	2.7	4.3	2.4	2.9	2.7	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	93.2	1.7	3.1	1.0	0.2	0.2	0.2	0.2	1.1	0.6	2.2	1.4	322
30-44	94.7	1.3	2.8	1.0	0.0	0.0	1.0	0.6	0.5	0.3	1.1	0.7	360
45-59	88.5	1.9	6.1	1.4	3.0	0.8	2.2	0.8	0.3	0.2	0.0	0.0	379
60-69	76.3	3.0	11.5	2.4	4.5	1.6	3.9	1.3	2.1	1.0	1.7	0.9	177
70-79	68.7	3.6	13.6	2.8	8.4	2.0	6.2	1.8	3.0	1.2	0.0	0.0	132
80+	53.5	7.6	13.2	5.4	4.2	2.5	11.6	4.4	14.1	5.6	3.3	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	86.5	1.0	6.2	0.6	2.3	0.4	2.4	0.4	1.5	0.3	1.1	0.5	1,419

**6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	85.0	1.7	9.6	1.3	3.2	0.6	1.3	0.4	0.4	0.2	0.5	0.4	663
Female	80.7	1.7	9.5	1.2	4.9	0.8	2.6	0.7	0.7	0.3	1.7	0.8	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	84.3	1.6	9.7	1.2	3.0	0.5	1.5	0.4	0.4	0.2	1.1	0.7	921
Rural	79.9	2.0	9.3	1.6	6.1	1.0	2.8	0.7	0.9	0.4	1.2	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	59.7	3.2	17.5	2.7	11.7	2.0	7.8	1.8	2.7	1.0	0.6	0.4	238
Q2	81.9	2.6	9.4	2.3	5.1	1.2	2.8	0.9	0.2	0.2	0.6	0.4	259
Q3	88.0	2.0	9.8	2.0	1.9	0.7	0.3	0.3	0.0	0.0	0.0	0.0	264
Q4	87.9	2.0	7.0	1.6	1.7	0.6	0.5	0.3	0.0	0.0	2.9	1.3	293
Q5(richest)	91.3	2.2	5.8	1.8	1.7	0.7	0.0	0.0	0.3	0.3	0.9	0.7	311
M	83.9	5.9	9.1	4.4	3.4	2.2	0.0	0.0	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	95.9	1.5	1.7	0.8	0.2	0.2	0.0	0.0	0.0	0.0	2.2	1.4	322
30-44	91.8	1.8	5.6	1.3	1.5	0.6	0.0	0.0	0.0	0.0	1.1	0.7	360
45-59	76.6	2.2	15.6	2.1	4.9	1.1	2.4	0.9	0.2	0.2	0.2	0.2	379
60-69	72.6	4.0	12.1	2.9	9.6	2.4	2.3	1.0	1.7	1.0	1.7	0.9	177
70-79	67.4	4.1	17.8	3.3	8.3	2.2	5.3	1.7	0.8	0.6	0.3	0.3	132
80+	54.8	7.8	10.6	4.2	11.4	4.4	15.3	6.5	5.9	2.9	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	82.7	1.2	9.5	0.9	4.1	0.5	2.0	0.4	0.6	0.2	1.1	0.5	1,419

**6.15 Problem with 'sleeping' in last 30 days - q2080**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	70.1	2.1	15.3	1.6	8.8	1.2	4.7	0.8	0.6	0.3	0.5	0.4	663
Female	56.0	1.9	16.9	1.3	15.5	1.2	8.8	1.1	1.5	0.5	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	63.3	1.8	17.1	1.4	11.5	0.9	6.3	0.9	0.9	0.4	0.9	0.5	921
Rural	61.3	2.5	14.5	1.6	13.9	1.5	8.1	1.2	1.3	0.5	0.9	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	40.3	3.4	19.1	2.6	23.8	3.0	14.0	2.2	2.8	1.1	0.0	0.0	238
Q2	58.4	3.0	16.5	2.3	14.1	2.2	8.8	1.8	1.5	0.8	0.6	0.4	259
Q3	67.6	3.0	15.3	2.3	10.2	1.9	6.2	1.6	0.6	0.5	0.0	0.0	264
Q4	72.5	3.3	14.4	2.4	6.5	1.2	4.3	1.2	0.0	0.0	2.3	1.1	293
Q5(richest)	69.8	2.9	16.3	2.3	9.5	2.0	2.7	0.8	0.9	0.7	0.9	0.7	311
M	61.5	7.1	14.6	5.1	11.8	5.0	8.5	4.4	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	76.6	3.0	13.6	2.3	6.1	1.6	1.5	0.6	0.0	0.0	2.2	1.4	322
30-44	75.6	2.5	12.0	1.8	8.9	1.8	2.6	0.8	0.2	0.2	0.6	0.6	360
45-59	57.3	2.8	17.9	2.0	14.1	1.9	8.9	1.4	1.8	0.8	0.0	0.0	379
60-69	47.0	4.1	19.9	3.4	18.8	2.8	11.0	2.1	1.6	1.0	1.7	0.9	177
70-79	39.1	4.2	23.3	3.1	17.4	2.9	18.4	3.2	1.8	1.1	0.0	0.0	132
80+	35.4	7.8	17.3	5.4	27.7	7.4	13.0	4.3	4.5	3.3	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	62.6	1.5	16.2	1.1	12.3	0.8	6.9	0.7	1.1	0.3	0.9	0.4	1,419

**6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	52.4	2.4	27.4	2.2	13.8	1.8	5.2	0.9	0.5	0.2	0.8	0.4	663
Female	42.6	2.0	29.0	1.8	17.8	1.6	8.4	1.1	0.9	0.3	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	47.3	2.0	29.4	1.8	14.9	1.8	7.1	1.0	0.5	0.2	0.9	0.5	921
Rural	47.0	2.9	26.3	2.3	17.9	1.9	6.6	1.2	1.2	0.4	1.2	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	38.5	3.4	22.6	2.6	21.7	2.6	14.9	2.4	2.3	0.7	0.0	0.0	238
Q2	48.2	3.4	25.6	2.6	18.2	2.5	6.5	1.7	0.9	0.5	0.6	0.4	259
Q3	50.4	3.2	31.1	2.8	11.3	1.9	6.3	1.3	0.3	0.3	0.5	0.5	264
Q4	47.6	3.4	27.2	3.3	16.7	2.8	5.9	1.5	0.3	0.3	2.3	1.1	293
Q5(richest)	49.7	3.6	33.5	3.0	12.7	2.2	3.0	1.1	0.2	0.2	0.9	0.7	311
M	47.8	7.1	27.7	8.1	16.7	7.0	4.2	2.9	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	51.4	3.0	31.1	2.8	11.7	2.1	3.2	1.0	0.3	0.3	2.2	1.4	322
30-44	47.1	3.3	36.3	3.4	10.8	2.3	4.5	1.4	0.2	0.2	0.9	0.7	360
45-59	48.6	2.5	24.2	2.3	18.7	2.2	7.9	1.3	0.6	0.3	0.0	0.0	379
60-69	41.3	4.1	23.7	3.1	23.1	3.3	8.6	1.9	1.7	0.9	1.7	0.9	177
70-79	46.5	4.4	19.1	2.8	16.4	2.9	16.0	3.2	2.0	1.2	0.0	0.0	132
80+	31.8	6.8	22.8	6.8	32.2	7.2	9.9	4.2	1.3	1.4	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	47.2	1.6	28.3	1.4	15.9	1.3	6.9	0.7	0.7	0.2	1.0	0.4	1,419

**6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	71.0	2.1	16.9	1.6	8.3	1.2	2.6	0.6	0.5	0.3	0.7	0.4	663
Female	56.9	2.1	22.4	1.6	13.1	1.4	5.7	0.8	0.6	0.2	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	66.5	2.0	17.3	1.4	10.3	1.1	4.3	0.6	0.6	0.3	1.1	0.5	921
Rural	57.9	2.3	24.5	1.9	11.9	1.7	4.3	1.1	0.5	0.1	0.9	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	44.7	3.5	24.0	3.1	19.6	2.9	10.5	1.8	1.2	0.4	0.0	0.0	238
Q2	62.0	3.2	20.9	2.6	10.8	2.0	5.5	1.4	0.0	0.0	0.8	0.6	259
Q3	67.6	2.6	23.0	2.6	5.1	1.1	4.0	1.2	0.0	0.0	0.3	0.3	264
Q4	66.6	3.6	17.7	2.7	11.4	2.2	1.2	0.5	0.8	0.4	2.3	1.1	293
Q5(richest)	73.0	3.0	15.9	2.3	8.5	1.9	1.5	0.8	0.2	0.2	0.9	0.7	311
M	61.6	10.9	15.4	5.9	11.6	7.5	4.7	3.0	3.1	3.1	3.6	3.6	54
<b>Age</b>													
18-29	71.0	3.6	16.3	2.5	8.3	1.9	1.6	0.7	0.5	0.5	2.2	1.4	322
30-44	71.5	3.0	18.9	2.6	7.0	1.5	1.9	0.7	0.1	0.1	0.6	0.6	360
45-59	58.8	2.7	23.3	2.3	13.1	1.8	4.2	1.2	0.6	0.3	0.0	0.0	379
60-69	54.6	4.1	17.6	2.5	14.4	2.9	10.8	2.2	1.0	0.8	1.7	0.9	177
70-79	56.4	4.4	22.6	3.3	12.8	3.0	6.6	2.1	0.6	0.6	0.9	0.7	132
80+	42.8	7.4	23.0	6.6	20.2	6.9	10.4	3.6	1.5	1.5	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	63.5	1.5	19.8	1.1	10.9	0.9	4.3	0.6	0.5	0.2	1.0	0.4	1,419

**6.18 Problem with 'worry or anxiety' in last 30 days - q2091**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	63.0	2.2	21.0	1.9	10.2	1.2	4.3	0.8	0.8	0.4	0.7	0.4	663
Female	48.0	2.1	25.9	1.6	15.6	1.4	7.8	1.0	1.4	0.5	1.2	0.6	756
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	55.0	2.3	24.2	1.7	12.8	1.4	5.7	0.8	1.3	0.4	1.0	0.5	921
Rural	55.1	2.6	22.5	1.9	13.7	1.6	6.9	1.2	0.8	0.4	1.0	0.5	498
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	44.0	3.1	23.2	2.6	17.4	2.7	12.4	2.3	2.7	1.2	0.3	0.3	238
Q2	56.6	3.4	21.7	2.8	12.7	2.1	6.4	1.7	1.9	1.0	0.8	0.6	259
Q3	58.5	3.2	24.2	2.9	12.8	1.9	4.3	1.3	0.2	0.2	0.0	0.0	264
Q4	57.7	3.4	22.9	2.6	12.4	2.3	4.1	1.4	0.6	0.4	2.3	1.1	293
Q5(richest)	58.7	3.6	25.4	2.8	10.7	2.2	3.8	1.1	0.5	0.4	0.9	0.7	311
M	44.1	9.6	25.1	7.6	15.7	4.1	11.4	4.6	0.0	0.0	3.6	3.6	54
<b>Age</b>													
18-29	63.3	3.4	19.5	2.5	9.3	2.0	5.2	1.5	0.5	0.4	2.2	1.4	322
30-44	58.4	3.3	24.0	2.6	13.0	2.2	3.5	1.1	0.5	0.4	0.6	0.6	360
45-59	48.3	2.7	28.3	2.2	15.5	1.8	6.7	1.1	1.2	0.6	0.0	0.0	379
60-69	47.6	3.9	21.3	2.5	17.6	3.2	8.9	2.8	2.9	1.6	1.7	0.9	177
70-79	56.3	4.3	20.6	3.3	8.8	2.4	12.1	2.9	1.5	1.5	0.8	0.6	132
80+	52.4	7.2	27.2	7.3	14.9	4.7	2.0	1.5	1.5	1.5	2.0	2.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	55.1	1.7	23.6	1.2	13.1	1.0	6.1	0.7	1.1	0.3	1.0	0.4	1,419

## 7.1 Self-reported utilization of health services and perceived unmet need

	<u>Ambulatory Adult</u>		<u>Ambulatory Child</u>		<u>Inpatient Adult</u>		<u>Inpatient Child</u>		<u>Informal Care</u>		<u>Unmet Care</u>		<u>Non use</u>	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>														
Female	55.6	1.9	10.6	1.1	10.3	1.2	0.6	0.3	4.8	1.0	0.9	0.4	17.2	1.6
Male	48.2	2.3	9.2	1.4	13.3	1.9	1.7	0.6	3.3	0.8	1.7	0.7	22.6	2.2
<b>Area of Residence</b>														
Urban	53.4	2.2	8.5	1.0	12.6	1.3	0.5	0.3	4.5	0.9	1.3	0.4	19.2	1.8
Rural	50.9	2.6	13.0	1.7	9.4	1.6	2.2	0.7	3.6	0.8	1.1	0.6	19.9	2.2
<b>Age Group</b>														
18-29y	57.0	3.6	13.5	2.2	6.0	1.6	2.0	1.0	2.7	1.7	1.6	0.8	17.1	2.6
30-44y	36.8	3.2	31.3	3.0	7.6	1.9	2.9	1.2	1.3	0.7	2.5	1.2	17.5	2.8
45-59y	60.8	3.3	3.6	1.0	11.2	2.0	0.2	0.2	2.4	0.8	0.9	0.5	20.8	2.4
60-69y	59.8	3.5	0.0	0.0	13.9	2.7	0.0	0.0	3.9	1.4	0.4	0.4	22.0	3.2
70+y	47.1	3.6	0.0	0.0	20.5	3.4	0.0	0.0	12.4	2.4	0.4	0.4	19.6	2.7
<b>Education Years</b>														
0														
1-5	50.3	8.0	2.3	2.3	17.1	7.0	0.0	0.0	6.7	3.8	0.0	0.0	23.5	7.2
6-11	50.4	2.3	11.6	1.4	11.8	1.5	1.1	0.4	4.8	1.0	1.1	0.4	19.2	1.9
12+	54.6	2.4	9.0	1.2	11.0	1.5	1.0	0.4	3.6	0.9	1.3	0.4	19.4	2.1
<b>Income Quintiles</b>														
Q1	52.4	3.6	7.5	1.7	11.1	2.1	0.2	0.2	8.1	2.1	0.7	0.5	19.9	2.9
Q2	47.3	3.6	8.6	1.8	14.1	2.4	1.3	0.8	3.8	1.3	2.0	1.0	23.0	3.0
Q3	54.6	3.3	9.9	1.9	12.8	2.1	1.6	1.0	2.7	1.0	0.0	0.0	18.5	2.3
Q4	57.2	3.4	10.6	1.9	8.9	1.8	0.7	0.5	2.8	1.6	2.2	0.9	17.7	2.5
Q5	52.0	3.6	14.6	2.2	9.7	2.6	1.6	0.9	4.5	1.3	0.7	0.7	16.8	3.2
<b>Ownership</b>														
Government	60.5	1.9	10.6	1.1	17.2	1.6	1.6	0.5	4.4	0.8	0.8	0.3	5.0	0.9
Private	78.3	5.4	11.6	4.1	0.0	0.0	0.0	0.0	1.7	1.7	0.0	0.0	8.5	4.3
NGO														
Other														
TOTAL	52.6	1.7	10.0	0.9	11.5	1.0	1.0	0.3	4.2	0.6	1.2	0.3	19.4	1.4

This table presents the percent of people reporting utilization of health services when they last needed care. It combines information from Q7000, Q7001 and Q7016. Utilization was recorded in the WHS in three separate categories: ambulatory care, hospital inpatient care, and informal care. Time cut-offs for utilization recall used here are 1 year for ambulatory care and informal care, and 3 years for hospital inpatient care. Utilization of ambulatory and inpatient care services

have been further sub-divided into "adult" and "child" categories. In addition to utilization, two other related aspects are being reported here - unmet need and non-use (regardless of expressed need) of health services. Unmet need was estimated on the basis of responses to Q7000 and Q7004. Non-use of health services is simply the percent of respondents who reported no need in Q7000 or did not utilize services in the time cut-offs mentioned above.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, and ownership (of health facility). Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership was constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services.

## 7.2 Patient assessed characteristics of ambulatory health services

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate					Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination		
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
<b>Sex</b>																
Female	85.0	2.8							21.5	1.9						
Male	82.3	3.2							21.2	2.6						
<b>Area of Residence</b>																
Urban	83.0	3.0							20.8	2.0						
Rural	86.3	3.6							23.0	2.7						
<b>Age Group</b>																
18-29y	78.3	4.9							21.8	3.5						
30-44y	80.5	4.2							25.3	3.4						
45-59y	85.4	3.7							19.9	2.5						
60-69y	92.0	3.3							23.0	4.1						
70+y	86.4	5.1							15.7	2.2						
<b>Education Years</b>																
0																
1-5	100.0	0.0							19.9	3.6						
6-11	85.2	4.0							23.5	2.7						
12+	82.5	2.8							20.0	1.9						
<b>Income Quintiles</b>																
Q1	94.1	3.2							18.7	2.9						
Q2	88.1	4.1							22.5	3.6						
Q3	86.3	4.9							22.3	4.5						
Q4	76.8	4.6							24.0	2.7						
Q5	79.1	4.7							19.9	3.0						
<b>Condition</b>																
Child-Visit	97.9	2.2							20.9	6.1						
Dental	38.0	7.2							20.6	3.6						
Arthritis	87.4	8.5							25.2	8.4						
Asthma	59.6	29.7							18.8	4.8						
Heart-Disease	100.0	0.0							23.2	6.2						
Minor-Surgery	100.0	0.0							26.4	7.7						

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
Other	90.9	2.8							22.3	2.7						
<b>Ownership</b>																
Government	87.7	2.3							21.7	1.9						
Private	46.6	8.4							18.5	3.1						
NGO																
Other																
<b>Self-Assessed Health</b>																
Very-Good	73.5	6.9							14.7	2.1						
Good	81.5	3.3							21.2	2.3						
Moderate	88.7	3.2							23.6	3.1						
Bad	89.2	5.4							22.4	4.4						
Very-Bad	100.0	0.0							35.0	14.2						
TOTAL	83.9	2.4							21.4	1.7						

This table looks at various characteristics of a visit to an ambulatory care facility, for those who utilized such services in the previous 12 months (Q7204 and Q7205). Characteristics of a visit reported here include:

- (1) The percent of patients visiting a medical doctor (of those visiting any ambulatory provider) (Q7302).
- (2) Patients' rating of *skills*, *equipment*, and *drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7304, Q7305 and Q7306, respectively.
- (3) Barriers or facilitators to care access have been reported here for those who utilized ambulatory care services in the last 12 months. They include:
  - (a) *Mean time taken to reach facility*: patient reported times from Q7307 were used to calculate averages across sample sub-groups.
  - (b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7308.
  - (c) *Mean cost of visit*: patient reported costs in Q7309-Q7313 were summed up for an individual and then averaged across sample sub-groups.
  - (d) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination listed in Q7328-Q7334.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include: sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents could select one of the

following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

### 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>												
Female			56.4	3.7	10.1	2.1			12.3	2.1		
Male			56.9	3.6	5.3	1.6			8.1	2.0		
<b>Area of Residence</b>												
Urban			55.1	3.5	8.1	2.0			9.8	1.8		
Rural			60.3	4.0	8.2	2.2			12.5	2.4		
<b>Age Group</b>												
18-29y			67.4	5.0	10.9	3.4			9.8	3.0		
30-44y			51.6	5.1	9.2	3.1			9.5	3.0		
45-59y			53.7	4.8	7.1	2.3			12.0	3.0		
60-69y			60.3	7.3	5.1	2.6			14.4	3.7		
70+y			46.9	6.6	7.5	4.7			6.4	3.1		
<b>Education Years</b>												
0												
1-5			55.2	18.4	9.5	9.2			0.0	0.0		
6-11			55.7	3.8	7.4	2.4			12.2	2.6		
12+			57.2	3.9	8.6	2.0			9.8	2.0		
<b>Income Quintiles</b>												
Q1			49.4	7.1	10.3	4.8			10.1	3.7		
Q2			63.2	5.2	5.7	2.6			12.8	4.1		
Q3			53.9	6.3	7.0	2.8			13.5	3.5		
Q4			63.0	5.0	8.5	3.3			7.5	2.4		
Q5			54.6	5.0	9.6	2.9			10.4	3.2		
<b>Condition</b>												
Child-Visit			62.6	9.9	5.6	3.9			10.9	5.1		
Dental			35.2	6.5	8.4	4.4			4.6	2.6		
Arthritis			43.2	13.1	0.0	0.0			20.8	11.0		
Asthma			40.4	29.7	0.0	0.0			0.0	0.0		
Heart-Disease			59.7	13.0	6.7	4.6			7.6	5.1		
Minor-Surgery			87.5	11.9	28.2	17.0			13.4	12.6		
Other			54.3	4.2	6.4	2.2			8.3	2.4		
<b>Ownership</b>												
Government			59.6	2.8	9.0	1.7			11.6	1.6		
Private			28.9	7.8	1.3	1.3			0.0	0.0		
NGO												
Other												
<b>Self-Assessed Health</b>												
Very-Good			50.7	8.0	3.1	2.2			1.4	1.4		
Good			55.5	3.8	6.4	1.6			10.0	2.1		
Moderate			58.5	4.3	10.9	3.2			13.7	3.2		
Bad			56.1	10.1	2.6	2.6			15.7	6.2		
Very-Bad			100.0	0.0	49.1	25.5			0.0	0.0		
TOTAL			56.6	2.7	8.1	1.6			10.6	1.4		

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

### 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Autonomy		Confidentiality				Choice of Healthcare Provider		Basic Amenities			
	Treatment Information %	Involvement SE	Talk Privately %	Confidentiality of Records SE	Confidentiality of Records %	Confidentiality of Records SE	Healthcare Provider %	Healthcare Provider SE	Cleanliness %	Space SE	Cleanliness %	Space SE
<b>Sex</b>												
Female		18.1	2.6	17.4	2.2		13.9	2.3	15.6	2.7		
Male		14.3	2.8	15.4	2.8		17.2	3.2	18.6	2.9		
<b>Area of Residence</b>												
Urban		14.0	2.2	16.3	2.1		12.4	2.1	19.2	2.6		
Rural		23.1	3.7	17.3	2.9		22.2	4.0	10.9	2.7		
<b>Age Group</b>												
18-29y		18.2	4.6	23.6	3.8		19.0	4.2	28.2	4.7		
30-44y		17.5	4.1	16.3	3.9		15.5	4.1	17.5	4.0		
45-59y		18.3	4.0	16.5	3.4		18.1	3.6	14.7	3.9		
60-69y		17.0	4.1	15.0	4.4		11.0	4.1	14.5	5.0		
70+y		8.3	3.3	7.2	3.4		7.2	3.0	2.9	2.0		
<b>Education Years</b>												
0												
1-5		9.5	9.2	24.6	15.5		18.3	12.4	0.0	0.0		
6-11		19.7	2.9	14.7	2.8		14.3	2.9	10.8	2.3		
12+		14.7	2.7	17.7	2.4		15.8	2.6	21.5	3.1		
<b>Income Quintiles</b>												
Q1		10.5	3.8	6.7	3.3		13.5	4.3	8.1	3.5		
Q2		19.6	4.5	21.5	5.0		15.8	4.3	22.2	5.0		
Q3		16.0	3.7	13.6	3.3		10.9	3.6	12.3	4.0		
Q4		13.5	3.6	22.8	4.2		14.5	3.6	18.9	4.2		
Q5		20.9	4.5	16.5	3.5		20.9	4.2	18.8	3.9		
<b>Condition</b>												
Child-Visit		5.5	3.6	24.3	8.8		15.3	6.5	19.2	7.2		
Dental		19.1	6.2	15.9	5.6		6.4	3.0	23.3	6.7		
Arthritis		26.3	11.6	9.5	6.7		28.6	12.2	17.5	8.8		
Asthma		0.0	0.0	36.6	28.9		0.0	0.0	0.0	0.0		
Heart-Disease		3.7	3.6	16.5	9.8		17.9	8.9	11.6	5.6		
Minor-Surgery		21.8	14.6	47.9	19.0		28.1	17.9	46.1	19.2		
Other		12.2	2.8	14.4	2.8		11.5	2.6	13.1	3.0		
<b>Ownership</b>												
Government		16.4	2.0	18.1	1.9		16.8	2.1	17.1	2.3		
Private		14.6	6.4	5.8	3.3		3.2	2.3	8.1	3.7		
NGO												
Other												
<b>Self-Assessed Health</b>												
Very-Good		7.1	3.6	13.6	5.2		8.4	3.8	16.1	4.9		
Good		16.8	3.2	12.5	2.4		15.5	2.9	17.8	3.0		
Moderate		18.7	3.4	22.0	3.5		16.7	3.5	16.1	3.4		
Bad		17.2	6.5	11.0	5.3		14.2	5.7	11.8	5.3		
Very-Bad		49.1	25.5	49.1	25.5		49.1	25.5	0.0	0.0		
TOTAL		16.6	2.0	16.6	1.7		15.2	1.9	16.8	2.0		

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.4 Patient assessed characteristics of hospital inpatient health services

	Patients staying for more than 6 days on last hospital visit		Patients rating characteristics of health care provider as adequate / more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Patients waiting more than a month to be admitted		Mean number of patients sleeping in the same room		Mean cost for a hospital visit		Percent of population perceiving discrimination	
	%	SE	Skills	Equipment	Drug supplies				Minutes	SE	%	SE	%	SE	No	SE	mean cost	SE	%	SE
<b>Sex</b>																				
Female	59.4	4.1						40.6	3.2			0.7	0.4	4.9	0.6					
Male	61.8	4.4						38.1	3.1			3.7	1.6	5.1	0.7					
<b>Area of Residence</b>																				
Urban	56.9	4.6						36.2	3.0			1.5	0.9	5.0	0.7					
Rural	66.5	3.9						46.0	3.4			2.7	1.1	5.0	0.6					
<b>Age Group</b>																				
18-29y	34.3	6.3						34.9	3.4			3.2	3.2	4.7	0.8					
30-44y	45.0	6.0						37.1	4.6			1.2	0.9	4.4	0.7					
45-59y	65.8	4.8						44.8	4.7			2.1	1.2	5.0	0.7					
60-69y	80.7	4.7						44.4	5.8			1.3	1.3	5.3	0.7					
70+y	72.8	5.9						36.3	3.6			1.7	1.2	5.5	0.8					
<b>Education Years</b>																				
0																				
1-5	76.7	10.9						39.1	10.1			0.0	0.0	6.0	1.0					
6-11	70.1	3.6						40.1	2.7			1.4	0.7	5.2	0.6					
12+	49.9	4.6						39.2	3.6			2.4	1.2	4.8	0.7					
<b>Income Quintiles</b>																				
Q1	75.8	4.0						44.2	3.8			0.0	0.0	5.6	0.7					
Q2	73.9	5.0						45.0	4.6			2.9	1.7	5.1	0.7					
Q3	63.4	5.0						37.1	4.1			0.8	0.8	5.2	0.8					
Q4	47.2	7.7						31.2	3.8			1.4	1.0	4.4	0.8					
Q5	41.5	5.6						41.4	6.2			4.9	3.2	4.6	0.8					
<b>Condition</b>																				
Child-Visit	33.6	8.8						42.4	6.6			0.0	0.0	4.1	0.9					
Fever	61.7	14.0						34.5	7.8			0.0	0.0	4.9	1.0					
Childbirth	60.6	7.2						27.1	3.4			0.0	0.0	4.6	0.8					
Arthritis	83.7	8.0						47.6	10.9			9.3	6.1	4.1	0.8					
Asthma	77.3	13.1						60.0	16.8			0.0	0.0	4.7	1.1					
Heart-Disease	83.9	7.4						38.9	6.0			0.0	0.0	5.4	0.8					
Injury	40.4	8.7						34.3	5.6			0.0	0.0	4.7	0.9					
Minor-Surgery	42.2	6.4						44.0	5.3			4.7	2.4	4.6	0.7					
Other	69.1	4.2						40.0	3.5			0.9	0.6	5.5	0.7					
<b>Ownership</b>																				
Government	60.3	3.4						39.9	2.5			1.9	0.7	5.0	0.6					
Private	0.0	0.0						15.5	0.5			0.0	0.0	1.5	0.5					
NGO																				
Other																				
<b>Self-Assessed Health</b>																				
Very-Good	29.6	7.7						33.4	5.8			0.0	0.0	5.0	0.8					
Good	50.9	5.0						35.5	2.6			1.2	0.7	4.7	0.7					
Moderate	63.3	5.0						41.5	3.9			4.1	1.8	5.1	0.7					
Bad	80.0	4.5						46.4	6.2			0.0	0.0	5.3	0.7					
Very-Bad	85.7	7.1						43.5	6.4			0.0	0.0	5.7	0.9					
TOTAL	60.3	3.4						39.6	2.4			1.9	0.7	5.0	0.6					

This table looks at various characteristics of a visit to a hospital inpatient care facility, for those who utilized such services in the past 3 years (Q7200, Q7201 and Q7402). Characteristics of a visit reported here include:

(1) The percent of patients who stayed 6 days or more at an inpatient facility during their last visit (Q7404).

(2) Patients' rating of *skills, equipment, and drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7405, Q7406 and Q7407, respectively.

(3) Barriers or facilitators to care access have been reported here for those who utilized inpatient care services in the last 3 years. They include:

(a) *Mean time taken to reach facility*: patient reported times from Q7408 were used to calculate averages across sample sub-groups.

(b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7409.

(c) *Patients waiting more than a month to be admitted*: percent of patients who reported waiting more than a month to be admitted in Q7410.

(d) *Mean number of patients sleeping in same room*: averages across sample sub-groups have been constructed for numbers reported in Q7417.

(e) *Mean cost of visit*: patient reported costs in Q7411-Q7415 were summed up for an individual and then averaged across sample sub-groups.

(f) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination asked in Q7433-Q7439.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

**7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1**

	Prompt Attention				Dignity				Communication				Autonomy			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions		Treatment Information		Involvement	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>																
Female			30.4	3.4	16.0	2.5			19.8	3.1					33.7	3.4
Male			30.3	3.5	9.8	2.4			14.5	3.1					24.0	3.8
<b>Area of Residence</b>																
Urban			28.7	3.1	12.6	2.1			19.0	2.9					29.1	2.9
Rural			33.3	3.8	15.5	3.1			15.4	2.8					31.5	3.9
<b>Age Group</b>																
18-29y			40.8	5.8	25.2	4.4			29.6	5.3					48.8	6.3
30-44y			25.7	4.9	10.7	3.3			11.1	3.3					21.3	4.2
45-59y			36.3	4.6	13.2	3.4			16.8	4.1					32.3	4.3
60-69y			31.3	6.6	15.3	5.1			13.9	5.0					25.0	5.4
70+y			20.1	4.6	6.9	2.5			18.3	5.8					24.5	5.6
<b>Education Years</b>																
0																
1-5			19.3	13.8	4.6	4.6			11.8	7.3					16.2	8.9
6-11			33.3	3.7	16.6	2.7			16.2	2.9					30.0	3.3
12+			28.2	3.3	11.2	2.2			19.5	3.0					30.6	3.4
<b>Income Quintiles</b>																
Q1			28.3	5.4	18.3	3.9			19.2	3.7					25.9	4.9
Q2			23.8	4.3	11.7	3.6			14.9	3.9					34.4	4.5
Q3			30.9	4.6	14.6	3.8			24.9	4.3					32.4	5.0
Q4			33.1	6.6	13.6	4.4			9.8	4.1					30.5	7.0
Q5			36.6	6.0	8.2	3.6			13.2	4.1					22.3	4.5
<b>Condition</b>																
Child-Visit			36.3	9.4	12.9	5.2			6.5	4.1					36.8	10.2
Fever			19.7	9.7	8.4	6.0			0.0	0.0					22.6	10.0
Childbirth			14.1	5.2	27.7	6.7			19.7	5.6					37.3	7.3
Arthritis			31.7	9.5	23.6	9.9			26.8	8.9					31.0	10.4
Asthma			25.6	13.2	0.0	0.0			6.7	6.7					22.2	15.0
Heart-Disease			21.3	6.4	6.3	4.0			10.8	4.9					21.6	6.0
Injury			41.8	9.5	12.9	6.0			25.0	7.3					40.4	9.6
Minor-Surgery			34.5	5.3	12.9	3.2			19.4	4.4					25.4	4.5
Other			32.2	3.8	12.3	2.7			17.1	3.1					31.5	4.1
<b>Ownership</b>																
Government			30.2	2.5	13.7	1.8			18.0	2.2					30.3	2.4
Private			0.0	0.0	0.0	0.0			0.0	0.0					0.0	0.0
NGO																
Other																
<b>Self-Assessed Health</b>																
Very-Good			29.6	7.9	6.8	4.0			24.1	7.9					35.8	9.7
Good			26.4	4.0	11.2	2.8			9.6	2.3					24.5	4.0
Moderate			34.4	4.1	16.0	2.9			20.9	4.1					34.4	4.4
Bad			31.2	5.7	11.3	3.6			17.2	4.7					25.7	4.6
Very-Bad			29.4	9.7	25.9	10.6			33.8	10.8					34.3	9.7
TOTAL			30.4	2.5	13.6	1.8			17.7	2.2					29.9	2.3

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Confidentiality		Choice of health care provider		Basic Amenities				Social Support	
	Talk Privately	Confidentiality of records			Cleanliness	Space			Family Visit	External Contact
	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>										
Female	31.1	3.2			30.1	3.0	34.0	3.5	3.8	1.2
Male	25.1	3.0			36.9	4.6	30.5	3.7	2.2	1.2
<b>Area of Residence</b>										
Urban	27.7	3.1			31.8	3.7	37.5	3.6	4.4	1.3
Rural	30.6	4.0			34.6	3.9	23.8	3.2	1.0	0.7
<b>Age Group</b>										
18-29y	42.0	6.0			38.8	5.7	44.5	6.7	9.2	3.4
30-44y	22.9	4.8			27.4	4.8	31.2	5.8	2.8	2.0
45-59y	27.2	3.8			37.2	4.6	34.1	4.5	2.1	1.5
60-69y	23.1	7.0			34.2	6.2	33.0	6.5	0.0	0.0
70+y	29.4	5.8			27.6	5.4	23.6	4.6	2.3	1.6
<b>Education Years</b>										
0										
1-5	13.6	7.9			37.6	14.1	11.8	7.3	0.0	0.0
6-11	26.4	3.5			32.3	3.5	29.4	3.1	3.1	1.3
12+	31.8	3.4			33.0	4.2	37.0	4.0	3.4	1.3
<b>Income Quintiles</b>										
Q1	32.8	5.3			32.3	4.6	25.8	5.0	3.0	1.8
Q2	29.3	5.4			33.3	5.6	31.9	5.7	6.9	3.0
Q3	23.7	4.9			36.1	5.6	34.8	4.7	3.9	2.2
Q4	24.2	6.2			34.2	6.5	36.2	6.6	0.0	0.0
Q5	28.9	5.4			30.4	5.9	38.0	6.6	2.1	1.6
<b>Condition</b>										
Child-Visit	26.8	7.5			30.8	8.6	17.8	5.9	10.1	5.3
Fever	32.9	13.2			33.7	13.3	15.4	10.4	12.7	9.6
Childbirth	26.9	6.3			18.6	5.2	40.8	7.1	4.9	3.4
Arthritis	46.7	11.6			22.5	9.6	22.7	9.8	11.7	6.8
Asthma	28.9	14.3			24.9	13.8	34.4	19.1	0.0	0.0
Heart-Disease	24.7	6.9			32.3	8.3	28.0	6.8	0.0	0.0
Injury	21.6	7.9			58.9	8.5	26.8	8.3	4.0	4.0
Minor-Surgery	32.2	4.5			29.7	5.2	32.5	5.6	2.7	1.6
Other	27.4	3.6			33.7	4.1	35.5	4.1	2.0	1.2
<b>Ownership</b>										
Government	28.9	2.5			32.3	2.8	32.8	2.6	3.2	0.9
Private	0.0	0.0			0.0	0.0	0.0	0.0	0.0	0.0
NGO										
Other										
<b>Self-Assessed Health</b>										
Very-Good	27.3	8.4			29.4	8.0	31.8	8.7	4.3	4.2
Good	25.8	4.1			30.2	3.8	32.9	4.1	4.7	1.9
Moderate	28.3	4.1			36.7	4.7	30.5	4.2	2.6	1.3
Bad	34.4	6.8			30.4	6.3	34.5	6.3	1.9	1.8
Very-Bad	30.6	10.3			39.5	10.3	39.1	10.7	0.0	0.0
TOTAL	28.7	2.5			32.8	2.8	32.6	2.6	3.2	0.9

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"

	Prompt Attention		Dignity		Communication		Autonomy		Confidentiality		Choice		Basic Amenities		Social Support	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>																
Female																
Male																
<b>Area of Residence</b>																
Urban																
Rural																
<b>Age Group</b>																
18-29y																
30-44y																
45-59y																
60-69y																
70+y																
<b>Education Years</b>																
0																
1-5																
6-11																
12+																
<b>Income Quintiles</b>																
Q1																
Q2																
Q3																
Q4																
Q5																
<b>Ownership</b>																
Government																
Private																
NGO																
Other																
<b>Self-Assessed Health</b>																
Very-Good																
Good																
Moderate																
Bad																
Very-Bad																
<b>Type of Care</b>																
Ambulatory-Adult																
Ambulatory-Child																
Inpatient-Adult																
Inpatient-Child																
Non-Users																
TOTAL																

This table shows the percent of people who rated a responsiveness domain as "Extremely Important". All survey respondents were queried on their perception of the importance of each of the eight responsiveness domains in section 7100 of the WHS where each domain was described in detail (Q7100-Q7107). Respondents were asked to say whether the domain was "Extremely Important", "Very Important", "Moderately Important", "Slightly Important" or "Not

Important".

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, ownership (of health facility), self-assessed health, and type of care. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, self-assessed health and type of care sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad". Type of care is a constructed sample sub-group combining information from Q7000, Q7001 and Q7016. There are five categories of type of care: "Ambulatory adult", "Ambulatory child", "Inpatient adult", "Inpatient child" and "Non users".

---

## 8.0 Expenditure Background Indicators (2002)

---

GDP per capita (\$)	6,390
Total health expenditure per capita (\$)	496
Total population (000)	9,923
Life expectancy at birth (Both sexes)	72.6
Life expectancy at birth (male)	68.4
Life expectancy at birth (female)	76.8
Healthy life expectancy at birth (Both sexes)	64.9
Healthy life expectancy at birth (Male)	61.5
Healthy life expectancy at birth (Female)	68.2
Government funded health expenditure as a percentage of total health expenditure	70.2
Social security funded health expenditure as a percentage of total health expenditure	57.1
Private health insurance premiums as a percentage of total health expenditure	0.4
Out-of-pocket payment as a percentage of total health expenditure	26.3

## 8.1 Distribution of health payments across different population groups

	Mean Household expenditure (EXP) in local currency	% Poor	% Impoverished	% Catastrophic	OOP % EXP	OOP % non-subsistence spending	Mean OOP in local currency
<b>Catastrophic</b>							
No	76,350	9.2	1.7		6.9	11.2	5183
Yes	86,877	13.8	17.3		39.5	58.6	42073
<b>Poor</b>							
No	83,243		3.7	10.1	10.0	15.7	9805
Yes	23,250		0.0	15.0	13.2	20.1	2162
<b>Insurance</b>							
No	70,863	0.0	0.0	0.0	4.0	5.4	2854
Yes	77,470	9.7	3.4	10.6	10.3	16.2	9079
<b>Residence</b>							
rural	66,949	13.8	4.8	12.8	10.7	18.0	6455
urban	82,536	7.7	2.7	9.5	10.2	15.2	10331
<b>Income_Quintile</b>							
1st(poorest)	32,621	48.0	12.6	12.8	10.3	17.5	2526
2nd	54,114	0.0	2.6	9.0	9.4	16.7	4763
3rd	67,377	0.0	0.9	8.7	8.4	13.6	5251
4th	84,445	0.0	0.5	8.8	9.8	15.1	7994
5th(richest)	149,237	0.0	0.3	13.3	13.8	17.8	24892
total	77,459	9.6	3.4	10.5	10.3	16.2	9068

OOP: Out of pocket

*Household consumption expenditure (EXP)* comprises both monetary and in-kind payment on all goods and services and the money value of the consumption of home-made products.

*Poor* - a household is considered poor when its total household expenditure is smaller than its subsistence spending  
*Impoverishment* - a non-poor household is impoverished by health payment when it becomes poor after paying for health services.

*Catastrophic health expenditure* occurs when a household's total out-of-pocket health payments equals or exceeds 40% of the households capacity to pay or non-subsistence spending.

*Out of pocket health payments (OOP)* refer to payments made by households at the point they receive health services. It includes doctor's consultation fees, purchases of medication and hospital bills. Spending on alternative and/or traditional medicine is included. Expenditure on health related transportation and special nutrition is excluded. Insurance reimbursements are also excluded.

*Subsistence spending* - household subsistence spending is the minimum requirement to maintain a basic life in a society. A poverty line is used in the analysis as subsistence spending.

*Food expenditure* is the amount spent on all foodstuffs by the household plus the value of the family's own food production consumed within the household. It excludes expenditure on alcoholic beverages, tobacco and food consumption outside the home (e.g hotel and restaurants).

*Household capacity to pay* - is defined as a household's non-subsistence spending.

---

## 8.2 Percent distribution of structure of out-of-pocket health payments

---

	Inpatient	Outpatient	Traditional	Drugs	Other	Total
<b>Catastrophic</b>						
No	1.3	2.4	1.2	74.2	21.0	100
Yes	9.9	5.5	0.2	21.0	63.4	100
<b>Poor</b>						
No	5.6	4.0	0.7	47.2	42.5	100
Yes	1.8	0.4	0.7	88.6	8.5	100
<b>Insurance</b>						
No	0.3	0.3	0.3	98.1	0.9	100
Yes	5.5	3.9	0.7	48.2	41.7	100
<b>Residence</b>						
rural	3.6	2.7	0.6	65.6	27.5	100
urban	6.1	4.3	0.8	42.9	46.0	100
<b>Income_Quintile</b>						
1st(poorest)	2.3	0.7	0.4	88.4	8.2	100
2nd	3.1	2.9	0.0	86.2	7.9	100
3rd	1.8	1.1	1.0	70.5	25.6	100
4th	7.8	2.8	1.1	66.3	22.0	100
5th(richest)	6.3	5.4	0.7	26.2	61.4	100
total	5.5	3.9	0.7	48.2	41.7	100

---

---

### 8.3 Financial sources used by households for paying for health services

---

	Savings	Sold Items	Borrow from relatives &	Borrow from others	Health insurance	Current Income	Other
<b>OOP%non_subistence_spending</b>							
less than 10%	7.2	0.5	1.0	0.1	2.8	85.2	2.1
10-20%	8.2	0.5	1.8	0.5	1.9	89.7	1.9
20-40%	13.4	0.7	3.1	2.5	4.0	96.1	4.3
above 40%	19.0	4.4	2.1	4.6	7.4	88.3	4.3
<b>Hospitalization</b>							
No	8.2	0.9	1.3	0.5	3.3	89.8	2.3
Yes	15.6	1.2	3.1	3.4	3.6	84.8	4.2
<b>Insurance</b>							
No	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Yes	9.9	0.9	1.7	1.2	3.4	88.8	2.7
<b>Residence</b>							
rural	10.0	0.7	2.1	1.2	4.2	94.5	3.2
urban	9.8	1.1	1.6	1.2	3.0	85.8	2.5
<b>Income_Quintile</b>							
1st(poorest)	6.4	0.8	1.8	0.9	2.9	88.5	2.6
2nd	12.5	0.5	1.3	0.8	5.5	93.6	2.6
3rd	10.5	1.0	1.2	1.9	3.1	89.9	2.2
4th	6.7	0.5	1.0	2.0	2.7	85.0	3.7
5th(richest)	13.4	1.9	3.3	0.2	2.7	86.3	2.5
total	9.9	0.9	1.7	1.2	3.4	88.6	2.7

## 8.4 Insurance Coverage

	Insured	SE	Mandatory Insurance	SE	Voluntary Insurance	SE	Insured as dependent	SE	Mandatory and Voluntary	SE
Q1(poorest)	16.6	1.3	15.2	6.4	16.6	1.3	0.0	0.0	15.2	6.4
Q2	19.4	1.4	9.7	4.8	19.4	1.4	0.0	0.0	9.7	4.8
Q3	20.6	1.4	30.2	6.2	20.6	1.4	0.0	0.0	30.2	6.2
Q4	21.2	1.4	18.0	5.2	21.2	1.4	100.0	0.0	18.0	5.2
Q5(richest)	22.2	1.5	26.9	7.0	22.2	1.5	0.0	0.0	26.9	7.0
Total	100		100		100		100		100	
Number	1,988.0		92.0		1,988.0		2.0		92.0	









