

REPORT OF LATVIA

TABLE OF CONTENTS

1	Introduction
2	Methodology
2.1	Sampling
2.2.1	Geographic Component of the Survey
2.2.2	The map
2.3	Survey Metrics
2.3.1	Sample deviation index
2.3.2	HH Sample deviation index
2.3.3	Response rate
2.3.4	Missing data
2.3.5	Reliability
3	Socio-Demographic Profile
3.1	Household composition by residence and age groups according to income quintile
3.2	Selected age distributions by income quintile
3.3	Characteristics of head of household by income quintile
3.4	Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex
3.5	Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education
4	Risk Factors
4.1	Prevalence of smoking and average daily tobacco consumption
4.2	Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers
4.3	Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity
4.4	Access to improved drinking-water: percent distribution of households by

- type of access to drinking water
- 4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility
- 4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used
- 5 Coverage
 - 5.01 Tuberculosis
 - 5.02 HIV Prevention and Screening
 - 5.03 Antenatal Care and Mother to Child Transmission of HIV
 - 5.04 Birth Delivery Care
 - 5.05 Child's Health - Common Illness
 - 5.06 Child's Health - Danger Signs
 - 5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card
 - 5.08 Vaccination Status of Children 12-23 Months
 - 5.09 Women's Cancer Screening
 - 5.10 Oral Health and Injuries
 - 5.11 Self Reported Non-Communicable Conditions
 - 5.12 Mental Health Conditions
 - 5.13 Cataracts
- 6 Health State Descriptions
 - 6.01 In general, rate your health today - q2000
 - 6.02 Difficulty with 'work and household activities' in last 30 days - q2001
 - 6.03 Difficulty with 'moving around' in last 30 days - q2010
 - 6.04 Difficulty with 'vigorous activities' in last 30 days - q2011
 - 6.05 Difficulty with 'self-care' in last 30 days - q2020
 - 6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021
 - 6.07 Amount of 'bodily aches or pains' in last 30 days - q2030

- 6.08 Amount of 'bodily discomfort' in last 30 days - q2031
- 6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050
- 6.10 Difficulty with 'learning a new task' in last 30 days - q2051
- 6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060
- 6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061
- 6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071
- 6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072
- 6.15 Problem with 'sleeping' in last 30 days - q2080
- 6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081
- 6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090
- 6.18 Problem with 'worry or anxiety' in last 30 days - q2091
- 6.19 Adjusted for vignette responses
- 7 Health System Responsiveness
 - 7.1 Self-reported utilization of health services and perceived unmet need
 - 7.2 Patient assessed characteristics of ambulatory health services
 - 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1
 - 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2
 - 7.4 Patient assessed characteristics of hospital inpatient health services
 - 7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1
 - 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2
 - 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"
- 8 Health Expenditure, Insurance and Human Resources for Health

- 8.0 Expenditure Background Indicators (2002)
- 8.1 Distribution of health payments across different population groups
- 8.2 Percent distribution of structure of out-of-pocket health payments
- 8.3 Financial sources used by households for paying for health services
- 8.4 Insurance Coverage
- 8.5 Human resources for health
- 9.1 Adult Mortality Chart 1
- 9.2 Adult Mortality Chart 2
- 9.3 Adult Mortality Chart 3
- 9.4 Adult Mortality Chart 4
- 10 Stress and Social Capital
- 10.1 Inability to control the important things in life - q8000
- 10.2 Inability to cope - q8001
- 10.3 Satisfaction with your health - q8002
- 10.4 Feeling safe when alone at home - q8011
- 10.5 Feeling safe when walking down the street alone after dark - q8012

1. Introduction

The World Health Organization initiated the World Health Survey (WHS) as a *data collection platform* to obtain information on the health of populations and on health systems. The WHS responds to the need of countries for a detailed and sustainable health information system and gathers data through surveys to measure essential population health parameters. The WHS uses standard survey procedures and instruments for general population surveys to gather comparable data across WHO Member States. These methods and instruments have been refined through scientific review of literature, extensive consultations with international experts and large-scale pilot tests. WHS is designed to evolve through its implementation by continuous input from collaborators, including policy makers, survey institutions, scientists and other interested parties. The data is jointly owned by the countries and the WHO, and there is a commitment for a long-term data collection, building local capacity and using this information for health policy.

The objectives of the World Health Survey were to:

- develop a **means** of providing:
 - **valid, relevant, reliable** and **comparable** information on **outcomes, functions** and **inputs** to health systems
 - to **supplement** the information provided by routine **HIS**
- build the **evidence base** necessary for policy-makers:
 - to **routinely monitor** if health systems are achieving the **desired goals**
 - and in some countries, to assess if the **increase in health resources** is having the desired effect

The introductory chapter of each country report will provide details on the current sources of data from routine health information systems and ad hoc surveys. It will describe the health situation in the country and identify gaps. It will also present a brief overview on the health system organization within the country.

2. Sampling

The WHO sampling guidelines emphasized scientific principles of survey sampling as explicit standards for quality, included examples of good sampling plans, and identified quality assurance standards for countries to adhere to. WHO and technical advisers provided technical support to countries when needed. The following principles were used for sampling in the WHS:

1. The sample should target the *de facto* population (i.e. all people living in that country including guest workers, immigrants and refugees) and not *de jure* population (i.e. the citizens of that country alone) with complete geographic coverage of the country, if possible, or a representative geographic segment of the country.
2. A *sample frame* with 90% coverage that included all key subgroups of interest. Enumeration or listing of households to update the frame was recommended.
3. The WHS sample identified all adult members of the general population aged 18 years or older as its sample. Households are selected using a random, stratified sampling procedure. One individual per household is then selected through a random selection procedure using the Kish table method.
4. The WHS used a sampling strategy which encompasses a known non-zero selection probability for any individual included in the study. This makes it possible to extrapolate the data to the whole population. Households and individuals are selected with a known probability in order to provide selection probabilities for each household and individual in the data set.
5. The WHS sampling strategy was without replacement and categories of non-respondents were clearly defined.
6. The WHS used GPS devices in certain countries and digitized maps to geocode the data within specified guidelines.

Each national report will present a detailed description of the sampling strategy and the allocation tables

2.2.1 Geographic Component of the Survey

Geographic Information refers to any information associated with a location on the Earth's surface. This location can be associated to a coordinate (latitude/longitude), a place's name, a postal address or any other geographic object.

Household Surveys represent an important source of information. If the geographic location of the surveyed clusters is collected at the same time that the interview is conducted users are then able to integrate the spatial dimension in their analysis

The geographic component of the WHO WHS is stored in two sections of the questionnaire:


- 0100 Sampling Information allowing the link between the location of each cluster and the sampling frame used for the survey. This section has also been used in order to collect the type of setting in which each household was located at the time of the survey. This section of the questionnaire has been used for all the countries part of the WHO WHS
- 0200 Geocoding Information for the collection of the latitude and longitude of each household. This section has only been used in 27 countries where GPS devices were available or have been provided. The choice of these countries has been done on the basis of the availability of digital maps delimiting the surveyed clusters, the field data collection planning and their distribution within the different WHO regions. Specific training and field material have been prepared and used by the interviewers or field supervisors in these countries in order to collect the latitude and longitude of each of the surveyed household.

As the sampling design may not always include the administrative structure of the country and as the codes entered in the data set for the sampling level needed to be labelled, a key correspondence table containing this information has been collected after the survey itself.

Once all the survey data and the key correspondence table available a specific protocol has been applied to the two sections mentioned earlier in order to clean eventual data collection or data entry mistakes as well as to fill the gaps in close collaboration with the survey institution.


The questionnaire as well as the training material and data cleaning protocols used can be downloaded from the WHO WHS web site (<http://www.who.int/healthinfo/survey/en/index.html>) in the "Survey Instrument and Related Documents" section.

2.3.1 Sample deviation index



SDI

2.3.2 HH Sample deviation index



HDI

2.3.3 Response rate


Household

Selected	929
Interviewed	924
Percentages	100


Individual

Selected	911
Interviewed	856
Percentages	94

2.3.4 Missing data



2.3.5 Reliability



3.1 Household composition by residence and age groups according to income quintile

	Percent Rural	SE	Mean Number of									
			Household members	SE	Children < 2 years	SE	Children < 5years	SE	Children 5 -17 years	SE	Persons 18+ years	SE
Q1(poorest)	57.5	4.57	1.6	0.06	0.0	0.01	0.0	0.02	0.1	0.02	1.4	0.04
Q2	21.2	4.00	1.5	0.06	0.0	0.00	0.0	0.01	0.1	0.02	1.3	0.04
Q3	21.7	3.43	1.8	0.09	0.0	0.01	0.0	0.01	0.2	0.04	1.6	0.07
Q4	21.4	4.11	2.1	0.11	0.0	0.01	0.1	0.02	0.3	0.06	1.8	0.08
Q5(richest)	26.1	4.22	2.7	0.11	0.0	0.02	0.1	0.02	0.3	0.05	2.3	0.09
All	30.2	1.11	1.8	0.05	0.0	0.00	0.0	0.01	0.2	0.02	1.6	0.04

Note: table based on de jure members of household i.e. usual residents

3.2 Selected age distributions by income quintile

	Females Age 18–49	Persons Age 50+	Persons Age 60+
Q1 (poorest)	33	221	164
Q2	50	225	163
Q3	74	169	115
Q4	88	144	98
Q5 (richest)	77	94	47
All	323	852	587

Note: table based on de jure members of household i.e. usual residents

3.3 Characteristics of head of household by income quintile

	Mean Age of Head	SE	Median Education of Head	Percent Female Head	SE
Q1 (poorest)	63	1.12	5	60.6	4.36
Q2	58	1.91	5	67.2	4.28
Q3	53	2.11	5	61.9	5.06
Q4	51	1.83	5	57.6	5.09
Q5 (richest)	44	1.44	5	38.4	4.41
All	55	0.99	5	59.2	2.29

Education

1=no formal education

2=less than primary school

3=primary school completed

4=secondary school completed

5=high school (or equivalent) completed

6=college/pre-university/university completed

7=post graduate degree completed

Note: table based on de jure members of household i.e. usual residents

3.4 Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex

	Male	SE	Female	SE	Missing	Total	SE
Age							
0-4	2.4	0.50	2.5	0.40		2.40	0.40
5-14	8.6	1.10	5.7	0.60		6.90	0.60
15-29	23.1	1.90	13.8	1.30		17.70	1.30
30-44	17.2	1.40	14.4	1.10		15.50	1.00
45-59	21.4	2.10	23.1	1.60		22.40	1.40
60-69	16.5	2.00	15.3	1.40		15.80	1.30
70-79	7.9	1.10	19.7	1.70		14.80	1.20
80+	2.9	0.80	5.5	1.10		4.40	0.70
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Residence							
Urban	67.3	1.50	67.0	1.10		67.10	0.20
Rural	32.7	1.50	33.0	1.10		32.90	0.20
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Marital_Status							
Never	37.1	2.10	23.8	1.80		29.30	1.50
Currently	37.0	2.10	26.3	1.50		30.80	1.60
Separated	1.7	0.70	3.3	0.80		2.60	0.60
Divorced	10.6	1.60	13.0	1.30		12.00	1.10
Widowed	6.8	1.50	28.9	1.90		19.80	1.50
CoHabiting	6.7	1.00	4.5	0.70		5.40	0.80
Missing	0.1	0.10	0.2	0.10		0.20	0.10
Education							
NoFormal	5.9	0.90	5.1	0.90		5.40	0.70
LessthanPrimary	8.8	1.10	9.9	1.00		9.40	0.80
Primary	21.2	2.10	13.0	1.40		16.40	1.40
Secondary	6.7	0.90	6.7	1.00		6.70	0.70
HighSchool	47.6	2.30	52.2	2.00		50.30	1.80
College	9.5	1.10	12.8	1.50		11.50	1.10
PostGraduate	0.1	0.10	0.2	0.10		0.20	0.10
Missing	0.1	0.10	0.2	0.10		0.10	0.10
Total	100.0		100.0			100.00	
Number	921.0		1,307.0			2,228.00	
Number of Households						912.00	

3.5 Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education

	Male	SE	Female	SE	Missing	Total	SE
Age							
18-29	24.9	3.10	19.5	2.20		21.90	2.00
30-44	29.4	3.50	24.8	2.10		26.90	2.10
45-59	24.1	3.20	23.5	2.10		23.80	2.00
60-69	12.5	1.90	15.2	1.80		14.00	1.50
70-79	7.1	1.40	12.2	1.20		9.90	1.00
80+	2.0	0.80	4.9	1.20		3.60	0.80
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Residence							
Urban	65.2	3.20	68.0	1.50		66.70	1.00
Rural	34.8	3.20	32.0	1.50		33.30	1.00
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Marital_Status							
Never	28.2	3.00	17.1	2.20		22.10	1.90
Currently	43.6	3.30	35.1	2.30		38.90	2.00
Separated	2.0	0.90	4.3	1.20		3.30	0.80
Divorced	12.5	2.10	14.4	1.60		13.60	1.30
Widowed	6.6	1.90	23.3	2.10		15.80	1.60
CoHabiting	7.1	1.80	5.7	1.10		6.30	1.00
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Education							
NoFormal	0.2	0.20	0.2	0.10		0.20	0.10
LessthanPrimary	3.4	1.10	4.7	1.00		4.10	0.70
Primary	24.8	3.20	16.4	1.90		20.20	1.90
Secondary	40.5	3.60	44.3	2.70		42.60	2.40
HighSchool	20.1	2.70	21.0	2.10		20.60	1.90
College	11.0	1.90	12.8	1.80		12.00	1.50
PostGraduate	0.0	0.00	0.5	0.30		0.30	0.10
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Total	100.0		100.0			100.00	
Number	381.0		474.0			855.00	

4.1 Prevalence of smoking and average daily tobacco consumption

	Current daily smoker	SE	Smoker, not daily	SE	Non-smoker	SE	Missing	Average daily tobacco consumption*	SE	Number of respondents
Sex										
Male	55.8	3.60	8.4	1.90	35.6	3.30	0.2	18.2	0.80	381
Female	18.5	1.80	5.6	1.00	75.8	2.00	0.0	11.9	0.80	474
Missing										0
Residence										
Urban	36.4	2.70	7.2	1.30	56.3	2.50	0.1	15.8	0.80	571
Rural	32.6	3.70	6.2	1.70	61.2	3.80	0.0	17.5	1.10	284
Missing										0
Income Quintile										
Q1	36.7	3.60	7.7	2.50	55.5	3.70	0.0	17.8	1.20	157
Q2	45.5	5.70	4.0	1.70	50.5	5.40	0.0	16.5	1.20	179
Q3	36.5	5.10	4.2	1.60	59.3	4.90	0.0	15.8	1.20	159
Q4	28.9	4.00	8.7	2.30	62.0	4.00	0.4	15.9	1.70	174
Q5	28.8	4.60	9.1	2.30	62.1	4.50	0.0	15.4	1.40	184
Missing										2
Age										
18-29	42.0	4.50	8.9	2.30	49.1	4.30	0.0	14.9	1.00	187
30-44	48.5	4.10	7.7	1.90	43.8	3.60	0.0	18.0	1.10	230
45-59	33.6	4.50	8.5	2.30	57.9	4.40	0.0	15.6	1.30	203
60-69	29.8	5.10	4.3	1.70	65.8	4.90	0.0	16.0	1.40	119
70-79	6.4	2.30	2.4	1.90	90.4	3.00	0.7	16.0	4.30	85
80+	3.9	2.80	0.0	0.00	96.1	2.80	0.0	3.1	1.00	31
Missing										0
All	35.1	2.20	6.9	1.00	57.9	2.10	0.1	16.3	0.60	855

* Denominator used is current daily smoker

4.2 Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers

	Life time abstainers	SE	Non-heavy drinkers	SE	Infrequent Heavy Drinkers*	SE	Frequent Heavy drinkers**	SE	Missing	Number of respondents
Sex										
Male	5.7	1.50	74.3	3.40	15.8	2.80	4.2	1.80	0.0	381
Female	13.9	1.90	82.8	2.00	2.7	0.90	0.1	0.10	0.5	474
Missing										0
Residence										
Urban	9.9	1.50	79.2	2.20	8.4	1.70	2.3	1.10	0.1	571
Rural	10.9	2.60	78.5	3.70	8.8	2.10	1.3	1.20	0.5	284
Missing										0
Income Quintile										
Q1	12.8	2.80	72.8	4.10	12.5	3.20	2.0	1.20	0.0	157
Q2	5.5	1.50	80.5	4.00	11.5	3.80	1.7	1.10	0.8	179
Q3	10.9	2.60	79.9	4.30	5.7	2.50	3.4	3.40	0.0	159
Q4	11.3	3.20	80.2	3.90	8.0	2.70	0.0	0.00	0.5	174
Q5	11.3	2.70	80.6	4.10	5.4	1.90	2.7	1.90	0.0	184
Missing										2
Age										
18-29	6.3	2.40	77.6	4.70	8.9	2.40	6.7	3.40	0.5	187
30-44	5.8	1.60	81.5	3.40	11.8	3.10	0.6	0.60	0.3	230
45-59	10.4	2.70	77.0	4.40	11.5	3.70	0.8	0.50	0.4	203
60-69	10.9	2.50	84.4	3.20	3.8	2.20	0.9	0.90	0.0	119
70-79	19.1	3.50	79.1	3.60	1.8	1.40	0.0	0.00	0.0	85
80+	40.0	10.90	60.0	10.90	0.0	0.00	0.0	0.00	0.0	31
Missing										0
All	10.3	1.30	79.0	1.90	8.6	1.30	1.9	0.80	0.3	855

*Infrequent heavy drinkers: 1-2 days with 5 or more standard drinks per week(last 7 days)

**Frequent heavy drinkers: 3 or more days with 5 or more standard drinks per week(last 7 days)

4.3 Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

	Fruit and vegetable intake						Physical Activity					Number of respondents
	*Insufficient	SE	Sufficient	SE	Missing	**Insufficient	SE	Sufficient	SE	Missing		
Sex												
Male	73.9	4.00	26.1	4.00	0.0	0.0	0.00	0.0	0.00	100.0	381	
Female	77.3	2.70	22.7	2.70	0.0						474	
Missing											0	
Residence												
Urban	77.1	3.20	22.9	3.20	0.0						571	
Rural	73.1	4.10	26.9	4.10	0.0						284	
Missing											0	
Income Quintile												
Q1	80.9	3.80	19.1	3.80	0.0						157	
Q2	82.2	4.30	17.8	4.30	0.0						179	
Q3	74.7	5.00	25.3	5.00	0.0						159	
Q4	74.8	3.90	25.2	3.90	0.0						174	
Q5	66.8	5.70	33.2	5.70	0.0						184	
Missing											2	
Age												
18-29	68.2	4.80	31.8	4.80	0.0						187	
30-44	75.2	3.90	24.8	3.90	0.0						230	
45-59	81.0	3.80	19.0	3.80	0.0						203	
60-69	72.7	5.10	27.3	5.10	0.0						119	
70-79	82.3	4.10	17.7	4.10	0.0						85	
80+	85.1	7.40	14.9	7.40	0.0						31	
Missing											0	
All	75.7	2.50	24.3	2.50	0.0						855	

*Insufficient intake of fruit/vegetables: less than 5 servings of fruit/vegetables per one typical day

**Insufficient physical activity: less than 150 minutes per week(last 7 days) spent on walking/moderate activity/vigorous activity

4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water

	Piped into households	SE	other improved sources	SE	Total improved sources	SE	No improved sources	SE	Missing	Number of households
Residence										
Urban	88.2	1.80	6.5	1.40	94.6	1.20	0.8	0.50	4.6	636
Rural	56.0	5.40	35.1	5.00	91.1	2.30	4.3	1.70	4.7	276
Missing										0
Income_Quintile										
Q1(poorest)	41.1	5.20	48.8	4.90	89.9	2.60	5.7	2.10	4.4	208
Q2	88.8	3.30	5.5	2.20	94.2	2.50	1.8	1.30	3.9	240
Q3	87.4	3.20	7.4	3.00	94.8	1.40	0.0	0.00	5.2	187
Q4	92.1	2.40	2.3	0.90	94.4	2.20	0.5	0.40	5.1	161
Q5(richest)	90.6	2.80	4.8	2.10	95.3	1.90	0.0	0.00	4.7	114
Missing										1
Total	78.4	2.10	15.1	1.90	93.5	1.10	1.9	0.60	4.6	912

other improved sources: public standpipe/protected tube well/borehole/protected dug well/protected spring/rainwater collection

no improved sources: unprotected well/unprotected spring/water taken directly from a pond or stream/tanker truck water/vendor provided water

4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility

	Flush toilet to sewage system	SE	Other improved toilet facilities	SE	Total improved sanitation	SE	No improved sanitation	SE	Missing	Number of households
Residence										
Urban	70.2	3.90	8.1	1.90	78.2	3.30	17.1	3.40	4.6	636
Rural	51.7	5.60	15.9	3.20	67.6	4.50	27.4	4.40	5.0	276
Missing										0
Income_Quintile										
Q1(poorest)	26.9	4.50	15.5	3.70	42.4	4.80	53.2	4.90	4.4	208
Q2	67.0	6.20	12.6	3.60	79.6	4.70	16.0	4.60	4.3	240
Q3	73.9	5.00	7.7	2.10	81.6	4.60	13.2	4.50	5.2	187
Q4	84.2	3.00	5.8	1.60	90.0	2.70	4.6	1.80	5.4	161
Q5(richest)	85.1	3.50	7.7	2.60	92.7	2.50	2.6	1.70	4.7	114
Missing										1
Total	64.6	3.20	10.4	1.60	75.0	2.60	20.2	2.70	4.8	912

other improved sanitation facilities: connection to septic system/pour-flush latrine/covered dry latrine(with privacy) [provided facilities not shared]
no improved sanitation: uncovered dry latrine(without privacy)/bucket latrine/no facilities(open defecation)

4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used

	Electricity, Gas	SE	Kerosene	SE	Solid fuels on improved stove	SE	Solid fuels on traditional stove	SE	Total solid fuel use	SE	Missing	Number of households
Residence												
Urban	92.0	1.50	0.2	0.20	3.1	1.00	0.2	0.10	3.2	1.00	4.6	636
Rural	70.4	4.00	0.0	0.00	22.6	3.80	2.4	1.00	24.9	4.00	4.7	276
Missing												0
Income_ Quintile												
Q1(poorest)	60.3	4.80	0.0	0.00	32.7	4.60	2.6	1.30	35.3	4.60	4.4	208
Q2	91.5	2.60	0.4	0.40	3.5	1.30	0.7	0.50	4.2	1.50	3.9	240
Q3	93.4	1.60	0.0	0.00	1.4	0.80	0.0	0.00	1.4	0.80	5.2	187
Q4	93.1	2.40	0.0	0.00	1.5	1.10	0.3	0.30	1.8	1.10	5.1	161
Q5(richest)	95.3	1.90	0.0	0.00	0.0	0.00	0.0	0.00	0.0	0.00	4.7	114
Missing												1
Total	85.5	1.60	0.1	0.10	9.0	1.40	0.8	0.30	9.8	1.40	4.6	912

5.01 Tuberculosis

	Total	Tuberculosis		
		Need	Coverage	SE
Sex				
Female	474	9.3	40.0	7.5
Male	381	9.5	35.4	10.7
Missing	0			
Residence				
Urban	571	10.2	39.6	7.8
Rural	284	7.8	33.5	9.9
Missing	0			
Income_Quintile				
Q1(poorest)	157	11.7	26.2	8.7
Q2	179	9.0	37.7	16.5
Q3	159	14.3	37.1	12.8
Q4	174	5.4	59.9	13.5
Q5(richest)	184	7.6	40.3	14.3
Missing	2			
Age				
18-29	187	5.9	46.2	14.8
30-44	230	8.8	24.3	11.3
45-59	203	11.0	54.1	13.6
60-69	119	12.9	36.5	12.1
70-79	85	10.4	32.5	12.2
80+	31	9.3	0.0	0.0
Missing	0			
Total	855	9.4	37.9	6.3

M = % Missing of Covered SE= Standard Error of % Covered

Tuberculosis Indicators:

"Need" = % of all respondents with self-reported continuous cough lasting 3 weeks, OR had blood in phlegm, OR had coughed blood in the last 12 months

"Coverage" = % of respondents with self reported symptoms who indicated receiving a TB screening test in the last 12 months.

5.02 HIV Prevention and Screening

Insufficient / No data available

5.03 Antenatal Care and Mother to Child Transmission of HIV

	Antenatal Care											Maternal to Child HIV Trans.				
	Women 18-49	Birth in last 5 years	Cov	SE	Visit 1x	Visit 2x	Visit 3x	Blood pressure checked	Blood sample taken	Told Signs of Pregnancy Complications	M	Birth in last 2 years	Cov	SE	M	
Residence																
Urban	166	20.8	42.8	10.2	11.7	11.7	11.7	0.0	1.8	47.4	13.4	20.0	27.5	8.2	0.0	
Rural	78	20.8	71.7	15.9	0.0	0.0	0.0	0.0	0.0	71.7	0.0	23.8	30.8	10.2	0.0	
Missing																
Income_Quintile																
Q1(poorest)	27	31.0	18.2	17.6	12.2	12.2	12.2	0.0	0.0	18.2	12.2	31.0	17.9	12.4	0.0	
Q2	35	10.2	50.6	21.8	0.0	0.0	0.0	0.0	0.0	50.6	0.0	10.2	0.0	0.0	0.0	
Q3	58	27.8	56.4	17.5	6.5	6.5	6.5	0.0	0.0	62.8	6.5	27.0	33.5	14.0	0.0	
Q4	60	19.1	57.1	17.0	4.9	4.9	4.9	0.0	5.3	62.0	10.2	19.1	40.2	16.5	0.0	
Q5(richest)	62	21.9	67.2	14.0	10.4	10.4	10.4	0.0	0.0	67.2	10.4	20.4	27.5	13.5	0.0	
Missing	2															
Total	244	21.8	52.9	9.0	7.6	7.6	7.6	0.0	1.1	55.9	8.8	21.2	28.7	6.4	0.0	

M = % Missing of Covered SE= Standard Error of %Covered

Antenatal Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least 3 times or more by health care professional, received blood pressure, blood sample taken, and told signs about complications for their last pregnancy (aggregate antenatal care indicator).

"Visit 1x" "Visit 2x" "Visit 3x" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least once, twice or three times or more respectively by a health care professional during their last pregnancy.

"Blood pressure checked" = % of female respondents of reproductive age with live birth in the last 5 years who had blood pressure taken when they saw health care professional during their last pregnancy.

"Blood sample taken" = % of female respondents of reproductive age with live birth in the last 5 years who had a blood sample taken when they saw health care professional during their last pregnancy.

"Told Signs of Pregnancy Complications" = % of female respondents of reproductive age with live birth in the last 5 years who had been told signs of pregnancy complications when they saw health care professional during their last pregnancy.

Maternal to Child HIV Transmission Indicators:

"Birth in last 2 years" = % of female respondents of reproductive age indicating having had a live birth in the last 2 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 2 years who indicated having been counseled and tested and given results for HIV during their last pregnancy.

5.04 Birth Delivery Care

	Women 18-49	Delivery Care				Place of Delivery						
		Need	Coverage	SE	M	Public Health Facility	SE	Private Health Facility	SE	Home	SE	M
Residence												
Urban	166	20.8	100.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Rural	78	20.8	91.8	7.1	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Missing												
Income Quintile												
Q1(poorest)	27	31.0	82.1	12.4	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Q2	35	10.2	100.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Q3	58	27.8	100.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Q4	60	19.1	100.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Q5(richest)	62	21.9	100.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Missing	2											
Total	244	21.8	97.1	2.7	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0

M = % Missing of Covered SE= Standard Error of % Covered Birth Delivery Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Coverage" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having received assistance with birth delivery from a health care professional (doctor, nurse or midwife) during their last birth.

"Public Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a public health facility.

"Private Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a private health facility.

"Home" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered home or in the field (etc.)

5.09 Women's Cancer Screening

	Total Females	Percent Females 18+	PAP Smear	SE	M	Percent Females 40+	Mammography / Breast Exam	SE	M
Residence									
Urban	322	83.2	72.4	3.3	5.2	47.8	36.8	4.3	10.1
Rural	152	82.2	75.9	3.7	4.3	45.3	32.0	6.3	9.9
Missing	0								
Income_Quintile									
Q1(poorest)	81	68.7	56.4	7.1	12.5	39.8	22.0	6.6	23.3
Q2	99	74.5	66.6	6.6	3.8	51.0	25.7	5.9	12.9
Q3	106	83.8	77.5	4.9	4.0	49.6	40.7	8.3	6.3
Q4	101	90.5	74.3	5.0	3.0	51.3	39.4	6.9	6.9
Q5(richest)	84	95.8	85.5	3.7	3.9	40.5	48.4	9.1	4.0
Missing	2								
Age									
18-29	92	19.5	77.2	6.1	4.4				
30-39	78	16.5	81.7	4.4	1.2				
40-49	74	15.6	82.3	4.1	4.8	15.6	38.5	6.6	6.9
50-59	77	16.2	74.1	4.7	5.0	16.2	35.7	5.9	8.8
60-69	72	15.2	50.0	6.4	9.6	15.2	31.6	5.0	14.6
70+	81	17.1		0.0		17.1		0.0	
Missing	0								
Total	474	82.9	73.5	2.5	4.9	47.0	35.3	3.5	10.0

M = % Missing of those with Pap Smear/Mammography SE= Standard Error of % with Pap Smear/Mammography

***Note: Values in the age row under the "Females 18 plus" and "Females 40 plus" columns are column percent values.

Women's Cancer Screening Indicators:

"Papsmear need"= % of all female respondents aged 18-69

"Papsmear" coverage = % of those female respondents aged 18-69 who self-reported receiving a pap smear test during pelvic examination in the last 3 years.

"Mammography need" = % of all female respondents aged 40-69

"Mammography" coverage = % of those female respondents aged 40-69 who self-reported receiving a mammography test or breast examination in the last 3 years.

****In several countries women above age 69 were asked the cancer screening questions, and for this reason they are included in the table. For countries that did not ask this questions of women above age 69, these values are expected to be missing.

5.10 Oral Health and Injuries

	Oral Health Problems					Road Traffic Accidents				Injury			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
Sex													
Female	474	48.4	77.9	3.2	0.0	1.4	9.2	9.3	0.0	6.9	22.9	6.7	0.0
Male	381	44.8	62.7	4.8	0.0	3.1	36.9	22.0	0.0	14.7	9.4	5.6	0.0
Missing	0												
Residence													
Urban	571	44.8	71.6	3.5	0.0	1.3	36.4	22.7	0.0	9.9	20.5	6.8	0.0
Rural	284	50.8	71.2	4.2	0.0	3.9	20.4	16.3	0.0	11.3	3.4	2.4	0.0
Missing	0												
Income_Quintile													
Q1(poorest)	157	45.2	55.7	7.1	0.0	0.3	0.0	0.0	0.0	12.4	13.7	8.1	0.0
Q2	179	42.6	61.6	5.7	0.0	3.1	0.0	0.0	0.0	12.7	6.5	4.3	0.0
Q3	159	44.7	78.3	5.0	0.0	3.3	10.3	10.7	0.0	8.5	33.5	21.2	0.0
Q4	174	54.4	76.4	5.2	0.0	2.5	100.0	0.0	0.0	6.0	10.1	7.3	0.0
Q5(richest)	184	47.0	82.4	5.1	0.0	1.4	0.0	0.0	0.0	12.2	13.2	6.8	0.0
Missing	2												
Age													
18-29	187	46.4	78.9	5.9	0.0	3.5	26.2	22.5	0.0	13.5	3.1	3.2	0.0
30-44	230	54.3	76.3	4.6	0.0	4.1	22.0	21.0	0.0	7.7	19.4	11.1	0.0
45-59	203	47.1	67.5	5.7	0.0	0.6	54.1	35.1	0.0	12.8	17.4	10.8	0.0
60-69	119	45.8	62.1	6.6	0.0	0.5	100.0	0.0	0.0	10.0	16.6	10.1	0.0
70-79	85	40.0	61.9	7.7	0.0	0.0	0.0	0.0	0.0	5.0	47.6	17.9	0.0
80+	31	13.1	62.2	22.1	0.0	2.5	0.0	0.0	0.0	12.0	0.0	0.0	0.0
Missing	0												
Total	855	46.8	71.4	2.7	0.0	2.2	26.9	14.2		10.4	14.3	4.4	

M = % Missing of % Covered SE= Standard Error of % Covered

Indicators:

Oral Health:

"Need" = % of all respondents indicating any problems in the last 12 months with their mouth and/or teeth.

"Coverage" = % of respondents reporting problems with their mouth and teeth who indicated receiving dental or medical care in last 12 months.

Road Traffic Accident:

"Need" = % of all respondents indicating having suffered bodily injury from a road traffic accident in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury from a road traffic accident in last 12 months who indicating having received treatment for their injuries on-site or at health facility within 1 hour of accident .

Injuries from Non-traffic Accidents:

"Need" = % of all respondents indicating having suffered bodily injury due to non-traffic accidents in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury due to non-traffic accidents in last 12 months who indicating having received treatment for injuries on-site or at health facility within 1 hour of accident

5.11 Self Reported Non-Communicable Conditions

	Angina					Osteo-Arthritis				Asthma				Diabetes			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
Sex																	
Female	474	21.9	59.8	5.2	0.6	15.8	43.8	5.9	0.0	4.6	54.7	11.5	0.0	6.9	79.6	6.8	3.3
Male	381	14.6	69.2	7.5	0.0	9.8	58.4	10.6	0.0	3.5	42.4	18.3	0.0	3.7	82.8	10.2	3.3
Missing	0																
Residence																	
Urban	571	17.8	73.9	5.4	0.0	12.5	50.5	7.3	0.0	4.7	47.3	12.6	0.0	6.7	86.4	5.1	4.0
Rural	284	20.4	44.3	7.6	1.1	14.4	45.3	9.0	0.0	3.0	58.2	17.4	0.0	3.1	55.4	18.3	0.0
Missing	0																
Income_Quintile																	
Q1(poorest)	157	26.0	53.6	6.9	0.0	16.8	37.6	10.9	0.0	2.2	28.0	17.2	0.0	4.6	44.2	16.8	0.0
Q2	179	21.2	57.5	8.9	0.0	15.3	53.3	10.3	0.0	5.3	23.5	14.2	0.0	5.9	73.5	14.8	10.1
Q3	159	22.1	62.2	9.6	1.8	13.7	59.1	11.5	0.0	7.2	74.6	11.9	0.0	8.1	97.3	2.7	0.0
Q4	174	17.5	86.5	7.0	0.0	12.5	54.4	12.4	0.0	3.3	57.7	27.3	0.0	6.2	92.2	6.0	4.2
Q5(richest)	184	7.9	56.4	10.9	0.0	7.6	37.9	15.4	0.0	2.7	49.5	24.7	0.0	2.9	80.2	13.6	0.0
Missing	2																
Age																	
18-29	187	3.4	0.0	0.0	0.0	4.3	0.0	0.0	0.0	2.0	0.0	0.0	0.0	0.0	82.3	13.4	2.9
30-44	230	6.4	53.8	15.5	4.2	3.1	40.4	19.8	0.0	3.0	48.1	29.0	0.0	1.0	87.0	6.6	0.0
45-59	203	16.9	42.9	11.1	0.0	12.6	49.1	12.3	0.0	3.1	74.6	18.8	0.0	5.4	84.5	8.4	12.9
60-69	119	37.4	74.4	5.5	0.0	26.0	47.3	11.0	0.0	7.0	46.7	19.2	0.0	13.1	56.5	20.7	
70-79	85	47.6	72.3	6.0	0.0	37.4	60.2	8.1	0.0	9.7	60.5	12.7	0.0	11.1	0.0	0.0	
80+	31	62.1	81.4	9.4	0.0	27.1	62.6	18.6	0.0	4.9	36.6	32.8	0.0	27.2	0.0	0.0	
Missing	0																
Total	855	18.6	63.1	4.6	0.4	13.1	48.6	5.7	0.0	4.1	49.9	10.7	0.0	5.5	80.6	5.8	3.3

M = % Missing for Covered

SE= Standard Error of %Covered

Self Reported Non-Communicable Conditions Indicators:

"Need" = % of total respondents indicating having received a diagnosis of the condition (angina, arthritis, asthma, diabetes) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (angina, arthritis, asthma, diabetes) indicating they have received treatment or screening for the condition in the last 2 weeks.

5.12 Mental Health Conditions

	Total	Depression				Schizophrenia			
		Need	Coverage	SE	M	Need	Coverage	SE	M
Sex									
Female	474	7.9	48.3	8.3	0.0	0.2	39.0	33.6	0.0
Male	381	2.7	54.4	21.5	0.0	1.2	60.6	29.6	0.0
Missing	0								
Residence									
Urban	571	6.0	44.0	8.7	0.0	0.7	85.3	15.0	0.0
Rural	284	4.7	64.3	15.6	0.0	0.7	0.0	0.0	0.0
Missing	0								
Income_Quintile									
Q1(poorest)	157	7.0	50.6	18.6	0.0	1.8	100.0	0.0	0.0
Q2	179	7.3	36.3	13.0	0.0	1.3	0.0	0.0	0.0
Q3	159	4.8	47.7	22.7	0.0	0.2	100.0	0.0	0.0
Q4	174	4.9	68.9	17.5	0.0	0.0	0.0	0.0	
Q5(richest)	184	3.9	55.0	16.1	0.0	0.0	0.0	0.0	0.0
Missing	2								
Age									
18-29	187	1.5	100.0	0.0	0.0	0.0	0.0	0.0	
30-44	230	5.4	35.8	16.3	0.0	1.9	43.6	31.6	0.0
45-59	203	6.9	53.1	16.1	0.0	0.5	100.0	0.0	0.0
60-69	119	6.6	34.0	14.9	0.0	0.0	0.0	0.0	0.0
70-79	85	9.3	70.9	13.7	0.0	0.4	100.0	0.0	0.0
80+	31	9.3	26.1	22.5	0.0	0.0	0.0	0.0	0.0
Missing	0								
Total	855	5.6	49.7	8.0	0.0	0.7	57.1		0.0

M = % Missing for Covered

SE= Standard Error of %Covered

Mental Health Conditions Indicators:

"Need" = % of total respondents indicating having received a diagnosis of the condition (depression, schizophrenia) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (depression, schizophrenia) indicating they have received treatment or screening for the condition in the last 2 weeks.

5.13 Cataracts

		Cataracts			
	Total 60+	Need	Coverage	SE	M
Sex					
Female	153	20.4	41.3	9.1	0.0
Male	82	15.1	30.3	13.1	0.0
Missing	0				
Residence					
Urban	165	20.4	38.4	9.0	0.0
Rural	70	14.2	37.3	15.9	0.0
Missing	0				
Income_Quintile					
Q1(poorest)	67	14.2	52.4	15.2	0.0
Q2	61	16.1	65.3	12.6	0.0
Q3	46	20.4	29.8	13.3	0.0
Q4	44	25.5	14.0	10.4	0.0
Q5(richest)	18	21.2	24.0	20.6	0.0
Missing	0				
Age					
60-69	119	10.2	28.8	12.5	0.0
70-79	85	18.6	45.3	10.5	0.0
80+	31	50.9	38.3	15.1	0.0
Missing	0				
Total	235	18.5	38.2	7.8	0.0

M = % Missing of Covered SE= Standard Error of %Covered

Cataracts Indicators:

Total = Number of respondents 60 years of age or more.

"Need" = % in need (respondents 60 years of age or more with self-reported diagnosis of cataracts in the last 5 years).

"Coverage" = % respondents 60 years of age or more with self-reported diagnosis of cataracts having indicated receiving eye surgery to remove cataracts in the last 5 years)

6.01 In general, rate your health today - q2000

	Very Good	SE	Good	SE	Moderate	SE	Bad	SE	Very Bad	SE	Missing	SE	N
Gender													
Male	8.8	2.3	37.8	3.4	34.3	3.2	16.4	2.5	2.7	1.0	0.0	0.0	381
Female	3.1	0.9	29.9	2.4	48.5	2.5	15.4	1.5	3.2	0.7	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	6.1	1.6	30.1	2.4	45.9	2.8	14.9	1.7	3.0	0.7	0.0	0.0	571
Rural	4.6	1.6	40.1	3.9	34.7	3.3	17.7	2.6	2.9	1.1	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	2.5	1.3	23.5	3.8	44.2	4.2	22.4	3.6	7.4	2.2	0.0	0.0	157
Q2	4.1	3.2	25.4	4.4	44.9	5.1	23.5	4.0	2.1	0.9	0.0	0.0	179
Q3	3.8	1.8	28.0	4.9	47.5	5.2	17.6	3.4	3.1	1.1	0.0	0.0	159
Q4	6.9	2.5	32.4	5.2	48.4	4.9	10.7	2.4	1.6	1.1	0.0	0.0	174
Q5(richest)	10.2	3.4	55.0	4.2	27.7	3.8	6.2	2.0	1.0	0.7	0.0	0.0	184
M	0.0	0.0	75.1	22.8	0.0	0.0	0.0	0.0	24.9	22.8	0.0	0.0	2
Age													
18-29	11.7	3.1	59.8	4.1	25.7	4.1	2.0	1.2	0.8	0.6	0.0	0.0	187
30-44	6.9	2.4	46.6	4.2	34.5	4.2	9.8	2.6	2.1	1.1	0.0	0.0	230
45-59	4.4	2.9	22.5	3.2	51.0	4.6	19.1	2.9	2.9	1.3	0.0	0.0	203
60-69	1.0	0.7	12.5	3.2	58.7	4.5	22.6	4.0	5.1	2.0	0.0	0.0	119
70-79	0.0	0.0	7.2	2.6	51.7	4.7	34.6	4.3	6.5	2.2	0.0	0.0	85
80+	0.0	0.0	0.0	0.0	50.6	10.8	44.4	11.0	5.0	3.5	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	5.6	1.2	33.4	2.1	42.2	2.2	15.8	1.4	3.0	0.6			855

6.02 Difficulty with 'work and household activities' in last 30 days - q2001

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	44.4	3.0	21.2	2.7	21.9	2.8	8.8	2.0	3.6	1.4	0.0	0.0	381
Female	32.2	2.3	23.3	2.0	28.8	2.1	12.5	1.5	2.4	0.7	0.8	0.6	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	40.4	2.2	22.1	2.0	24.2	2.0	9.9	1.5	2.7	0.8	0.7	0.5	571
Rural	32.2	3.3	22.9	3.4	28.8	3.6	12.8	2.3	3.3	1.4	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	23.8	3.7	19.6	3.8	32.0	3.9	16.6	2.9	6.4	2.5	1.5	1.5	157
Q2	34.2	4.6	21.5	3.9	28.0	4.3	14.7	3.8	1.7	0.8	0.0	0.0	179
Q3	40.3	5.0	16.1	3.1	26.1	4.2	13.6	2.8	3.5	2.0	0.4	0.4	159
Q4	40.1	3.9	27.3	3.8	23.2	3.6	5.9	2.0	3.1	1.5	0.3	0.3	174
Q5(richest)	48.1	4.8	26.3	3.5	20.5	3.6	4.3	1.3	0.6	0.6	0.2	0.2	184
M	43.1	30.8	32.0	27.6	0.0	0.0	24.9	22.8	0.0	0.0	0.0	0.0	2
Age													
18-29	63.3	4.6	23.0	3.9	11.5	3.2	1.3	0.7	0.9	0.9	0.0	0.0	187
30-44	41.0	4.1	28.2	4.4	20.7	3.5	6.5	2.3	2.2	1.2	1.3	1.0	230
45-59	32.2	4.1	22.2	3.2	30.9	3.8	11.6	2.6	3.0	1.7	0.2	0.2	203
60-69	25.2	4.2	20.3	4.2	33.2	4.2	16.2	3.5	4.7	1.8	0.5	0.5	119
70-79	13.4	3.6	13.2	3.4	42.3	4.9	26.5	3.7	4.6	1.8	0.0	0.0	85
80+	7.7	4.7	9.9	5.0	40.8	10.8	32.7	9.8	8.9	5.0	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	37.6	1.8	22.4	1.8	25.7	1.8	10.9	1.2	2.9	0.7	0.5	0.3	855

6.03 Difficulty with 'moving around' in last 30 days - q2010

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	61.9	3.3	17.3	2.9	8.7	1.8	8.4	1.7	3.7	1.4	0.0	0.0	381
Female	59.0	2.4	13.3	1.7	14.6	1.6	9.7	1.3	3.2	0.8	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	61.1	2.3	15.1	2.1	11.7	1.4	8.3	1.2	3.9	1.0	0.0	0.0	571
Rural	58.8	2.6	15.1	2.3	12.6	2.3	11.0	2.2	2.6	1.2	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	44.3	4.5	19.2	4.1	18.3	3.2	13.3	2.6	5.0	2.3	0.0	0.0	157
Q2	51.2	4.5	16.6	4.4	16.7	3.6	11.2	2.8	4.3	2.0	0.0	0.0	179
Q3	63.3	4.5	11.0	3.1	10.0	2.5	10.7	2.3	4.9	2.1	0.0	0.0	159
Q4	65.9	3.9	12.6	2.8	10.3	2.6	8.4	2.5	2.8	1.4	0.0	0.0	174
Q5(richest)	75.0	3.5	16.1	3.5	5.1	1.7	3.1	1.1	0.6	0.6	0.0	0.0	184
M	75.1	22.8	0.0	0.0	24.9	22.8	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	89.7	2.9	7.4	2.6	1.3	0.7	0.7	0.7	0.9	0.9	0.0	0.0	187
30-44	67.9	4.3	17.5	3.3	6.6	2.2	4.8	1.8	3.2	1.5	0.0	0.0	230
45-59	59.7	4.1	20.1	3.8	9.1	2.3	7.4	2.2	3.6	1.7	0.0	0.0	203
60-69	41.4	4.4	16.5	3.7	22.7	3.2	14.8	3.0	4.7	1.7	0.0	0.0	119
70-79	21.9	4.5	13.7	3.2	33.1	5.3	26.5	4.3	4.8	1.5	0.0	0.0	85
80+	8.8	5.2	8.5	4.3	36.7	9.9	34.9	8.9	11.2	7.9	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	60.3	1.8	15.1	1.6	12.0	1.2	9.2	1.1	3.4	0.8			855

6.04 Difficulty with 'vigorous activities' in last 30 days - q2011

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	42.3	3.7	22.5	3.7	15.5	2.6	8.4	1.9	10.5	2.1	0.7	0.6	381
Female	36.9	2.4	18.5	1.8	15.3	1.6	14.2	1.7	14.0	1.8	1.2	0.8	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	41.6	2.5	17.6	2.1	15.1	2.0	11.5	1.5	12.8	1.7	1.5	0.8	571
Rural	34.8	3.7	25.7	4.4	16.0	2.7	11.8	2.1	11.7	2.4	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	29.0	4.1	19.2	3.6	19.0	3.3	15.9	3.1	14.4	3.3	2.5	1.7	157
Q2	36.2	4.6	17.5	4.2	19.5	4.6	10.0	2.5	15.9	3.0	0.8	0.7	179
Q3	41.4	4.8	11.9	2.6	9.7	2.4	19.6	3.6	17.0	3.3	0.4	0.4	159
Q4	40.8	4.5	23.3	3.8	14.2	3.2	9.4	2.5	11.0	3.5	1.4	1.2	174
Q5(richest)	47.9	5.2	28.2	6.1	14.5	3.2	4.5	1.4	5.0	1.4	0.0	0.0	184
M	43.1	30.8	32.0	27.6	0.0	0.0	24.9	22.8	0.0	0.0	0.0	0.0	2
Age													
18-29	68.2	5.0	18.6	4.6	9.0	2.7	2.4	1.1	1.3	1.0	0.4	0.4	187
30-44	46.7	4.1	28.1	3.8	10.9	2.6	5.2	1.6	8.0	2.2	1.2	0.9	230
45-59	31.0	4.0	23.8	3.5	18.3	3.3	16.7	3.7	10.2	2.4	0.0	0.0	203
60-69	21.1	3.4	15.6	3.4	23.1	4.6	17.7	3.2	21.2	4.3	1.4	1.0	119
70-79	10.9	3.0	7.5	2.7	24.7	4.6	23.0	4.4	32.8	4.6	1.1	1.1	85
80+	12.6	9.0	2.3	2.3	12.7	5.5	26.8	8.3	37.9	10.4	7.8	7.4	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	39.3	2.1	20.3	2.0	15.4	1.6	11.6	1.2	12.4	1.4	1.0	0.5	855

6.05 Difficulty with 'self-care' in last 30 days - q2020

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	83.8	2.3	6.9	1.7	6.7	1.7	2.0	0.8	0.6	0.5	0.0	0.0	381
Female	79.9	2.0	6.7	1.1	7.8	1.3	4.4	1.1	1.2	0.5	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	82.8	1.8	6.8	1.0	6.5	1.2	2.6	0.8	1.3	0.5	0.0	0.0	571
Rural	79.2	3.0	6.8	2.0	9.0	1.8	4.7	1.6	0.3	0.2	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	71.5	3.7	9.3	2.7	11.6	2.8	6.1	2.0	1.5	1.0	0.0	0.0	157
Q2	81.1	3.9	7.1	2.9	8.3	2.3	3.1	1.9	0.5	0.5	0.0	0.0	179
Q3	78.0	3.8	8.3	2.2	8.1	2.8	3.8	1.3	1.9	1.0	0.0	0.0	159
Q4	83.0	3.6	6.6	2.1	5.8	2.2	3.5	1.8	1.1	0.9	0.0	0.0	174
Q5(richest)	92.5	1.9	3.4	1.2	3.5	1.5	0.6	0.6	0.0	0.0	0.0	0.0	184
M	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	97.1	1.5	1.5	1.1	0.9	0.9	0.0	0.0	0.4	0.4	0.0	0.0	187
30-44	89.4	2.9	3.8	2.2	3.6	1.6	2.2	1.2	0.9	0.7	0.0	0.0	230
45-59	85.0	3.1	5.8	1.7	6.8	2.2	1.0	0.6	1.3	0.9	0.0	0.0	203
60-69	67.8	3.9	15.9	3.2	10.7	2.5	4.7	2.0	0.9	0.6	0.0	0.0	119
70-79	59.0	4.6	13.5	3.5	19.3	3.9	7.5	2.2	0.8	0.6	0.0	0.0	85
80+	22.9	7.8	13.7	5.6	30.8	9.5	29.5	11.2	3.1	3.1	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	81.6	1.6	6.8	1.0	7.3	1.0	3.3	0.7	1.0	0.3			855

6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	87.0	2.3	8.5	2.1	3.1	1.0	1.0	0.6	0.4	0.4	0.0	0.0	381
Female	83.1	1.9	7.6	1.3	5.5	1.1	3.0	1.0	0.9	0.5	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	86.6	1.6	7.6	1.4	3.5	0.8	1.8	0.6	0.6	0.4	0.0	0.0	571
Rural	81.4	3.0	8.8	2.4	6.2	1.6	2.7	1.3	0.8	0.6	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	72.6	3.7	14.8	3.4	7.4	1.8	4.0	1.7	1.2	0.9	0.0	0.0	157
Q2	84.8	3.7	8.7	3.1	5.3	1.9	1.2	1.2	0.0	0.0	0.0	0.0	179
Q3	80.6	3.9	9.8	3.0	7.0	2.3	1.2	0.7	1.5	1.0	0.0	0.0	159
Q4	89.2	2.9	3.8	1.6	2.9	1.3	3.1	1.8	0.9	0.9	0.0	0.0	174
Q5(richest)	94.8	1.6	3.9	1.4	0.3	0.3	1.0	0.7	0.0	0.0	0.0	0.0	184
M	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	97.0	1.6	2.1	1.3	0.9	0.9	0.0	0.0	0.0	0.0	0.0	0.0	187
30-44	89.8	2.9	4.8	2.4	1.2	0.7	2.7	1.3	1.5	0.9	0.0	0.0	230
45-59	88.9	2.6	6.9	2.3	3.4	1.3	0.0	0.0	0.8	0.8	0.0	0.0	203
60-69	76.4	3.7	14.7	3.1	5.9	2.2	3.0	1.6	0.0	0.0	0.0	0.0	119
70-79	66.8	4.0	16.4	3.5	12.9	3.2	3.1	1.4	0.8	0.5	0.0	0.0	85
80+	30.1	9.2	25.2	7.6	27.4	10.2	17.2	9.5	0.0	0.0	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	84.9	1.5	8.0	1.2	4.4	0.8	2.1	0.6	0.7	0.3			855

6.07 Amount of 'bodily aches or pains' in last 30 days - q2030

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	44.4	3.4	24.5	3.5	18.7	2.6	8.9	1.8	3.5	1.3	0.0	0.0	381
Female	36.0	2.5	25.6	2.4	24.4	2.0	9.0	1.2	5.0	1.0	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	42.4	2.6	25.4	2.6	19.7	2.1	7.6	1.2	4.9	1.0	0.0	0.0	571
Rural	34.4	3.6	24.5	3.2	26.2	3.2	11.8	2.2	3.1	1.1	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	27.3	4.3	21.3	4.2	29.3	3.9	14.7	3.0	7.4	2.6	0.0	0.0	157
Q2	36.7	4.8	25.9	4.7	20.9	3.7	10.7	2.8	5.8	2.0	0.0	0.0	179
Q3	43.8	5.3	18.9	3.9	22.3	3.9	10.4	2.6	4.6	2.1	0.0	0.0	159
Q4	43.3	4.4	23.6	3.6	25.5	3.8	4.9	1.6	2.8	1.2	0.0	0.0	174
Q5(richest)	46.7	4.5	34.3	4.8	12.7	2.9	5.0	1.6	1.3	0.9	0.0	0.0	184
M	32.0	27.6	43.1	30.8	0.0	0.0	24.9	22.8	0.0	0.0	0.0	0.0	2
Age													
18-29	57.8	5.3	28.4	5.2	11.0	2.9	1.5	1.1	1.3	0.7	0.0	0.0	187
30-44	45.0	4.2	30.2	4.7	14.3	2.8	7.1	1.9	3.3	1.5	0.0	0.0	230
45-59	42.2	4.5	21.9	3.3	20.7	3.3	10.5	2.5	4.7	1.9	0.0	0.0	203
60-69	22.1	3.6	22.8	4.1	36.5	3.8	13.5	2.9	5.1	1.9	0.0	0.0	119
70-79	12.9	3.2	19.8	4.5	42.3	5.3	18.9	3.6	6.2	1.8	0.0	0.0	85
80+	17.3	9.1	11.9	6.6	37.8	9.1	13.8	5.8	19.2	8.6	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	39.8	2.1	25.1	2.1	21.8	1.8	9.0	1.1	4.3	0.8			855

6.08 Amount of 'bodily discomfort' in last 30 days - q2031

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	39.7	3.6	29.8	3.9	18.4	2.6	8.5	1.8	3.4	1.3	0.2	0.2	381
Female	34.9	2.5	24.3	2.3	26.5	2.2	9.9	1.4	4.0	0.8	0.3	0.3	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	39.5	2.7	26.2	2.6	21.3	2.1	8.3	1.4	4.5	1.0	0.2	0.2	571
Rural	32.1	3.6	27.9	4.2	26.1	3.6	11.3	2.3	2.3	1.0	0.3	0.3	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	28.6	4.2	18.6	4.0	30.2	4.2	16.6	3.5	5.9	2.4	0.0	0.0	157
Q2	34.9	4.8	28.6	5.0	20.4	3.6	10.2	2.7	5.2	1.8	0.8	0.8	179
Q3	39.9	4.8	21.6	3.9	23.6	3.8	9.6	2.5	5.4	2.2	0.0	0.0	159
Q4	38.0	4.5	27.5	4.4	25.3	3.6	7.4	2.2	1.4	0.7	0.4	0.4	174
Q5(richest)	43.0	4.8	35.6	4.9	16.5	3.3	3.6	1.2	1.3	0.9	0.0	0.0	184
M	32.0	27.6	43.1	30.8	0.0	0.0	24.9	22.8	0.0	0.0	0.0	0.0	2
Age													
18-29	54.5	5.4	30.9	4.9	11.9	3.2	1.5	1.1	1.3	0.7	0.0	0.0	187
30-44	43.5	4.0	31.2	4.8	13.3	2.8	7.7	2.4	4.2	1.7	0.0	0.0	230
45-59	37.0	4.4	25.0	3.6	25.1	3.4	9.6	2.2	3.3	1.7	0.0	0.0	203
60-69	21.4	4.6	22.7	4.1	38.1	4.5	13.2	2.9	4.0	1.6	0.6	0.6	119
70-79	10.9	3.2	23.1	4.4	39.7	5.4	19.0	3.5	5.6	1.8	1.7	1.6	85
80+	15.5	9.0	6.3	4.4	42.1	9.1	24.6	9.3	11.5	5.6	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	37.0	2.2	26.8	2.3	22.9	1.8	9.3	1.2	3.7	0.7	0.3	0.2	855

6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	57.7	3.6	24.1	3.6	13.3	2.2	4.2	1.1	0.6	0.5	0.0	0.0	381
Female	49.7	2.6	25.3	2.2	16.6	1.8	7.3	1.1	1.2	0.6	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	54.6	3.3	25.6	2.8	13.7	1.8	5.3	1.0	0.9	0.4	0.0	0.0	571
Rural	50.6	3.4	23.1	3.6	18.1	2.9	7.2	1.5	1.0	0.8	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	47.1	4.7	20.7	3.6	17.2	3.5	13.2	2.6	1.8	1.2	0.0	0.0	157
Q2	48.8	5.2	27.1	4.7	18.2	3.4	5.9	1.7	0.0	0.0	0.0	0.0	179
Q3	47.9	5.2	25.6	4.5	19.5	3.9	5.5	2.0	1.5	1.1	0.0	0.0	159
Q4	62.8	4.1	22.4	3.9	10.9	2.4	2.3	1.1	1.7	0.9	0.0	0.0	174
Q5(richest)	58.1	4.4	27.5	4.2	10.9	2.6	3.6	1.4	0.0	0.0	0.0	0.0	184
M	68.0	27.6	32.0	27.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	70.9	4.9	22.8	4.3	3.3	1.3	3.0	1.5	0.0	0.0	0.0	0.0	187
30-44	56.7	5.1	32.4	5.2	6.6	1.9	2.2	1.1	2.1	1.1	0.0	0.0	230
45-59	54.6	4.9	21.2	3.6	20.5	4.0	3.8	1.5	0.0	0.0	0.0	0.0	203
60-69	47.9	5.2	19.4	3.8	24.8	4.0	7.4	2.3	0.5	0.5	0.0	0.0	119
70-79	25.4	4.0	24.9	4.4	31.9	4.6	15.9	3.6	1.9	1.2	0.0	0.0	85
80+	9.3	5.0	24.0	10.9	32.0	9.8	31.5	9.3	3.1	3.1	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	53.2	2.4	24.8	2.2	15.1	1.5	5.9	0.8	0.9	0.4			855

6.10 Difficulty with 'learning a new task' in last 30 days - q2051

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	59.4	3.7	25.7	3.6	9.3	1.7	3.8	1.1	1.4	0.7	0.4	0.3	381
Female	57.8	2.5	22.1	2.3	11.6	1.5	6.4	1.3	1.9	0.5	0.3	0.3	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	60.8	3.0	23.9	2.8	8.7	1.2	4.5	0.9	2.0	0.6	0.1	0.1	571
Rural	53.8	3.8	23.4	4.3	14.4	2.1	6.7	1.8	0.9	0.5	0.8	0.6	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	49.3	4.3	17.4	3.8	16.6	2.8	12.8	3.0	3.3	1.4	0.6	0.6	157
Q2	56.7	5.0	25.5	5.0	10.1	2.5	5.8	2.0	0.9	0.5	1.1	0.8	179
Q3	56.5	5.0	26.2	4.6	11.0	2.9	4.5	1.8	1.7	0.9	0.0	0.0	159
Q4	64.4	5.3	24.2	5.1	7.1	1.9	1.5	0.7	2.8	1.2	0.0	0.0	174
Q5(richest)	64.3	4.5	24.6	3.9	8.8	2.4	2.4	1.1	0.0	0.0	0.0	0.0	184
M	56.9	30.8	43.1	30.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	77.0	4.9	19.3	4.6	1.7	1.0	2.0	1.2	0.0	0.0	0.0	0.0	187
30-44	64.2	5.3	28.3	5.5	4.4	1.6	1.4	0.8	1.7	1.0	0.0	0.0	230
45-59	59.8	3.9	23.3	3.3	13.1	3.0	3.4	1.5	0.0	0.0	0.4	0.4	203
60-69	50.1	5.2	19.8	3.9	19.8	3.9	6.4	2.2	2.8	1.2	1.1	1.1	119
70-79	29.8	4.3	27.9	4.6	23.1	3.7	11.8	3.0	6.6	2.6	0.9	0.8	85
80+	6.3	4.2	22.3	10.8	23.8	8.6	43.2	10.5	4.5	3.4	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	58.5	2.4	23.7	2.4	10.6	1.1	5.2	0.8	1.7	0.4	0.3	0.2	855

6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	66.8	3.7	19.5	3.4	10.7	2.4	2.1	1.0	0.5	0.4	0.3	0.3	381
Female	65.7	2.3	20.9	1.9	8.1	1.4	2.9	0.9	1.9	0.6	0.4	0.2	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	70.3	2.8	20.7	2.6	6.3	1.5	1.3	0.6	1.1	0.5	0.3	0.2	571
Rural	58.1	4.2	19.5	3.3	15.3	2.9	5.0	1.6	1.6	0.8	0.5	0.3	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	55.8	4.6	20.8	3.7	14.8	3.3	6.9	2.1	0.9	0.9	0.8	0.8	157
Q2	65.9	5.3	23.6	4.8	8.7	2.9	1.1	0.6	0.7	0.6	0.0	0.0	179
Q3	68.0	4.2	16.7	3.4	8.8	2.8	4.5	2.2	1.9	1.0	0.0	0.0	159
Q4	69.6	5.0	21.0	4.9	6.3	2.3	0.6	0.4	1.8	1.5	0.8	0.6	174
Q5(richest)	71.1	4.0	19.2	3.5	8.1	2.8	0.4	0.3	1.0	0.8	0.0	0.0	184
M	32.0	27.6	0.0	0.0	43.1	30.8	0.0	0.0	0.0	0.0	24.9	22.8	2
Age													
18-29	73.4	5.0	16.1	3.9	9.4	2.9	0.0	0.0	1.0	0.7	0.0	0.0	187
30-44	61.1	5.0	28.9	5.2	7.0	2.4	0.7	0.4	1.8	0.9	0.5	0.5	230
45-59	67.9	4.0	15.9	3.3	10.6	2.5	3.8	1.8	1.3	0.9	0.5	0.4	203
60-69	70.7	4.4	18.3	3.5	6.3	2.1	3.9	1.7	0.8	0.6	0.0	0.0	119
70-79	58.8	4.6	22.7	4.0	12.4	3.4	4.7	2.0	0.4	0.4	1.0	1.0	85
80+	53.1	10.8	12.0	5.4	19.7	9.6	12.2	6.1	3.1	3.1	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	66.2	2.3	20.3	2.0	9.3	1.4	2.6	0.6	1.3	0.4	0.4	0.2	855

6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	62.5	4.0	20.7	3.1	14.2	3.2	2.2	1.1	0.0	0.0	0.3	0.3	381
Female	65.6	2.2	21.0	1.9	9.4	1.4	2.0	0.6	1.5	0.6	0.5	0.3	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	68.1	2.8	18.4	1.9	10.7	2.4	1.9	0.8	0.7	0.3	0.2	0.2	571
Rural	56.6	4.0	25.8	3.5	13.2	2.2	2.5	1.0	1.1	0.7	0.7	0.5	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	57.7	4.5	24.7	3.8	14.6	3.3	2.3	1.2	0.0	0.0	0.8	0.8	157
Q2	65.5	4.9	21.2	4.3	10.7	3.5	1.9	1.2	0.0	0.0	0.7	0.7	179
Q3	70.3	4.5	15.1	3.1	8.2	2.8	3.8	2.1	2.6	1.3	0.0	0.0	159
Q4	69.5	4.9	15.9	3.1	11.4	3.0	2.1	1.5	0.5	0.4	0.5	0.5	174
Q5(richest)	59.1	4.7	26.5	3.8	13.0	3.7	0.2	0.2	1.2	0.8	0.0	0.0	184
M	0.0	0.0	56.9	30.8	0.0	0.0	43.1	30.8	0.0	0.0	0.0	0.0	2
Age													
18-29	63.3	5.4	22.6	4.1	12.7	3.1	0.4	0.4	1.0	0.7	0.0	0.0	187
30-44	59.9	4.8	21.2	3.4	13.5	5.2	3.4	1.5	1.5	0.8	0.5	0.5	230
45-59	68.0	4.1	21.0	3.4	9.2	2.7	1.9	1.4	0.0	0.0	0.0	0.0	203
60-69	68.0	4.6	20.8	3.7	7.0	2.2	2.8	1.5	0.3	0.3	1.1	1.1	119
70-79	61.5	4.6	21.1	4.0	15.5	3.9	0.4	0.4	0.6	0.6	1.0	1.0	85
80+	71.1	9.6	6.8	4.3	12.9	7.6	6.1	4.4	3.1	3.1	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	64.2	2.3	20.9	1.8	11.6	1.8	2.1	0.6	0.8	0.3	0.4	0.2	855

6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	66.2	3.8	12.1	2.5	10.0	2.1	2.8	1.2	0.7	0.4	8.2	2.5	381
Female	65.8	2.4	11.8	1.6	8.4	1.4	4.4	0.9	1.9	0.5	7.8	1.6	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	65.1	3.0	12.7	1.7	8.7	1.3	3.2	0.8	1.5	0.5	8.7	2.2	571
Rural	67.7	3.5	10.5	2.1	9.9	2.5	4.6	1.5	1.0	0.5	6.4	2.3	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	56.3	4.0	15.1	3.3	17.8	3.9	3.9	1.6	1.5	0.8	5.5	2.1	157
Q2	60.7	5.5	10.6	3.6	11.6	2.6	4.8	1.6	2.0	1.0	10.4	3.9	179
Q3	59.1	4.9	13.9	3.2	5.6	1.7	5.6	2.3	2.5	1.1	13.3	4.0	159
Q4	75.9	3.5	10.0	2.2	4.7	1.8	2.1	1.0	1.0	0.7	6.4	2.3	174
Q5(richest)	75.9	4.0	10.4	2.7	6.7	2.1	2.4	1.0	0.0	0.0	4.7	2.1	184
M	75.1	22.8	24.9	22.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	71.8	4.9	11.7	3.4	5.7	2.2	0.6	0.6	0.0	0.0	10.2	3.1	187
30-44	77.6	3.7	6.2	1.8	5.2	2.2	0.9	0.5	0.6	0.4	9.6	2.6	230
45-59	64.5	4.5	13.5	3.4	7.6	2.1	3.7	1.8	1.4	1.0	9.3	2.6	203
60-69	58.7	5.3	16.9	3.1	14.0	3.6	5.9	2.0	1.2	0.9	3.3	1.7	119
70-79	47.7	4.6	15.8	3.5	16.1	3.8	11.4	3.1	4.1	1.6	4.9	2.9	85
80+	32.6	10.2	15.9	8.1	30.4	9.5	13.0	6.5	8.2	4.6	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	66.0	2.3	11.9	1.4	9.1	1.2	3.7	0.8	1.3	0.4	8.0	1.7	855

6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	72.3	3.5	6.8	1.5	8.5	2.0	3.4	1.2	0.9	0.8	8.0	2.5	381
Female	71.0	2.3	9.9	1.5	7.4	1.3	3.0	0.7	1.0	0.4	7.8	1.6	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	71.1	2.6	8.7	1.2	7.1	1.3	3.0	0.8	1.4	0.6	8.6	2.2	571
Rural	72.5	3.3	8.1	1.6	9.4	2.6	3.6	1.6	0.0	0.0	6.4	2.3	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	61.9	3.8	10.4	2.4	17.7	3.5	4.0	1.6	0.4	0.4	5.5	2.1	157
Q2	67.5	4.7	11.2	2.8	7.3	2.5	3.7	1.4	0.0	0.0	10.4	3.9	179
Q3	64.5	4.8	8.6	2.3	6.5	2.1	4.1	1.9	3.0	2.0	13.3	4.0	159
Q4	78.2	3.6	8.0	2.2	3.7	1.5	2.2	1.1	1.5	0.9	6.4	2.3	174
Q5(richest)	83.3	3.1	4.8	1.4	5.3	2.0	2.3	1.3	0.0	0.0	4.2	2.1	184
M	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	83.2	3.7	2.8	1.2	3.3	1.8	0.6	0.6	0.0	0.0	10.2	3.1	187
30-44	80.7	3.4	6.0	1.9	3.0	1.6	0.3	0.3	0.4	0.4	9.6	2.6	230
45-59	64.5	3.7	10.1	2.7	9.4	2.3	4.7	1.9	2.5	1.6	8.9	2.6	203
60-69	67.8	4.8	9.5	2.3	16.3	3.8	1.8	0.9	1.3	0.9	3.3	1.7	119
70-79	52.3	5.0	17.2	3.5	13.5	3.6	11.3	3.6	0.7	0.8	4.9	2.9	85
80+	47.5	10.1	23.6	9.2	14.3	6.1	14.6	7.8	0.0	0.0	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	71.6	2.0	8.5	1.0	7.9	1.2	3.2	0.7	0.9	0.4	7.9	1.7	855

6.15 Problem with 'sleeping' in last 30 days - q2080

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	52.0	3.6	22.7	3.6	15.3	2.3	7.8	1.7	2.3	1.1	0.0	0.0	381
Female	38.4	2.4	25.6	2.2	20.2	2.0	11.0	1.5	4.8	1.0	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	43.6	2.9	26.8	2.6	15.5	1.8	9.6	1.3	4.5	1.1	0.0	0.0	571
Rural	46.1	2.9	19.3	2.7	23.2	2.6	9.4	2.0	2.0	0.9	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	38.5	4.0	26.2	3.9	17.0	3.0	16.4	3.2	2.0	1.1	0.0	0.0	157
Q2	38.5	5.2	23.0	4.6	22.3	4.2	12.1	2.8	4.0	1.6	0.0	0.0	179
Q3	42.4	4.7	20.1	3.7	19.8	3.6	11.4	2.8	6.3	2.5	0.0	0.0	159
Q4	57.2	4.3	20.8	3.2	12.8	2.8	3.8	1.2	5.5	2.1	0.0	0.0	174
Q5(richest)	44.7	4.7	31.2	4.0	18.1	2.9	5.1	1.7	0.9	0.5	0.0	0.0	184
M	75.1	22.8	0.0	0.0	24.9	22.8	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	59.9	4.6	25.7	4.0	10.2	3.0	3.9	1.7	0.3	0.3	0.0	0.0	187
30-44	45.3	4.7	31.3	5.2	14.6	3.3	7.4	2.1	1.5	0.9	0.0	0.0	230
45-59	45.3	4.6	22.5	3.1	19.6	3.5	8.4	2.0	4.3	2.0	0.0	0.0	203
60-69	37.1	4.7	18.1	3.3	27.0	3.9	11.0	2.9	6.8	2.2	0.0	0.0	119
70-79	25.9	4.6	18.6	4.0	30.6	4.2	17.7	3.6	7.1	2.4	0.0	0.0	85
80+	18.3	7.0	15.5	7.1	12.3	5.2	39.3	9.7	14.6	9.1	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	44.4	2.2	24.3	2.0	18.0	1.5	9.5	1.1	3.7	0.8			855

6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	46.0	3.4	24.8	3.2	18.2	3.0	8.7	2.0	1.8	1.0	0.5	0.5	381
Female	31.1	2.5	30.7	2.6	24.2	2.2	11.2	1.6	2.9	0.8	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	39.0	2.9	29.7	2.4	18.9	2.0	9.4	1.5	2.6	0.8	0.4	0.4	571
Rural	35.3	3.7	24.8	3.6	26.8	4.1	11.3	2.6	1.9	0.8	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	36.0	4.2	25.3	3.8	21.4	3.8	14.9	4.2	2.4	1.3	0.0	0.0	157
Q2	36.0	4.8	33.2	5.1	19.4	3.4	9.1	2.6	2.3	1.0	0.0	0.0	179
Q3	37.8	5.1	22.3	3.9	20.0	3.4	14.3	3.1	4.3	2.2	1.3	1.3	159
Q4	43.9	4.2	28.0	3.6	19.9	3.3	5.7	2.2	2.5	1.5	0.0	0.0	174
Q5(richest)	34.5	5.1	30.7	4.4	26.7	5.0	7.3	2.2	0.8	0.8	0.0	0.0	184
M	75.1	22.8	0.0	0.0	24.9	22.8	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	53.4	4.9	26.5	3.9	16.1	3.8	2.8	1.4	1.2	0.9	0.0	0.0	187
30-44	36.4	4.6	33.2	4.6	20.2	3.4	7.1	2.5	2.1	1.3	0.9	0.9	230
45-59	37.4	4.2	27.8	3.7	22.7	3.7	8.9	2.7	3.2	1.6	0.0	0.0	203
60-69	32.5	4.5	26.2	3.5	23.0	3.3	14.2	3.0	4.1	1.7	0.0	0.0	119
70-79	21.9	3.9	25.4	4.4	33.0	4.3	17.6	3.8	2.2	1.2	0.0	0.0	85
80+	18.3	7.0	14.5	6.3	21.0	8.0	46.2	10.9	0.0	0.0	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	37.7	2.3	28.0	2.0	21.6	1.9	10.0	1.3	2.4	0.6	0.2	0.2	855

6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	50.6	3.3	20.5	3.5	20.8	2.9	5.9	1.5	2.2	1.1	0.0	0.0	381
Female	38.8	2.4	31.6	2.6	19.2	1.7	7.2	1.4	3.2	1.0	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	48.2	2.7	26.5	2.5	16.9	1.8	5.7	1.1	2.7	0.8	0.0	0.0	571
Rural	35.9	2.8	26.9	4.0	26.0	3.7	8.4	2.2	2.8	1.3	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	32.3	4.1	24.7	4.8	28.3	4.6	9.9	2.8	4.7	1.9	0.0	0.0	157
Q2	44.2	5.6	28.8	4.6	17.6	4.0	5.8	1.8	3.7	1.7	0.0	0.0	179
Q3	43.2	5.1	25.9	4.3	18.3	3.4	7.9	2.1	4.8	2.2	0.0	0.0	159
Q4	53.8	4.2	26.1	4.1	14.8	2.8	4.8	1.9	0.6	0.4	0.0	0.0	174
Q5(richest)	45.7	4.1	27.4	4.1	21.1	4.2	5.3	2.2	0.5	0.4	0.0	0.0	184
M	32.0	27.6	24.9	22.8	43.1	30.8	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	57.7	4.9	21.2	3.8	14.2	3.4	5.5	2.1	1.4	0.8	0.0	0.0	187
30-44	43.9	3.6	30.3	4.1	17.0	3.7	4.4	1.7	4.3	1.9	0.0	0.0	230
45-59	44.1	4.6	28.1	3.8	18.5	3.4	6.2	1.9	3.1	1.6	0.0	0.0	203
60-69	43.6	5.0	21.3	4.0	26.4	4.0	7.8	2.5	0.9	0.6	0.0	0.0	119
70-79	23.4	3.8	34.5	4.9	27.6	4.0	11.2	3.0	3.4	2.0	0.0	0.0	85
80+	21.4	9.2	21.3	8.9	39.9	10.1	15.6	6.8	1.8	1.8	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	44.1	2.1	26.6	2.1	19.9	1.7	6.6	1.0	2.7	0.7			855

6.18 Problem with 'worry or anxiety' in last 30 days - q2091

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	35.7	3.2	31.9	3.0	22.8	2.9	6.0	1.4	3.6	1.2	0.0	0.0	381
Female	27.8	2.4	35.5	2.5	21.2	2.2	11.4	1.5	4.2	1.1	0.0	0.0	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	35.2	2.7	33.6	2.5	19.6	2.1	7.1	1.3	4.4	1.1	0.0	0.0	571
Rural	23.4	3.3	34.4	3.4	26.4	3.2	12.8	2.3	2.9	1.1	0.0	0.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	22.9	3.5	28.7	4.8	26.7	4.2	16.8	3.5	5.0	2.1	0.0	0.0	157
Q2	30.7	4.9	34.6	4.7	22.3	4.5	8.2	2.3	4.1	1.8	0.0	0.0	179
Q3	33.3	4.9	27.1	4.2	21.3	4.7	10.8	2.4	7.6	2.6	0.0	0.0	159
Q4	41.4	4.6	32.6	4.3	15.9	3.1	7.7	2.3	2.3	1.1	0.0	0.0	174
Q5(richest)	28.2	3.9	44.8	4.6	23.5	3.9	2.5	1.5	1.1	0.7	0.0	0.0	184
M	0.0	0.0	32.0	27.6	24.9	22.8	43.1	30.8	0.0	0.0	0.0	0.0	2
Age													
18-29	45.1	5.4	32.0	4.3	14.1	3.3	6.0	2.1	2.9	1.3	0.0	0.0	187
30-44	29.2	3.8	35.0	3.7	25.7	4.3	6.1	1.8	3.9	1.6	0.0	0.0	230
45-59	27.1	3.5	41.3	3.6	19.7	3.5	7.9	2.0	4.1	1.9	0.0	0.0	203
60-69	35.0	5.0	24.7	3.9	22.1	3.9	14.9	2.9	3.3	1.6	0.0	0.0	119
70-79	16.4	3.3	33.7	4.8	29.4	4.5	14.1	3.4	6.4	2.9	0.0	0.0	85
80+	18.3	9.4	24.5	9.0	33.6	10.0	18.7	7.3	4.8	3.4	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	31.3	2.1	33.9	2.0	21.9	1.8	9.0	1.1	3.9	0.8			855

7.1 Self-reported utilization of health services and perceived unmet need

	<u>Ambulatory Adult</u>		<u>Ambulatory Child</u>		<u>Inpatient Adult</u>		<u>Inpatient Child</u>		<u>Informal Care</u>		<u>Unmet Care</u>		<u>Non use</u>	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female	44.4	2.8	4.2	1.0	7.2	1.1	1.3	0.6	7.4	1.3	1.0	0.5	34.4	2.6
Male	34.1	3.9	5.2	2.2	6.3	1.3	1.1	0.6	4.6	1.5	1.5	0.8	47.2	4.0
Area of Residence														
Urban	43.8	3.0	4.4	1.2	6.0	1.0	1.6	0.6	6.8	1.3	1.5	0.7	36.0	2.8
Rural	35.4	4.3	4.7	1.7	8.9	1.8	0.5	0.3	6.0	1.6	0.5	0.4	44.1	4.4
Age Group														
18-29y	39.4	4.8	5.6	2.1	4.4	1.8	1.8	1.1	5.3	1.9	0.5	0.5	42.9	5.6
30-44y	29.1	4.1	13.4	3.5	6.4	1.9	4.4	1.6	8.9	2.7	1.6	1.0	36.2	4.3
45-59y	46.3	4.9	1.2	0.8	7.3	1.8	0.0	0.0	2.9	1.5	0.7	0.5	41.7	4.5
60-69y	41.4	4.6	1.1	0.8	7.3	2.4	0.0	0.0	4.9	2.1	1.5	1.1	43.8	5.1
70+y	48.8	5.0	0.7	0.5	8.7	2.4	0.0	0.0	10.6	2.9	1.7	1.1	29.5	4.1
Education Years														
0														
1-5	18.7	7.6	0.0	0.0	10.4	5.5	0.0	0.0	12.4	5.9	0.0	0.0	58.5	9.8
6-11	38.3	3.1	5.0	1.7	7.9	1.4	1.1	0.5	7.2	1.8	0.5	0.3	40.0	3.2
12+	46.4	3.3	4.6	1.1	5.7	1.2	1.6	0.8	5.3	1.2	2.0	0.9	34.5	3.2
Income Quintiles														
Q1	33.2	4.5	2.6	2.0	8.4	2.2	1.6	1.6	6.9	2.1	0.5	0.5	46.8	4.5
Q2	38.0	4.8	7.1	2.6	6.4	1.8	1.2	0.7	4.8	1.5	0.3	0.3	42.2	4.8
Q3	45.4	5.1	3.9	1.7	5.9	1.9	1.5	0.9	7.2	2.8	0.4	0.4	35.6	4.3
Q4	44.7	5.0	2.8	1.3	7.4	2.3	1.1	0.8	8.4	2.8	1.7	1.1	33.8	3.9
Q5	44.4	5.7	5.5	2.2	6.4	1.8	0.8	0.8	5.5	2.0	2.8	1.8	34.5	4.9
Ownership														
Government	50.1	3.0	5.1	1.2	10.6	1.5	2.0	0.7	7.6	1.4	1.2	0.6	23.3	2.4
Private	69.0	6.6	2.0	2.0	4.1	2.8	0.0	0.0	4.9	4.1	2.5	2.3	17.6	4.9
NGO	68.6	19.4	12.4	13.5	0.0	0.0	0.0	0.0	19.0	15.5	0.0	0.0	0.0	0.0
Other	28.5	23.3	0.0	0.0	0.0	0.0	0.0	0.0	71.5	23.3	0.0	0.0	0.0	0.0
TOTAL	41.1	2.4	4.5	1.0	6.9	0.9	1.3	0.4	6.5	1.0	1.2	0.5	38.6	2.4

This table presents the percent of people reporting utilization of health services when they last needed care. It combines information from Q7000, Q7001 and Q7016. Utilization was recorded in the WHS in three separate categories: ambulatory care, hospital inpatient care, and informal care. Time cut-offs for utilization recall used here are 1 year for ambulatory care and informal care, and 3 years for hospital inpatient care. Utilization of ambulatory and inpatient care services

have been further sub-divided into "adult" and "child" categories. In addition to utilization, two other related aspects are being reported here - unmet need and non-use (regardless of expressed need) of health services. Unmet need was estimated on the basis of responses to Q7000 and Q7004. Non-use of health services is simply the percent of respondents who reported no need in Q7000 or did not utilize services in the time cut-offs mentioned above.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, and ownership (of health facility). Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership was constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services.

7.2 Patient assessed characteristics of ambulatory health services

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
Sex																
Female	78.8	4.0	94.6	1.7	91.7	2.0	91.8	2.7	22.8	1.8	20.6	4.8	13.0	2.8	9.6	2.6
Male	77.0	5.5	97.0	1.9	96.1	1.9	95.8	1.8	44.2	19.3	29.3	5.6	11.0	2.3	4.5	2.3
Area of Residence																
Urban	81.1	3.7	94.9	1.7	92.5	2.0	91.0	2.7	32.0	8.0	16.3	4.4	12.7	2.8	9.7	2.7
Rural	71.9	6.5	96.3	2.2	94.2	2.7	97.3	1.8	21.3	3.2	38.5	7.9	12.0	2.5	4.6	2.7
Age Group																
18-29y	65.2	9.3	97.0	2.1	92.0	3.8	85.7	7.3	44.7	27.1	36.1	8.6	6.7	1.6	7.8	4.2
30-44y	66.6	7.0	95.7	2.7	96.8	1.9	93.8	3.1	21.7	2.9	32.2	6.7	12.1	3.8	10.8	4.4
45-59y	83.8	5.0	98.7	0.9	93.1	3.3	97.9	1.2	27.2	3.5	13.6	5.6	14.3	3.3	5.6	2.9
60-69y	86.8	6.0	87.7	6.2	88.8	4.8	96.1	2.2	25.6	4.2	7.4	4.3	20.8	8.2	8.1	4.8
70+y	92.5	4.0	94.2	3.5	92.7	3.9	89.4	5.0	24.6	2.9	25.7	14.4	9.2	2.3	8.9	4.1
Education Years																
0																
1-5	81.2	13.2	95.9	4.2	95.9	4.2	87.6	9.0	28.4	8.2	7.6	7.2	11.8	4.7	0.0	0.0
6-11	78.4	5.5	93.4	2.3	90.5	2.8	89.5	3.8	35.3	12.6	24.2	6.7	10.4	2.4	9.4	3.2
12+	77.5	4.1	96.8	1.5	95.5	1.8	96.3	1.5	24.0	2.4	24.1	4.5	13.9	3.1	6.6	2.3
Income Quintiles																
Q1	81.7	7.9	98.8	1.2	92.2	4.0	91.9	4.4	23.0	3.4	4.5	3.3	11.1	3.7	7.2	4.5
Q2	72.7	7.6	90.5	4.0	94.3	2.9	90.0	6.4	27.8	3.9	10.4	3.9	11.4	2.1	14.9	5.1
Q3	79.0	7.7	94.5	2.8	92.3	4.2	96.5	2.1	24.6	3.2	23.2	7.3	14.7	6.3	7.1	4.0
Q4	79.5	6.2	95.5	2.8	89.5	4.4	93.0	3.0	20.1	2.8	45.1	10.0	13.1	4.6	2.4	1.7
Q5	79.8	6.0	98.9	1.1	96.4	2.3	93.4	3.3	48.2	26.7	26.0	7.2	11.6	2.5	8.1	4.0
Condition																
Child-Visit	100.0	0.0	79.9	19.3	100.0	0.0	100.0	0.0	34.4	16.6	15.6	15.8	4.2	1.3	20.1	19.3
Preventive	100.0	0.0	100.0	0.0	61.8	30.3	100.0	0.0	13.3	6.2	38.2	30.3	17.7	10.7	0.0	0.0
Childbirth	85.8	9.0	100.0	0.0	92.1	4.8	92.3	6.1	18.4	3.6	28.8	10.6	9.7	3.5	6.0	4.3
Acute	88.0	7.1	93.6	6.3	95.3	4.8	100.0	0.0	24.4	8.2	14.4	9.0	8.3	1.9	14.3	9.6
Injury	87.5	8.4	88.7	6.2	80.4	10.9	92.6	5.1	28.2	6.5	9.8	6.8	10.1	2.1	16.4	9.8
Chronic	76.7	4.4	94.9	1.9	93.7	1.9	94.9	1.9	34.6	11.0	22.8	6.0	12.6	3.4	7.0	2.4

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
Other																
Ownership																
Government	83.4	3.5	94.9	1.6	92.5	1.8	91.7	2.5	31.8	7.6	22.7	4.6	10.7	2.4	7.6	2.0
Private	59.1	8.2	96.3	3.0	94.8	2.9	99.2	0.8	20.5	2.6	27.2	8.0	19.0	4.4	11.0	5.3
NGO	100.0	0.0	100.0	0.0	60.6	34.3	0.0	0.0	20.7	9.1	0.0	0.0	11.7	1.2	0.0	0.0
Other	50.0	24.2	100.0	0.0	100.0	0.0	85.7	11.7	3.2	2.7	0.0	0.0	15.3	1.5	0.0	0.0
Self-Assessed Health																
Very-Good	79.3	13.8	100.0	0.0	100.0	0.0	100.0	0.0	32.7	11.4	5.8	6.1	9.1	1.8	0.0	0.0
Good	69.2	6.0	95.3	2.1	95.6	2.0	95.7	2.2	35.9	15.8	39.0	6.2	11.4	2.5	6.2	2.7
Moderate	81.5	4.6	95.0	2.2	92.3	2.6	92.2	3.3	24.8	2.4	18.3	6.0	13.7	3.7	9.9	3.5
Bad	90.9	4.6	96.1	2.9	88.0	5.2	89.5	5.2	25.1	4.2	4.4	3.1	12.2	3.3	7.6	4.5
Very-Bad	84.9	13.6	92.3	7.6	83.9	10.9	75.7	16.6	19.1	9.6	0.0	0.0	8.3	2.6	14.6	10.3
TOTAL	78.3	3.3	95.3	1.4	93.0	1.6	93.0	2.0	28.9	5.9	23.1	3.9	12.5	2.2	8.1	2.0

This table looks at various characteristics of a visit to an ambulatory care facility, for those who utilized such services in the previous 12 months (Q7204 and Q7205). Characteristics of a visit reported here include:

- (1) The percent of patients visiting a medical doctor (of those visiting any ambulatory provider) (Q7302).
- (2) Patients' rating of *skills*, *equipment*, and *drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7304, Q7305 and Q7306, respectively.
- (3) Barriers or facilitators to care access have been reported here for those who utilized ambulatory care services in the last 12 months. They include:
 - (a) *Mean time taken to reach facility*: patient reported times from Q7307 were used to calculate averages across sample sub-groups.
 - (b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7308.
 - (c) *Mean cost of visit*: patient reported costs in Q7309-Q7313 were summed up for an individual and then averaged across sample sub-groups.
 - (d) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination listed in Q7328-Q7334.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include: sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents could select one of the

following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention		Wait Time		Talked Respectfully		Dignity		Privacy		Communication	
	Travel Time											
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex												
Female	25.7	4.8	35.3	4.0	12.1	3.3	17.4	3.8	19.3	3.7	28.9	4.4
Male	29.3	6.6	28.8	5.3	11.1	4.2	10.2	4.2	24.2	5.8	18.1	5.9
Area of Residence												
Urban	28.9	5.0	35.3	4.1	10.6	2.8	15.0	3.7	17.8	3.6	24.3	4.4
Rural	21.7	6.1	28.8	5.7	14.7	7.5	15.8	4.8	27.5	7.1	29.0	7.3
Age Group												
18-29y	24.5	7.5	36.2	8.3	22.8	9.0	21.4	7.9	32.2	8.6	36.8	9.0
30-44y	20.2	6.5	34.7	6.8	11.8	6.0	13.9	4.8	15.4	5.8	25.0	6.9
45-59y	36.7	9.7	29.3	6.2	1.7	1.2	6.9	3.6	10.2	4.4	13.7	4.0
60-69y	23.2	7.8	35.5	7.8	18.3	6.4	28.0	8.6	31.5	8.0	34.8	7.8
70+y	24.5	7.4	32.3	9.1	9.0	4.3	12.3	5.0	21.4	6.5	24.2	7.0
Education Years												
0												
1-5	26.6	11.6	19.0	9.9	0.0	0.0	12.5	8.4	21.6	15.3	12.5	8.3
6-11	27.5	7.0	34.1	5.3	12.0	4.0	14.6	4.4	23.1	5.1	27.0	5.4
12+	25.5	4.6	33.4	4.5	13.0	4.2	16.7	3.9	19.3	4.4	25.9	4.8
Income Quintiles												
Q1	23.4	6.5	37.8	8.0	6.9	4.4	6.1	3.3	19.8	6.3	34.0	8.5
Q2	25.6	7.8	34.8	7.1	15.2	6.6	19.7	7.7	21.6	7.3	27.9	7.7
Q3	36.0	11.0	30.6	7.5	9.0	5.2	13.8	5.3	11.2	4.9	23.2	7.2
Q4	20.9	8.2	27.3	7.2	15.9	8.0	15.5	4.8	24.8	8.4	26.8	8.6
Q5	27.5	6.8	38.6	6.9	9.7	4.2	17.8	5.6	25.6	6.6	19.2	5.8
Condition												
Child-Visit	32.5	21.2	39.0	22.7	32.5	21.2	39.0	22.7	32.5	21.2	19.5	18.9
Preventive	46.8	31.6	15.1	15.8	0.0	0.0	15.1	15.8	15.1	15.8	0.0	0.0
Childbirth	18.3	10.5	44.5	10.8	13.8	9.0	10.9	7.3	35.8	11.5	32.6	10.5
Acute	36.2	16.4	17.4	9.3	7.9	7.6	0.0	0.0	2.3	2.3	19.6	10.3
Injury	31.1	12.5	7.0	5.0	3.6	3.6	13.6	7.0	14.4	6.8	23.6	10.5
Chronic	26.7	5.6	30.6	5.3	8.1	2.9	8.9	2.6	15.9	3.8	17.7	4.2
Other												
Ownership												
Government	29.2	4.7	35.1	4.1	13.4	3.7	16.5	3.6	20.3	4.1	27.7	4.5
Private	18.8	6.2	25.6	6.3	1.7	1.2	11.2	5.1	19.6	6.0	15.2	5.4
NGO	0.0	0.0	39.4	34.3	39.4	34.3	0.0	0.0	0.0	0.0	39.4	34.3
Other	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	11.8	11.8	11.8	11.8
Self-Assessed Health												
Very-Good	52.3	21.2	24.9	15.0	11.7	11.4	19.0	13.7	24.9	15.0	24.9	15.0
Good	19.0	5.6	27.6	5.0	10.0	5.2	12.9	3.6	23.8	5.8	25.7	6.2
Moderate	29.8	6.6	34.6	5.4	12.8	4.0	15.1	4.5	17.0	4.5	27.9	5.6
Bad	25.2	8.0	47.0	7.8	14.1	6.5	23.5	7.6	29.5	9.2	21.8	7.6
Very-Bad	39.3	18.9	40.7	17.3	7.7	7.6	8.3	8.2	7.7	7.6	7.7	7.6
TOTAL	26.7	4.0	33.3	3.3	11.8	3.0	15.2	3.0	20.7	3.4	25.7	3.8

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Autonomy				Confidentiality				Choice of		Basic Amenities			
	Treatment Information		Involvement		Talk Privately		Confidentiality of Records		Healthcare Provider		Cleanliness		Space	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female	38.1	4.3	38.5	4.3	32.8	5.2	30.4	4.8	41.1	4.5	16.2	3.4	25.5	4.1
Male	31.5	6.8	33.8	6.5	20.6	5.3	19.8	6.3	38.2	6.2	12.6	4.2	23.4	5.0
Area of Residence														
Urban	35.0	4.4	34.9	4.4	32.3	5.2	22.9	4.7	34.5	4.6	13.5	3.4	21.5	3.7
Rural	39.0	6.4	42.1	6.2	22.1	4.8	36.8	7.8	53.3	6.5	18.9	4.9	32.7	6.8
Age Group														
18-29y	51.8	9.0	46.9	8.6	30.8	8.4	39.5	9.2	42.8	8.5	34.3	9.0	49.6	8.2
30-44y	41.2	6.8	40.7	6.4	32.0	6.1	36.1	8.3	37.0	7.3	9.7	3.7	21.1	6.7
45-59y	26.9	5.7	23.4	5.4	15.4	4.6	19.4	5.6	35.8	7.4	9.7	3.8	17.5	5.0
60-69y	37.2	7.8	45.7	8.2	29.8	7.5	26.2	8.5	48.1	7.8	16.8	7.0	23.6	7.3
70+y	25.0	7.2	34.9	8.2	45.3	12.0	13.7	5.6	41.5	9.8	6.0	3.6	12.1	5.6
Education Years														
0														
1-5	6.1	6.1	44.3	13.9	8.4	7.6	9.5	8.4	37.5	17.4	17.5	15.2	44.0	18.6
6-11	35.6	5.7	35.5	5.7	33.4	6.7	23.1	5.6	39.5	6.2	14.4	4.6	22.1	5.0
12+	40.4	4.3	39.6	4.3	28.1	4.2	33.7	5.2	41.4	4.5	16.2	3.6	26.6	4.3
Income Quintiles														
Q1	40.5	8.3	45.8	8.9	27.0	8.1	18.4	6.6	54.7	8.8	12.0	6.2	18.4	7.3
Q2	35.3	8.0	35.8	7.8	30.9	7.4	29.0	8.0	44.9	7.7	19.4	7.2	31.0	7.9
Q3	21.9	6.6	33.5	8.4	18.7	6.5	33.0	8.9	35.3	8.0	13.8	6.1	20.3	7.0
Q4	39.5	8.4	32.5	7.6	38.5	10.0	27.8	8.9	28.8	7.8	10.3	3.5	19.6	6.6
Q5	45.7	7.2	42.0	7.0	29.4	6.4	25.9	6.4	43.6	7.2	19.0	5.8	33.2	7.1
Condition														
Child-Visit	39.0	22.7	39.6	22.8	55.2	22.4	74.7	18.0	87.7	13.0	28.6	19.5	28.6	19.5
Preventive	53.2	31.6	15.1	15.8	15.1	15.8	15.1	15.8	15.1	15.8	38.2	30.3	38.2	30.3
Childbirth	56.2	10.8	40.5	10.7	26.9	10.4	27.7	9.3	52.4	10.8	15.4	7.9	23.5	9.2
Acute	17.9	9.8	29.1	11.5	28.2	11.4	14.1	8.4	36.4	12.3	9.1	6.5	26.4	10.9
Injury	44.8	13.0	41.3	13.2	22.3	9.2	25.6	9.8	57.9	14.6	14.2	10.1	19.8	9.1
Chronic	25.9	5.0	27.1	4.7	24.1	6.0	17.6	4.7	28.5	4.8	10.0	3.2	21.0	4.3
Other														
Ownership														
Government	38.8	4.4	39.4	4.4	31.9	4.7	27.7	4.7	41.4	4.6	17.3	3.6	26.1	4.0
Private	24.5	6.2	27.2	6.2	17.5	5.5	27.3	7.0	35.2	6.7	7.0	4.0	19.5	5.5
NGO	39.4	34.3	39.4	34.3	39.4	34.3	39.4	34.3	39.4	34.3	39.4	34.3	39.4	34.3
Other	11.8	11.8	11.8	11.8	0.0	0.0	0.0	0.0	11.8	11.8	0.0	0.0	50.0	0.0
Self-Assessed Health														
Very-Good	32.0	17.2	24.9	15.0	19.0	13.7	19.8	14.3	19.0	13.7	0.0	0.0	5.7	5.9
Good	39.1	5.7	37.9	5.6	26.4	5.0	33.3	6.8	44.5	5.8	20.1	4.3	31.6	5.1
Moderate	36.3	5.6	37.4	5.5	32.0	6.8	23.7	5.5	35.8	5.4	13.6	4.4	20.6	4.7
Bad	31.7	8.7	42.3	9.8	30.7	8.0	28.3	8.3	55.0	9.1	13.6	8.0	28.8	9.4
Very-Bad	23.1	15.3	23.1	15.3	25.6	16.1	18.3	16.6	35.1	18.1	7.7	7.6	27.6	14.7
TOTAL	36.2	3.7	37.1	3.6	29.2	3.9	27.2	4.1	40.3	3.8	15.1	2.8	24.9	3.3

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.4 Patient assessed characteristics of hospital inpatient health services

	Patients staying for more than 6 days on last hospital visit		Patients rating characteristics of health care provider as adequate / more than adequate				Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Patients waiting more than a month to be admitted		Mean number of patients sleeping in the same room		Mean cost for a hospital visit		Percent of population perceiving discrimination			
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	%	SE	No	SE	mean cost	SE	%	SE
Sex																				
Female	70.3	4.0	87.7	2.2	87.5	2.4	75.9	3.2	44.4	5.8	34.2	3.8	1.3	0.8	4.4	0.7	41.2	8.8	14.4	2.4
Male	78.8	4.6	86.0	4.8	86.8	4.6	70.2	6.3	50.1	7.9	26.1	5.3	8.3	4.2	4.2	0.7	31.3	8.5	18.5	5.2
Area of Residence																				
Urban	70.9	4.0	89.3	2.4	88.5	2.5	72.8	3.6	38.7	4.3	27.1	3.9	3.6	1.8	4.3	0.7	31.3	5.7	15.0	2.8
Rural	76.5	5.3	82.5	4.0	84.6	3.8	77.9	5.1	61.9	11.8	43.0	5.7	2.5	1.8	4.4	0.8	52.7	15.6	16.8	4.2
Age Group																				
18-29y	56.5	8.1	80.1	5.7	76.3	6.4	82.9	5.6	26.9	5.3	35.4	7.6	0.0	0.0	3.8	0.8	26.5	10.0	20.9	5.6
30-44y	46.1	7.0	85.0	5.0	84.9	4.9	71.2	6.8	52.6	9.0	46.7	8.3	2.5	1.8	4.0	0.7	32.1	10.0	13.9	4.9
45-59y	87.9	4.2	93.6	3.5	94.2	3.5	75.9	7.4	62.1	12.7	20.3	6.2	5.0	3.4	4.8	0.9	23.7	4.1	11.8	4.2
60-69y	83.9	5.3	86.1	5.4	83.1	5.7	63.5	8.1	36.9	4.8	25.7	6.6	5.8	5.6	4.2	0.8	51.2	19.3	23.7	6.8
70+y	86.8	4.0	89.1	3.9	93.2	3.4	77.1	5.1	45.3	7.7	29.5	6.5	3.2	2.3	4.6	0.8	55.4	19.8	12.0	4.0
Education Years																				
0																				
1-5	95.1	5.0	100.0	0.0	81.9	12.2	74.8	13.9	23.7	8.6	12.5	8.0	0.0	0.0	5.1	0.8	22.1	8.8	17.2	11.6
6-11	75.3	4.3	89.4	2.9	89.7	2.9	73.9	4.3	47.4	6.0	30.4	4.8	6.0	2.6	4.6	0.7	34.7	8.0	14.0	3.3
12+	67.3	4.8	83.5	3.4	85.2	3.2	74.4	3.8	46.9	7.5	35.3	4.5	0.9	0.9	4.0	0.7	42.8	11.2	17.1	3.8
Income Quintiles																				
Q1	80.8	6.0	88.4	4.7	89.4	4.8	74.9	7.2	60.6	13.1	30.8	6.7	3.7	2.6	4.2	0.8	31.8	6.3	18.3	5.6
Q2	80.4	6.3	87.1	4.1	92.7	3.4	75.7	7.3	43.7	10.1	19.7	5.5	1.7	1.7	4.9	0.9	24.1	3.9	6.2	3.0
Q3	66.5	7.4	88.5	4.0	87.6	4.0	71.4	6.1	50.4	8.1	35.1	8.4	5.2	3.7	4.2	0.7	54.4	23.9	16.9	5.3
Q4	73.5	5.9	90.1	3.9	90.4	3.9	77.5	6.0	38.9	5.8	29.0	6.3	3.9	3.8	4.3	0.8	34.2	10.6	16.7	5.8
Q5	59.5	6.7	79.9	7.4	74.8	7.6	70.7	7.3	37.9	7.1	48.8	7.7	1.4	1.4	4.0	0.7	43.8	13.8	20.5	6.4
Condition																				
Child-Visit	45.8	13.6	51.6	14.4	55.4	17.4	57.9	17.4	24.8	5.9	61.2	13.8	0.0	0.0	3.9	1.1	9.3	2.7	9.5	9.3
Fever	49.8	13.4	69.9	11.9	64.6	12.5	61.9	12.6	29.1	4.2	32.8	13.8	0.0	0.0	4.2	0.9	8.7	2.5	22.9	11.4
Childbirth	28.0	12.5	97.9	2.2	95.7	3.1	71.9	11.1	14.9	3.8	35.1	8.6	0.0	0.0	2.8	0.9	18.2	7.6	19.2	9.1
Arthritis	100.0	0.0	100.0	0.0	82.6	18.7	100.0	0.0	17.5	6.6	17.4	18.7	0.0	0.0	6.3	0.7	21.0	14.4	0.0	0.0
Asthma	100.0	0.0	62.2	22.8	62.2	22.8	62.2	22.8	52.9	11.8	29.4	23.7	0.0	0.0	4.4	1.7	24.1	12.1	0.0	0.0
Heart-Disease	92.4	4.2	91.3	4.7	95.1	3.8	62.0	8.5	47.2	12.8	25.4	8.4	9.9	6.4	4.5	0.8	22.1	3.9	18.8	6.5
Injury	84.2	7.3	75.5	7.2	80.7	7.3	73.9	8.5	49.5	15.4	28.5	8.7	0.0	0.0	4.5	0.8	51.4	17.4	14.8	6.8
Minor-Surgery	55.5	6.7	95.0	2.9	92.8	3.0	93.5	3.5	51.6	8.5	40.4	6.7	4.8	2.8	3.8	0.7	56.8	22.6	7.4	3.3
Other	83.6	4.2	86.7	3.4	87.4	3.4	71.6	5.3	51.8	7.7	30.7	5.6	2.1	1.6	4.8	0.8	40.6	8.9	18.2	4.5
Ownership																				
Government	72.1	3.3	87.1	2.1	86.8	2.2	74.1	3.1	44.6	4.5	31.3	3.3	3.4	1.4	4.4	0.6	30.5	4.6	15.1	2.2
Private	77.6	13.5	83.2	15.3	100.0	0.0	87.0	12.6	85.8	29.3	65.9	18.0	0.0	0.0	2.4	0.9	219.9	84.3	27.7	17.4
NGO	100.0	0.0	100.0	0.0	100.0	0.0	72.2	28.8	52.2	9.2	0.0	0.0	0.0	0.0	2.8	0.8	58.7	13.8	0.0	0.0
Other																				
Self-Assessed Health																				
Very-Good	76.8	13.2	71.0	18.7	78.0	18.4	78.0	18.4	42.8	17.2	40.2	17.4	0.0	0.0	3.7	1.2	5.9	1.9	35.6	18.8
Good	38.8	6.9	85.7	5.0	85.3	5.1	76.1	6.7	31.3	5.0	33.6	6.5	0.0	0.0	3.7	0.8	20.4	4.4	8.8	3.6
Moderate	76.5	4.2	88.0	2.8	88.8	2.8	78.1	3.7	51.4	6.4	34.0	4.9	4.6	2.5	4.2	0.7	47.6	12.5	13.6	3.1
Bad	89.3	3.4	88.8	3.8	90.3	3.5	67.4	6.5	49.2	9.6	28.0	5.4	4.5	2.6	4.9	0.8	40.6	11.7	17.9	5.0
Very-Bad	73.0	11.7	89.1	8.2	73.0	12.4	66.5	12.9	35.8	9.1	22.3	10.4	0.0	0.0	5.1	0.9	33.8	5.4	30.1	12.1
TOTAL	72.6	3.2	87.2	2.0	87.3	2.1	74.3	3.0	46.0	4.9	32.0	3.3	3.3	1.4	4.3	0.6	38.1	6.7	15.6	2.3

This table looks at various characteristics of a visit to a hospital inpatient care facility, for those who utilized such services in the past 3 years (Q7200, Q7201 and Q7402). Characteristics of a visit reported here include:

(1) The percent of patients who stayed 6 days or more at an inpatient facility during their last visit (Q7404).

(2) Patients' rating of *skills, equipment, and drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7405, Q7406 and Q7407, respectively.

(3) Barriers or facilitators to care access have been reported here for those who utilized inpatient care services in the last 3 years. They include:

(a) *Mean time taken to reach facility*: patient reported times from Q7408 were used to calculate averages across sample sub-groups.

(b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7409.

(c) *Patients waiting more than a month to be admitted*: percent of patients who reported waiting more than a month to be admitted in Q7410.

(d) *Mean number of patients sleeping in same room*: averages across sample sub-groups have been constructed for numbers reported in Q7417.

(c) *Mean cost of visit*: patient reported costs in Q7411-Q7415 were summed up for an individual and then averaged across sample sub-groups.

(d) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination asked in Q7433-Q7439.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication				Autonomy			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions		Treatment Information		Involvement	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex																
Female	32.6	3.9	26.9	3.9	18.7	3.2	18.2	3.2	26.8	3.6	24.1	3.3	46.0	4.0	44.6	3.8
Male	41.3	6.4	30.9	6.0	25.1	5.4	21.2	5.3	24.2	5.6	20.2	5.1	48.5	6.1	48.9	6.4
Area of Residence																
Urban	35.1	4.2	29.7	3.9	21.9	3.6	20.2	3.3	26.9	3.6	22.0	2.9	44.3	3.9	44.3	4.2
Rural	34.8	5.7	24.2	5.8	17.3	4.4	16.3	4.2	24.1	5.5	25.1	5.0	52.3	6.9	49.2	5.4
Age Group																
18-29y	25.5	6.7	30.7	7.5	20.8	4.9	27.6	5.6	35.0	7.6	34.1	7.7	43.5	7.7	53.6	7.4
30-44y	34.9	6.0	34.6	6.6	26.0	7.6	17.4	5.1	34.3	7.5	23.7	5.0	55.6	7.2	44.2	6.3
45-59y	34.9	9.0	13.8	5.4	14.9	7.1	11.4	4.7	12.6	4.5	10.5	4.4	44.4	8.5	37.2	7.7
60-69y	53.0	7.5	34.9	7.3	30.8	7.5	21.3	6.9	22.7	6.4	23.5	5.8	46.5	7.8	51.0	7.7
70+y	30.5	5.7	27.1	5.8	13.9	4.0	19.5	5.6	25.5	5.8	24.4	4.8	43.1	7.2	45.9	6.9
Education Years																
0																
1-5	30.8	15.4	24.3	10.8	18.8	9.8	29.8	12.9	31.6	12.8	28.9	12.7	51.4	16.9	56.8	16.4
6-11	31.6	4.5	26.6	4.6	19.5	4.3	18.1	4.0	19.2	4.3	21.9	3.7	42.0	4.9	43.8	5.0
12+	38.6	4.5	30.2	4.5	21.9	3.9	19.0	3.8	32.7	4.6	23.8	3.8	51.6	5.1	47.2	4.8
Income Quintiles																
Q1	36.5	8.1	20.2	5.8	23.9	6.9	11.2	4.7	17.2	6.0	18.2	5.3	53.3	7.3	46.8	6.5
Q2	30.9	6.8	28.0	7.4	19.1	7.3	7.9	3.4	22.8	6.2	17.6	4.8	32.3	8.0	32.9	8.3
Q3	38.6	7.1	30.0	6.4	19.5	5.4	24.3	5.8	21.9	5.2	20.6	4.9	44.8	7.0	49.7	7.2
Q4	37.9	6.8	29.2	6.5	18.0	5.5	18.7	6.1	30.2	7.3	18.9	5.0	50.9	7.5	50.1	7.5
Q5	29.1	7.8	32.6	7.2	23.1	6.8	34.3	7.1	39.3	8.4	42.7	6.9	53.3	7.1	49.1	8.0
Condition																
Child-Visit	27.3	12.3	28.1	12.6	33.2	13.5	36.7	13.8	51.8	17.4	32.9	16.6	68.8	16.8	65.5	14.9
Fever	29.2	11.3	32.3	11.8	30.8	12.0	33.7	13.0	51.6	12.9	51.4	12.9	62.6	13.5	54.1	13.9
Childbirth	2.0	2.1	24.1	8.5	11.5	10.6	13.4	10.6	27.3	13.8	25.8	11.9	23.6	12.2	38.4	11.3
Arthritis	17.4	18.7	34.8	28.0	0.0	0.0	17.4	18.7	17.4	18.7	17.4	18.7	65.2	28.0	100.0	0.0
Asthma	67.0	20.6	53.9	22.5	29.4	23.7	13.4	13.0	0.0	0.0	0.0	0.0	29.4	23.7	29.4	23.7
Heart-Disease	28.1	8.5	26.3	8.4	23.5	8.0	23.9	8.8	19.8	7.6	7.4	3.9	56.3	9.8	65.1	8.3
Injury	29.4	7.7	18.0	6.3	24.4	7.8	30.8	8.2	27.9	7.8	44.5	8.6	58.3	8.9	53.4	8.6
Minor-Surgery	48.1	7.5	30.5	6.2	12.6	4.4	9.1	3.9	15.3	5.5	8.1	3.4	40.6	7.0	37.5	6.2
Other	39.5	5.6	28.5	4.9	22.6	5.0	17.4	3.9	30.2	5.7	26.6	4.8	43.9	5.8	38.1	5.8
Ownership																
Government	33.8	3.6	27.8	3.4	20.9	3.0	19.4	2.7	26.2	3.0	23.0	2.6	46.9	3.6	46.5	3.6
Private	56.0	20.6	39.2	18.7	16.8	15.3	13.0	12.6	25.7	17.1	25.7	17.1	53.4	20.4	36.6	18.8
NGO	72.2	28.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other																
Self-Assessed Health																
Very-Good	35.2	19.3	29.0	18.7	29.0	18.7	22.0	18.4	29.0	18.7	22.0	18.4	62.1	17.6	55.1	18.3
Good	16.4	5.2	34.2	7.5	20.1	5.7	26.6	6.8	43.2	7.5	34.5	7.4	51.2	7.6	49.6	7.4
Moderate	40.0	5.0	23.7	4.4	16.7	3.6	16.8	4.0	22.5	4.0	17.1	3.1	43.2	5.5	43.1	5.2
Bad	42.3	6.3	31.2	5.6	27.2	6.2	15.8	4.6	16.5	4.7	22.2	4.8	46.8	6.4	46.0	6.2
Very-Bad	22.5	9.5	26.3	11.0	13.4	9.0	23.9	12.2	41.7	12.9	37.0	12.8	48.8	12.8	47.3	13.1
TOTAL	35.0	3.4	28.0	3.2	20.5	2.9	19.0	2.6	26.1	3.0	23.0	2.5	46.7	3.4	45.8	3.4

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Confidentiality				Choice of health care provider		Basic Amenities				Social Support			
	Talk Privately	Confidentiality of records					Cleanliness	Space			Family Visit	External Contact		
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female	29.1	3.9	21.8	4.5	53.4	4.8	23.7	3.3	40.6	4.2	17.1	3.2	27.4	3.8
Male	28.6	5.9	31.6	7.1	48.1	5.8	25.4	5.7	31.6	5.8	21.8	5.2	24.8	5.2
Area of Residence														
Urban	27.9	4.0	24.1	4.8	54.2	4.7	23.6	3.5	40.1	4.1	20.0	3.3	27.3	4.0
Rural	31.4	6.3	26.3	7.3	46.8	7.1	25.5	5.3	33.5	5.8	14.7	4.7	25.2	5.7
Age Group														
18-29y	41.0	8.1	30.2	8.8	67.4	6.6	32.1	7.4	43.9	7.7	29.2	7.4	35.1	8.5
30-44y	35.7	7.7	35.7	8.1	57.2	7.3	41.7	6.6	40.4	6.2	29.2	7.3	25.8	6.6
45-59y	13.2	4.7	7.7	4.3	41.5	10.2	17.7	5.4	36.2	8.6	8.8	3.8	11.9	4.5
60-69y	32.2	7.2	34.1	10.1	54.9	8.2	16.5	5.0	37.0	7.2	22.7	6.5	32.3	7.7
70+y	25.5	5.5	22.1	7.9	43.5	7.2	13.9	4.8	34.9	6.4	7.5	2.9	29.8	5.9
Education Years														
0														
1-5	19.3	9.9	17.3	13.1	56.6	14.0	24.9	12.2	48.4	17.1	13.9	8.8	55.6	14.6
6-11	27.4	4.5	31.1	5.9	50.1	5.5	16.9	3.9	31.1	4.6	17.1	3.9	25.7	4.0
12+	31.9	4.7	18.7	4.6	53.9	5.8	31.7	4.7	44.5	5.3	20.4	3.6	25.1	4.4
Income Quintiles														
Q1	32.3	7.1	15.5	6.5	52.3	8.1	13.4	5.4	30.6	6.8	15.0	6.1	24.1	5.9
Q2	24.4	6.9	17.1	8.5	49.2	8.8	16.3	6.4	39.4	8.7	6.8	3.4	22.2	6.8
Q3	31.6	6.4	22.6	6.7	57.6	7.3	27.8	6.2	34.0	6.7	28.8	7.8	28.8	6.4
Q4	23.5	5.7	30.4	8.4	56.0	7.9	27.9	6.8	39.6	7.5	22.1	5.2	31.7	6.7
Q5	33.9	7.4	38.2	8.7	41.8	6.3	37.1	8.7	48.1	7.3	18.6	7.0	24.6	7.3
Condition														
Child-Visit	40.6	13.5	45.1	26.1	45.4	17.9	67.5	14.3	67.4	13.0	51.9	12.6	57.9	13.7
Fever	29.7	11.7	26.6	16.5	55.4	12.7	55.0	13.4	53.2	13.4	54.3	12.9	51.8	13.2
Childbirth	34.7	10.8	26.7	13.4	50.2	11.3	13.4	10.6	24.1	8.8	24.6	10.5	13.2	7.2
Arthritis	82.6	18.7	21.1	23.7	34.8	28.0	65.2	28.0	100.0	0.0	17.4	18.7	34.8	28.0
Asthma	53.9	22.5	0.0	0.0	86.6	13.0	53.9	22.5	13.4	13.0	0.0	0.0	13.4	13.0
Heart-Disease	22.3	6.8	28.6	11.2	49.9	10.3	18.2	6.5	34.2	7.8	10.9	4.6	30.2	9.1
Injury	44.0	8.6	43.4	12.0	60.8	10.4	27.0	7.1	36.8	8.0	26.5	9.0	44.1	9.4
Minor-Surgery	20.4	5.9	18.1	7.8	36.5	6.5	17.7	6.4	27.8	6.5	3.9	2.3	9.3	4.2
Other	27.8	5.0	21.8	5.4	57.5	6.0	21.7	4.8	44.8	5.8	19.2	4.0	27.7	4.9
Ownership														
Government	28.9	3.4	25.9	4.2	53.3	4.0	24.7	3.0	39.2	3.5	18.3	2.7	26.7	3.3
Private	36.6	18.8	0.0	0.0	27.7	17.4	16.8	15.3	23.3	16.1	27.7	17.4	36.6	18.8
NGO	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other														
Self-Assessed Health														
Very-Good	51.8	18.8	50.4	21.7	75.4	14.3	44.7	18.9	42.3	19.2	29.0	18.7	51.9	18.2
Good	41.6	7.5	40.7	9.0	56.7	7.7	33.6	8.2	37.6	7.0	21.6	7.3	18.5	5.8
Moderate	22.7	4.2	21.4	5.3	48.3	5.5	23.9	3.7	31.7	4.6	17.3	4.1	23.8	4.8
Bad	29.9	6.0	16.8	5.6	52.4	6.4	17.6	5.4	48.0	6.6	17.3	4.6	31.8	6.0
Very-Bad	13.8	7.9	12.2	8.8	45.0	13.0	10.1	9.4	41.6	12.8	13.1	9.0	34.9	12.4
TOTAL	29.0	3.4	24.8	4.0	51.9	3.9	24.2	2.9	38.1	3.4	18.4	2.7	26.6	3.2

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"

	Prompt Attention		Dignity		Communication		Autonomy		Confidentiality		Choice		Basic Amenities		Social Support	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex																
Female	37.4	2.9	38.6	2.9	54.3	3.0	34.4	3.1	39.6	3.0	31.2	2.9	31.9	3.2	35.7	3.4
Male	30.0	3.4	26.3	3.4	40.6	4.0	21.5	3.2	29.9	3.6	26.8	3.6	24.1	3.4	25.1	3.7
Area of Residence																
Urban	39.8	3.3	37.4	3.3	53.8	3.6	36.2	3.6	38.2	3.4	34.6	3.3	34.6	3.7	35.7	3.8
Rural	25.2	3.7	29.0	3.2	41.7	4.0	17.9	2.6	33.0	3.7	19.8	3.1	18.7	3.2	25.3	3.9
Age Group																
18-29y	45.7	5.2	34.7	5.2	56.7	5.8	27.0	4.5	46.8	5.5	26.7	4.6	36.8	5.5	37.7	5.6
30-44y	37.0	4.0	36.1	4.1	45.1	4.7	30.9	4.2	40.3	4.2	31.1	4.3	31.8	4.6	35.8	4.5
45-59y	27.9	4.0	36.9	4.6	51.2	5.0	28.3	4.5	43.6	4.7	32.2	4.2	27.7	4.6	29.9	4.4
60-69y	31.6	4.3	26.6	4.5	45.8	4.2	29.6	4.9	23.6	4.5	31.3	4.7	22.1	4.4	20.6	4.2
70+y	35.2	4.8	36.8	5.0	50.8	4.9	35.0	5.4	25.3	4.0	26.8	4.1	28.2	5.2	36.1	5.3
Education Years																
0																
1-5	27.9	9.5	29.3	9.5	38.5	10.4	30.1	10.5	24.4	9.2	26.5	8.7	18.9	7.1	24.8	9.5
6-11	34.9	3.6	34.7	3.5	44.3	3.7	33.4	3.6	31.8	3.5	25.0	3.1	31.2	3.9	31.2	3.6
12+	35.6	3.2	35.2	3.1	55.9	3.1	26.8	2.7	42.1	3.4	34.5	3.0	28.4	2.9	33.6	3.6
Income Quintiles																
Q1	22.0	3.4	23.8	4.3	37.6	4.9	19.5	3.4	21.4	3.1	21.7	4.3	15.5	3.3	24.6	4.3
Q2	33.2	4.9	32.2	4.9	51.1	4.8	29.1	4.3	34.5	4.8	22.6	3.9	25.1	4.4	26.8	4.2
Q3	39.8	4.8	40.7	5.2	53.4	5.6	37.5	5.1	43.7	5.1	37.3	5.3	37.2	5.8	35.2	5.4
Q4	39.4	5.0	37.9	5.3	49.7	5.1	34.1	5.2	38.8	5.1	31.9	4.4	34.1	5.5	35.5	5.5
Q5	40.8	4.6	38.2	4.5	57.2	4.4	29.9	4.4	44.2	4.6	35.7	4.9	34.9	4.9	39.4	5.3
Ownership																
Government	36.6	3.0	37.0	3.0	53.8	3.2	33.6	3.1	37.8	3.2	30.7	2.5	31.3	3.2	33.6	3.3
Private	36.5	6.7	30.0	6.7	49.6	7.2	20.4	5.6	40.1	6.6	25.1	6.2	25.4	6.3	32.0	6.5
NGO	100.0	0.0	0.0	0.0	68.6	19.4	49.5	32.4	0.0	0.0	49.5	32.4	12.4	13.5	49.5	32.4
Other	24.0	13.8	35.8	16.7	35.8	16.7	28.5	23.3	40.3	23.9	40.3	23.9	14.3	11.7	24.0	13.8
Self-Assessed Health																
Very-Good	22.0	8.0	31.3	9.1	28.5	8.6	20.7	7.9	44.0	11.2	35.6	10.8	19.4	7.6	21.7	7.8
Good	34.5	3.7	34.4	3.6	53.4	4.2	24.3	3.1	42.0	3.8	27.3	3.3	27.1	3.5	30.0	3.8
Moderate	33.9	3.3	35.6	3.5	49.4	3.6	34.0	3.7	35.9	3.5	28.8	3.3	30.6	4.0	33.9	3.8
Bad	38.6	4.5	32.7	4.6	50.8	4.8	31.2	4.5	26.1	4.7	32.7	4.5	31.3	4.4	32.9	4.8
Very-Bad	52.8	8.6	37.9	8.3	47.4	9.1	36.3	8.2	41.6	10.3	42.0	8.5	33.4	8.2	40.6	8.5
Type of Care																
Ambulatory-Adult	35.7	3.8	36.7	3.9	52.6	4.0	33.8	3.9	41.0	3.9	31.0	3.3	31.8	4.0	33.4	3.8
Ambulatory-Child	33.7	10.6	49.0	11.3	52.7	11.3	39.9	11.3	44.9	9.9	41.9	10.1	28.4	7.7	40.5	10.7
Inpatient-Adult	35.9	6.9	30.0	6.5	47.3	6.0	21.3	6.0	30.4	6.5	34.4	6.4	26.4	5.5	27.7	5.5
Inpatient-Child	49.7	18.8	26.9	12.6	37.5	16.2	32.7	16.0	38.5	16.6	32.7	16.0	24.6	14.5	57.2	19.4
Non-Users	34.4	3.4	32.4	3.3	48.3	3.8	27.5	3.4	32.7	3.2	27.4	3.7	28.3	3.6	30.7	4.0
TOTAL	35.0	2.5	34.6	2.5	49.8	2.7	30.2	2.5	36.5	2.6	29.8	2.4	29.4	2.7	32.3	2.9

This table shows the percent of people who rated a responsiveness domain as "Extremely Important". All survey respondents were queried on their perception of the importance of each of the eight responsiveness domains in section 7100 of the WHS where each domain was described in detail (Q7100-Q7107).

Respondents were asked to say whether the domain was "Extremely Important", "Very Important", "Moderately Important", "Slightly Important" or "Not

Important".

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, ownership (of health facility), self-assessed health, and type of care. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, self-assessed health and type of care sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad". Type of care is a constructed sample sub-group combining information from Q7000, Q7001 and Q7016. There are five categories of type of care: "Ambulatory adult", "Ambulatory child", "Inpatient adult", "Inpatient child" and "Non users".

8.0 Expenditure Background Indicators (2002)

GDP per capita (\$)	3,952
Total health expenditure per capita (\$)	203
Total population (000)	2,329
Life expectancy at birth (Both sexes)	70.3
Life expectancy at birth (male)	64.6
Life expectancy at birth (female)	75.8
Healthy life expectancy at birth (Both sexes)	62.8
Healthy life expectancy at birth (Male)	58.0
Healthy life expectancy at birth (Female)	67.5
Government funded health expenditure as a percentage of total health expenditure	64.1
Social security funded health expenditure as a percentage of total health expenditure	31.8
Private health insurance premiums as a percentage of total health expenditure	0.3
Out-of-pocket payment as a percentage of total health expenditure	35.6

8.1 Distribution of health payments across different population groups

	Mean Household expenditure (EXP) in local currency	% Poor	% Impoverished	% Catastrophic	OOP % EXP	OOP % non-subsistence spending	Mean OOP in local currency
Catastrophic							
No	115	9.5	2.8		6.6	10.5	8
Yes	87	26.4	19.2		49.2	62.3	37
Poor							
No	122		5.8	12.0	10.5	16.0	13
Yes	29		0.0	31.8	29.1	31.7	6
Insurance							
No	111	11.9	5.1	14.3	12.7	17.9	12
Residence							
rural	102	16.4	9.9	21.0	15.4	21.5	13
urban	115	9.9	3.0	11.4	11.4	16.3	11
Income_Quintile							
1st(poorest)	41	59.2	14.1	22.3	20.9	24.9	6
2nd	66	0.0	9.9	17.2	11.2	17.8	7
3rd	92	0.0	0.2	10.1	10.1	17.1	9
4th	124	0.0	1.4	13.8	11.4	17.0	13
5th(richest)	233	0.0	0.0	8.3	9.7	12.7	24
total	111	11.9	5.1	14.3	12.7	17.9	12

OOP: Out of pocket

Household consumption expenditure (EXP) comprises both monetary and in-kind payment on all goods and services and the money value of the consumption of home-made products.

Poor - a household is considered poor when its total household expenditure is smaller than its subsistence spending

Impoverishment - a non-poor household is impoverished by health payment when it becomes poor after paying for health services.

Catastrophic health expenditure occurs when a household's total out-of-pocket health payments equals or exceeds 40% of the households capacity to pay or non-subsistence spending.

Out of pocket health payments (OOP) refer to payments made by households at the point they receive health services. It includes doctor's consultation fees, purchases of medication and hospital bills. Spending on alternative and/or traditional medicine is included. Expenditure on health related transportation and special nutrition is excluded. Insurance reimbursements are also excluded.

Subsistence spending - household subsistence spending is the minimum requirement to maintain a basic life in a society. A poverty line is used in the analysis as subsistence spending.

Food expenditure is the amount spent on all foodstuffs by the household plus the value of the family's own food production consumed within the household. It excludes expenditure on alcoholic beverages, tobacco and food consumption outside the home (e.g hotel and restaurants).

Household capacity to pay - is defined as a household's non-subsistence spending.

8.2 Percent distribution of structure of out-of-pocket health payments

	Inpatient	Outpatient	Traditional	Drugs	Other	Total
Catastrophic						
No	1.7	5.3	3.6	65.7	23.7	100
Yes	9.3	4.0	1.2	51.2	34.4	100
Poor						
No	5.4	4.9	2.4	58.3	29.0	100
Yes	0.1	1.9	5.0	72.2	20.7	100
Insurance						
No	5.1	4.7	2.5	59.2	28.4	100
Residence						
rural	3.5	5.2	6.1	64.1	21.2	100
urban	5.8	4.5	0.8	56.9	32.0	100
Income_Quintile						
1st(poorest)	2.6	3.2	3.9	72.7	17.6	100
2nd	0.9	4.2	0.7	77.1	17.1	100
3rd	0.3	3.5	1.9	80.1	14.3	100
4th	8.4	4.4	0.5	66.3	20.5	100
5th(richest)	6.9	5.9	4.1	38.7	44.3	100
total	5.1	4.7	2.5	59.2	28.4	100

8.3 Financial sources used by households for paying for health services

	Savings	Sold Items	Borrow from relatives &	Borrow from others	Health insurance	Current Income	Other
OOP%non_subsistence_spending							
less than 10%	4.2	0.8	5.4	0.9	2.2	69.0	3.8
10-20%	6.5	1.7	11.0	5.7	9.0	89.1	1.2
20-40%	16.1	0.9	19.5	2.8	9.2	90.1	3.5
above 40%	10.0	2.0	25.3	6.1	6.6	97.5	1.6
Hospitalization							
No	5.4	0.9	10.1	1.9	3.0	77.8	3.1
Yes	18.7	2.4	20.6	7.5	16.8	94.7	2.0
Insurance							
No	7.7	1.2	12.0	2.9	5.4	80.8	2.9
Residence							
rural	6.0	2.4	15.4	2.4	7.3	87.2	2.8
urban	8.5	0.6	10.4	3.1	4.6	77.9	3.0
Income_Quintile							
1st(poorest)	6.5	1.2	13.2	1.4	2.4	75.2	2.6
2nd	6.7	0.9	13.1	4.9	4.7	83.4	0.7
3rd	8.2	0.0	11.7	3.0	7.0	81.9	1.4
4th	9.0	2.2	13.9	3.3	5.1	84.3	6.4
5th(richest)	8.2	1.7	7.9	1.9	7.8	78.9	3.6
total	7.7	1.2	12.0	2.9	5.4	80.8	2.9

10.1 Inability to control the important things in life - q8000

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
Gender													
Male	21.1	3.0	18.5	2.8	39.7	3.8	15.5	2.4	3.8	1.2	1.4	0.8	381
Female	16.1	2.1	17.7	2.1	43.7	2.7	16.0	1.8	4.8	0.9	1.8	0.6	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	19.5	2.8	18.2	2.1	43.5	2.8	13.6	1.8	3.9	0.9	1.4	0.6	571
Rural	15.9	2.3	17.8	3.2	38.8	3.6	20.1	2.8	5.3	1.5	2.1	1.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	13.4	2.8	20.2	4.1	29.6	4.1	29.3	4.1	6.2	2.0	1.3	1.0	157
Q2	19.6	5.4	11.4	3.4	43.8	5.4	19.6	4.2	4.1	1.3	1.5	1.4	179
Q3	26.1	5.0	10.5	2.4	44.8	5.4	9.4	2.0	7.5	2.6	1.7	0.9	159
Q4	17.4	3.7	27.3	3.9	39.4	4.0	12.0	2.9	2.7	1.2	1.2	0.9	174
Q5(richest)	15.4	3.0	20.6	3.6	51.0	5.4	9.0	2.6	1.8	1.0	2.3	1.2	184
M	24.9	22.8	0.0	0.0	0.0	0.0	75.1	22.8	0.0	0.0	0.0	0.0	2
Age													
18-29	24.6	5.4	19.4	3.4	46.3	5.1	6.7	2.6	1.9	1.0	1.1	0.8	187
30-44	16.3	3.0	17.3	3.5	44.9	4.3	15.4	2.9	4.5	1.6	1.6	1.2	230
45-59	19.3	4.4	18.2	3.2	37.5	4.1	18.7	3.0	5.2	1.8	1.1	0.7	203
60-69	16.7	4.2	18.0	4.0	37.7	4.6	20.3	4.4	5.3	2.3	1.9	1.4	119
70-79	15.5	3.5	17.8	3.6	41.5	5.4	16.0	3.5	5.2	1.8	4.0	2.9	85
80+	3.1	3.1	15.0	8.0	40.3	10.2	35.5	10.4	6.1	4.4	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	18.3	2.1	18.0	1.8	41.9	2.2	15.8	1.5	4.3	0.8	1.6	0.5	855

10.2 Inability to cope - q8001

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
Gender													
Male	14.4	2.8	23.4	3.3	38.9	3.3	14.5	2.3	5.2	1.5	3.5	1.6	381
Female	12.7	1.9	21.3	2.1	41.5	2.6	18.8	1.6	3.4	0.9	2.2	0.6	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	14.0	2.4	22.3	2.2	41.7	2.4	15.4	1.4	3.3	0.9	3.4	1.1	571
Rural	12.4	2.4	22.2	3.7	37.8	3.7	19.9	2.6	6.2	1.9	1.6	1.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	10.8	2.5	24.5	3.8	27.8	3.4	29.2	3.9	4.9	2.1	2.8	1.4	157
Q2	14.3	4.3	18.2	4.0	44.3	4.5	15.7	3.4	4.8	2.1	2.7	1.6	179
Q3	17.0	4.4	15.9	3.1	41.0	4.9	14.9	2.6	8.9	2.6	2.4	1.0	159
Q4	18.2	4.1	22.4	4.1	42.2	4.5	13.1	3.0	2.3	1.4	1.8	1.0	174
Q5(richest)	7.2	2.4	30.0	4.0	44.9	4.6	12.7	2.8	0.9	0.7	4.2	2.8	184
M	24.9	22.8	0.0	0.0	43.1	30.8	32.0	27.6	0.0	0.0	0.0	0.0	2
Age													
18-29	16.1	5.4	33.7	4.3	40.2	5.3	8.6	2.7	0.9	0.9	0.4	0.4	187
30-44	10.4	2.4	21.0	3.6	43.8	4.0	16.1	2.5	4.7	1.9	4.0	2.3	230
45-59	15.3	3.6	15.0	2.9	42.8	3.8	17.6	3.0	6.7	2.3	2.5	1.0	203
60-69	13.1	3.9	27.8	4.2	33.5	5.0	20.2	3.7	2.5	1.1	3.0	1.6	119
70-79	16.0	3.9	12.1	3.4	39.8	5.1	20.3	4.1	5.6	2.0	6.1	3.3	85
80+	2.1	2.1	16.5	8.3	27.8	9.0	45.7	10.5	7.9	4.8	0.0	0.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	13.4	1.8	22.3	1.9	40.4	2.0	16.9	1.3	4.2	0.9	2.8	0.8	855

10.3 Satisfaction with your health - q8002

	Very dissatisfied	SE	Dissatisfied	SE	Neither satisfied nor dissatisfied	SE	Satisfied	SE	Very Satisfied	SE	Missing	SE	N
Gender													
Male	6.0	2.0	19.1	3.0	21.9	2.6	43.7	3.8	6.5	1.8	2.8	1.1	381
Female	4.2	0.8	20.3	2.0	31.0	2.2	39.5	2.4	3.2	0.8	1.7	0.7	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	6.0	1.4	18.2	2.0	30.4	2.0	38.8	2.5	4.2	1.0	2.5	0.8	571
Rural	3.0	1.1	22.9	3.4	20.2	2.9	46.6	3.8	5.7	1.7	1.6	1.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	3.9	1.5	26.9	3.8	29.7	3.8	33.3	4.1	4.9	2.2	1.3	1.0	157
Q2	3.2	1.6	23.7	4.2	33.8	4.8	33.9	4.3	3.5	1.3	1.8	1.4	179
Q3	10.6	4.1	17.5	3.4	27.2	3.9	36.2	4.3	4.5	1.5	4.1	1.9	159
Q4	4.0	2.0	17.1	3.4	28.2	4.4	44.7	5.1	3.6	1.6	2.4	1.3	174
Q5(richest)	3.6	1.5	14.5	3.8	16.9	3.0	56.6	4.4	6.9	2.5	1.5	1.0	184
M	24.9	22.8	0.0	0.0	0.0	0.0	75.1	22.8	0.0	0.0	0.0	0.0	2
Age													
18-29	6.9	3.1	9.6	3.0	17.2	3.7	56.2	5.1	9.6	2.9	0.4	0.4	187
30-44	4.3	1.5	17.8	3.9	22.2	3.1	48.7	4.8	3.8	1.4	3.2	1.6	230
45-59	3.2	1.6	20.5	3.1	35.4	4.0	38.2	4.2	1.4	1.1	1.2	0.7	203
60-69	5.7	2.5	27.5	4.8	37.4	4.8	25.5	3.7	1.2	0.9	2.6	1.6	119
70-79	2.5	1.1	32.9	4.6	26.5	4.3	24.8	4.3	8.5	2.8	4.7	3.0	85
80+	13.6	6.7	24.8	8.9	26.9	9.8	25.7	9.7	5.9	4.2	3.1	3.1	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	5.0	1.0	19.8	1.7	27.0	1.7	41.4	2.1	4.7	0.9	2.2	0.6	855

10.4 Feeling safe when alone at home - q8011

	Completely safe	SE	Very safe	SE	Moderately safe	SE	Slightly safe	SE	Not safe at all	SE	Missing	SE	N
Gender													
Male	18.3	3.3	38.8	4.1	22.3	2.7	15.6	2.3	4.4	1.2	0.6	0.4	381
Female	8.7	1.6	28.0	2.6	32.8	2.4	18.5	1.9	11.8	1.8	0.2	0.2	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	13.7	1.9	30.7	2.7	28.7	2.1	18.0	1.7	8.6	1.5	0.3	0.2	571
Rural	11.7	3.4	37.1	4.8	26.9	3.3	15.5	2.6	8.2	2.5	0.5	0.5	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	11.4	2.8	30.8	4.4	22.5	4.1	22.5	3.7	11.8	3.0	0.9	0.9	157
Q2	9.8	3.4	33.2	4.7	25.4	3.3	22.7	4.2	8.9	2.9	0.0	0.0	179
Q3	13.9	4.0	28.1	4.7	31.7	4.2	18.8	3.5	6.5	1.7	1.1	0.8	159
Q4	14.5	3.2	39.0	4.7	28.9	3.3	10.2	2.4	7.4	2.0	0.0	0.0	174
Q5(richest)	15.5	3.9	31.9	5.6	31.7	4.3	12.8	3.0	8.1	2.2	0.0	0.0	184
M	0.0	0.0	75.1	22.8	24.9	22.8	0.0	0.0	0.0	0.0	0.0	0.0	2
Age													
18-29	13.3	3.3	37.7	5.2	30.3	5.0	13.3	2.9	4.9	2.3	0.4	0.4	187
30-44	17.2	3.8	34.6	4.0	27.8	3.7	14.7	3.2	5.8	1.8	0.0	0.0	230
45-59	12.6	2.5	32.2	4.1	23.7	3.4	23.9	3.5	7.6	1.8	0.0	0.0	203
60-69	8.9	2.5	25.8	4.0	36.0	4.1	16.8	3.3	11.3	3.2	1.2	1.2	119
70-79	12.0	2.8	28.0	4.9	25.7	4.4	16.5	3.5	17.8	3.9	0.0	0.0	85
80+	1.4	1.4	34.8	11.0	22.9	7.6	18.8	6.9	19.0	7.9	3.1	3.0	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	13.0	1.7	32.8	2.4	28.1	1.8	17.2	1.4	8.5	1.3	0.4	0.2	855

10.5 Feeling safe when walking down the street alone after dark - q8012

	Completely safe	SE	Very safe	SE	Moderately safe	SE	Slightly safe	SE	Not safe at all	SE	Missing	SE	N
Gender													
Male	16.6	3.3	26.1	3.0	26.0	3.2	20.7	2.6	9.1	2.0	1.5	0.6	381
Female	6.2	1.3	16.4	2.4	19.4	2.0	26.9	2.4	29.7	2.8	1.4	0.5	474
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	7.9	2.1	17.7	2.5	22.3	2.4	27.9	2.4	23.1	2.7	1.2	0.4	571
Rural	16.8	3.4	26.9	3.1	22.5	3.0	16.5	2.5	15.3	2.5	2.0	1.0	284
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	12.0	3.0	24.2	3.7	22.5	3.4	23.8	3.7	16.1	3.2	1.5	1.0	157
Q2	13.0	5.4	14.5	3.2	20.8	3.9	27.2	4.2	23.0	4.3	1.6	0.9	179
Q3	12.5	3.8	14.4	3.4	16.7	4.0	30.2	5.3	23.7	4.2	2.5	1.2	159
Q4	7.2	2.8	21.1	4.2	27.2	3.8	23.6	3.6	19.5	3.4	1.3	0.7	174
Q5(richest)	9.9	3.0	29.3	4.8	23.8	4.2	16.0	3.1	20.4	3.3	0.6	0.4	184
M	0.0	0.0	0.0	0.0	24.9	22.8	75.1	22.8	0.0	0.0	0.0	0.0	2
Age													
18-29	11.5	3.4	24.8	4.3	28.7	5.3	18.1	3.8	16.5	3.4	0.4	0.4	187
30-44	9.1	3.2	29.3	3.8	21.7	3.7	26.2	4.1	13.7	3.0	0.0	0.0	230
45-59	14.7	4.5	16.8	3.2	19.6	3.1	28.8	4.1	20.0	3.6	0.0	0.0	203
60-69	8.0	2.4	16.3	3.9	18.7	3.4	24.2	3.8	30.6	4.3	2.3	1.4	119
70-79	12.1	3.8	10.3	3.3	24.6	4.7	19.1	3.7	28.7	4.8	5.2	2.1	85
80+	1.4	1.4	5.2	3.9	14.3	8.1	27.0	9.1	37.2	9.4	14.9	6.1	31
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	10.8	1.8	20.8	2.0	22.3	1.9	24.1	1.8	20.5	2.0	1.5	0.4	855