

# **REPORT OF IRELAND**

## TABLE OF CONTENTS

1	Introduction
2	Methodology
2.1	Sampling
2.2.1	Geographic Component of the Survey
2.2.2	The map
2.3	Survey Metrics
2.3.1	Sample deviation index
2.3.2	HH Sample deviation index
2.3.3	Response rate
2.3.4	Missing data
2.3.5	Reliability
3	Socio-Demographic Profile
3.1	Household composition by residence and age groups according to income quintile
3.2	Selected age distributions by income quintile
3.3	Characteristics of head of household by income quintile
3.4	Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex
3.5	Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education
4	Risk Factors
4.1	Prevalence of smoking and average daily tobacco consumption
4.2	Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers
4.3	Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

- 4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water
- 4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility
- 4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used
- 5 Coverage
  - 5.01 Tuberculosis
  - 5.02 HIV Prevention and Screening
  - 5.03 Antenatal Care and Mother to Child Transmission of HIV
  - 5.04 Birth Delivery Care
  - 5.05 Child's Health - Common Illness
  - 5.06 Child's Health - Danger Signs
  - 5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card
  - 5.08 Vaccination Status of Children 12-23 Months
  - 5.09 Women's Cancer Screening
  - 5.10 Oral Health and Injuries
  - 5.11 Self Reported Non-Communicable Conditions
  - 5.12 Mental Health Conditions
  - 5.13 Cataracts
- 6 Health State Descriptions
  - 6.01 In general, rate your health today - q2000
  - 6.02 Difficulty with 'work and household activities' in last 30 days - q2001
  - 6.03 Difficulty with 'moving around' in last 30 days - q2010
  - 6.04 Difficulty with 'vigorous activities' in last 30 days - q2011
  - 6.05 Difficulty with 'self-care' in last 30 days - q2020
  - 6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021

- 6.07 Amount of 'bodily aches or pains' in last 30 days - q2030
- 6.08 Amount of 'bodily discomfort' in last 30 days - q2031
- 6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050
- 6.10 Difficulty with 'learning a new task' in last 30 days - q2051
- 6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060
- 6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061
- 6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071
- 6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072
- 6.15 Problem with 'sleeping' in last 30 days - q2080
- 6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081
- 6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090
- 6.18 Problem with 'worry or anxiety' in last 30 days - q2091
- 6.19 Adjusted for vignette responses
- 7 Health System Responsiveness
  - 7.1 Self-reported utilization of health services and perceived unmet need
  - 7.2 Patient assessed characteristics of ambulatory health services
    - 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1
    - 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2
  - 7.4 Patient assessed characteristics of hospital inpatient health services
    - 7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1
    - 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2
  - 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"
- 8 Health Expenditure, Insurance and Human Resources for Health

- 8.0 Expenditure Background Indicators (2002)
- 8.1 Distribution of health payments across different population groups
- 8.2 Percent distribution of structure of out-of-pocket health payments
- 8.3 Financial sources used by households for paying for health services
- 8.4 Insurance Coverage
- 8.5 Human resources for health
- 9.1 Adult Mortality Chart 1
- 9.2 Adult Mortality Chart 2
- 9.3 Adult Mortality Chart 3
- 9.4 Adult Mortality Chart 4
- 10 Stress and Social Capital
- 10.1 Inability to control the important things in life - q8000
- 10.2 Inability to cope - q8001
- 10.3 Satisfaction with your health - q8002
- 10.4 Feeling safe when alone at home - q8011
- 10.5 Feeling safe when walking down the street alone after dark - q8012

# 1. Introduction

The World Health Organization initiated the World Health Survey (WHS) as a *data collection platform* to obtain information on the health of populations and on health systems. The WHS responds to the need of countries for a detailed and sustainable health information system and gathers data through surveys to measure essential population health parameters. The WHS uses standard survey procedures and instruments for general population surveys to gather comparable data across WHO Member States. These methods and instruments have been refined through scientific review of literature, extensive consultations with international experts and large-scale pilot tests. WHS is designed to evolve through its implementation by continuous input from collaborators, including policy makers, survey institutions, scientists and other interested parties. The data is jointly owned by the countries and the WHO, and there is a commitment for a long-term data collection, building local capacity and using this information for health policy.

The objectives of the World Health Survey were to:

- develop a **means** of providing:
  - **valid, relevant, reliable** and **comparable** information on **outcomes, functions** and **inputs** to health systems
  - to **supplement** the information provided by routine **HIS**
- build the **evidence base** necessary for policy-makers:
  - to **routinely monitor** if health systems are achieving the **desired goals**
  - and in some countries, to assess if the **increase in health resources** is having the desired effect

The introductory chapter of each country report will provide details on the current sources of data from routine health information systems and ad hoc surveys. It will describe the health situation in the country and identify gaps. It will also present a brief overview on the health system organization within the country.

## 2. Sampling

The WHO sampling guidelines emphasized scientific principles of survey sampling as explicit standards for quality, included examples of good sampling plans, and identified quality assurance standards for countries to adhere to. WHO and technical advisers provided technical support to countries when needed. The following principles were used for sampling in the WHS:

1. The sample should target the *de facto* population (i.e. all people living in that country including guest workers, immigrants and refugees) and not *de jure* population (i.e. the citizens of that country alone) with complete geographic coverage of the country, if possible, or a representative geographic segment of the country.
2. A *sample frame* with 90% coverage that included all key subgroups of interest. Enumeration or listing of households to update the frame was recommended.
3. The WHS sample identified all adult members of the general population aged 18 years or older as its sample. Households are selected using a random, stratified sampling procedure. One individual per household is then selected through a random selection procedure using the Kish table method.
4. The WHS used a sampling strategy which encompasses a known non-zero selection probability for any individual included in the study. This makes it possible to extrapolate the data to the whole population. Households and individuals are selected with a known probability in order to provide selection probabilities for each household and individual in the data set.
5. The WHS sampling strategy was without replacement and categories of non-respondents were clearly defined.
6. The WHS used GPS devices in certain countries and digitized maps to geocode the data within specified guidelines.

Each national report will present a detailed description of the sampling strategy and the allocation tables

---

## 2.2.1 Geographic Component of the Survey

---

Geographic Information refers to any information associated with a location on the Earth's surface. This location can be associated to a coordinate (latitude/longitude), a place's name, a postal address or any other geographic object.

Household Surveys represent an important source of information. If the geographic location of the surveyed clusters is collected at the same time that the interview is conducted users are then able to integrate the spatial dimension in their analysis

The geographic component of the WHO WHS is stored in two sections of the questionnaire:

- 0100 Sampling Information allowing the link between the location of each cluster and the sampling frame used for the survey. This section has also been used in order to collect the type of setting in which each household was located at the time of the survey. This section of the questionnaire has been used for all the countries part of the WHO WHS
- 0200 Geocoding Information for the collection of the latitude and longitude of each household. This section has only been used in 27 countries where GPS devices were available or have been provided. The choice of these countries has been done on the basis of the availability of digital maps delimiting the surveyed clusters, the field data collection planning and their distribution within the different WHO regions. Specific training and field material have been prepared and used by the interviewers or field supervisors in these countries in order to collect the latitude and longitude of each of the surveyed household.


As the sampling design may not always include the administrative structure of the country and as the codes entered in the data set for the sampling level needed to be labelled, a key correspondence table containing this information has been collected after the survey itself.

Once all the survey data and the key correspondence table available a specific protocol has been applied to the two sections mentioned earlier in order to clean eventual data collection or data entry mistakes as well as to fill the gaps in close collaboration with the survey institution.

The questionnaire as well as the training material and data cleaning protocols used can be downloaded from the WHO WHS web site (<http://www.who.int/healthinfo/survey/en/index.html>) in the "Survey Instrument and Related Documents" section.




2.3.1 Sample deviation index



SDI

2.3.2 HH Sample deviation index



HDI

---

### 2.3.3 Response rate

---

---

#### Household

Selected	1,014
----------	-------

Interviewed	1,014
-------------	-------

Percentages	100
-------------	-----

#### Individual


Selected	1,014
----------	-------

Interviewed	1,014
-------------	-------

Percentages	100
-------------	-----

---

2.3.4 Missing data



---

### **2.3.5 Reliability**

---

**Insufficient / No data available**

### 3.1 Household composition by residence and age groups according to income quintile

	Percent Rural	SE	Mean Number of									
			Household members	SE	Children < 2 years	SE	Children < 5years	SE	Children 5 -17 years	SE	Persons 18+ years	SE
Q1(poorest)	35.8	4.08	1.8	0.08	0.0	0.01	0.1	0.02	0.1	0.03	1.6	0.06
Q2	34.8	3.16	2.7	0.13	0.1	0.02	0.2	0.05	0.5	0.07	2.0	0.07
Q3	41.8	3.19	3.2	0.12	0.0	0.01	0.2	0.03	0.6	0.07	2.4	0.09
Q4	47.8	4.03	3.5	0.11	0.0	0.01	0.2	0.04	0.8	0.07	2.6	0.09
Q5(richest)	39.8	3.92	3.9	0.11	0.0	0.01	0.2	0.03	1.0	0.08	2.7	0.09
All	39.8	1.18	3.0	0.06	0.0	0.01	0.2	0.02	0.6	0.03	2.3	0.04

Note: table based on de jure members of household i.e. usual residents

---

### 3.2 Selected age distributions by income quintile

---

	<b>Females Age 18–49</b>	<b>Persons Age 50+</b>	<b>Persons Age 60+</b>
Q1 (poorest)	62	193	141
Q2	126	138	100
Q3	181	115	47
Q4	178	154	49
Q5 (richest)	204	109	23
All	749	710	361

Note: table based on de jure members of household i.e. usual residents

---

### 3.3 Characteristics of head of household by income quintile

---

	Mean Age of Head	SE	Median Education of Head	Percent Female Head	SE
Q1 (poorest)	57	1.52	3	43.5	3.55
Q2	48	1.50	4	41.4	4.28
Q3	42	1.00	4	37.6	3.30
Q4	45	0.96	4	27.0	3.04
Q5 (richest)	43	0.82	4	33.4	3.57
All	47	0.57	4	36.6	1.87

#### Education

1=no formal education

2=less than primary school

3=primary school completed

4=secondary school completed

5=high school (or equivalent) completed

6=college/pre-university/university completed

7=post graduate degree completed

Note: table based on de jure members of household i.e. usual residents



### 3.4 Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex

	Male	SE	Female	SE	Missing	Total	SE
<b>Age</b>							
0-4	5.7	0.80	5.1	0.50		5.40	0.50
5-14	13.8	1.00	14.9	1.10		14.40	0.70
15-29	29.0	1.50	26.5	1.30		27.70	1.20
30-44	17.8	0.90	21.3	1.00		19.60	0.80
45-59	21.0	1.20	19.1	1.10		20.10	0.90
60-69	6.6	0.70	6.6	0.70		6.60	0.60
70-79	4.2	0.60	4.1	0.60		4.10	0.50
80+	0.9	0.30	1.6	0.40		1.20	0.30
Missing	1.0	0.30	0.8	0.30		0.90	0.20
<b>Residence</b>							
Urban	58.8	1.90	58.9	1.80		58.90	1.60
Rural	41.2	1.90	41.1	1.80		41.10	1.60
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Marital_Status</b>							
NoFormal	6.5	1.00	6.1	0.70		6.30	0.60
LessthanPrimary	10.7	1.10	9.7	1.00		10.20	0.80
Primary	25.7	1.50	23.1	1.50		24.40	1.40
Secondary	40.5	1.90	42.3	1.60		41.40	1.60
HighSchool	3.4	0.70	3.3	0.70		3.40	0.60
College	7.9	0.90	9.5	1.00		8.70	0.70
PostGraduate	2.1	0.40	1.5	0.30		1.80	0.30
Missing	3.2	0.80	4.4	1.00		3.80	0.80
Total	100.0		100.0			100.00	
Number	1,548.0		1,553.0			3,101.00	
Number of Households						1,014.00	

### 3.5 Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education

	Male	SE	Female	SE	Missing	Total	SE
<b>Age</b>							
18-29	29.0	2.30	26.8	2.60		27.90	1.80
30-44	27.9	2.50	27.2	1.90		27.50	1.70
45-59	24.2	2.00	23.5	2.10		23.90	1.30
60-69	10.2	1.60	10.1	1.40		10.10	1.10
70-79	6.3	1.10	7.7	1.20		7.00	0.80
80+	2.4	1.10	4.7	1.70		3.60	1.10
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Residence</b>							
Urban	59.7	2.10	60.9	2.50		60.30	1.40
Rural	40.3	2.10	39.1	2.50		39.70	1.40
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Education</b>							
NoFormal	0.5	0.30	0.2	0.20		0.40	0.20
LessthanPrimary	1.5	0.80	1.5	0.70		1.50	0.60
Primary	31.8	3.00	23.9	2.40		27.80	2.20
Secondary	50.0	3.00	55.0	2.80		52.60	2.30
HighSchool	3.4	0.90	3.3	0.90		3.40	0.70
College	9.0	1.60	13.9	2.10		11.50	1.30
PostGraduate	2.8	0.90	1.7	0.60		2.20	0.60
Missing	0.9	0.50	0.4	0.30		0.70	0.30
Total	100.0		100.0			100.00	
Number	497.3		515.7			1,013.00	

































## 5.09 Women's Cancer Screening

	Total Females	Percent Females 18+	PAP Smear	SE	M	Percent Females 40+	Mammography / Breast Exam	SE	M
<b>Residence</b>									
Urban	314	86.4	24.5	3.5	10.1	43.9	26.5	4.7	13.5
Rural	202	89.5	36.0	3.8	8.2	45.0	18.5	4.4	11.0
Missing	0								
<b>Income_Quintile</b>									
Q1(poorest)	74	56.2	16.6	5.7	12.2	36.1	10.0	4.8	12.5
Q2	97	72.6	21.0	4.3	5.1	38.7	11.3	4.9	11.9
Q3	110	95.5	33.5	4.9	7.1	39.1	24.7	8.5	3.8
Q4	106	100.0	37.5	4.7	4.3	59.2	30.3	6.3	8.9
Q5(richest)	111	100.0	25.7	4.7	13.4	44.8	29.0	7.0	23.0
Missing	17								
<b>Age</b>									
18-29	138	26.8	19.4	3.5	9.7				
30-39	85	16.5	47.9	5.1	7.3				
40-49	105	20.4	33.0	5.5	8.0	20.4	14.0	3.2	13.0
50-59	71	13.8	30.8	6.7	9.0	13.8	37.8	7.4	9.4
60-69	52	10.1	13.7	5.2	15.0	10.1	22.5	6.0	15.8
70+	64	12.4	0.0	0.0		12.4	0.0	0.0	
Missing	0								
Total	516	87.6	29.1	2.7	9.3	44.3	23.3	3.3	12.5

M = % Missing of those with Pap Smear/Mammography SE= Standard Error of % with Pap Smear/Mammography

\*\*\*Note: Values in the age row under the "Females 18 plus" and "Females 40 plus" columns are column percent values.

**Women's Cancer Screening Indicators:**

"Papsmear need" = % of all female respondents aged 18-69

"Papsmear" coverage = % of those female respondents aged 18-69 who self-reported receiving a pap smear test during pelvic examination in the last 3 years.

"Mammography need" = % of all female respondents aged 40-69

"Mammography" coverage = % of those female respondents aged 40-69 who self-reported receiving a mammography test or breast examination in the last 3 years.

\*\*\*\*In several countries women above age 69 were asked the cancer screening questions, and for this reason they are included in the table. For countries that did not ask this questions of women above age 69, these values are expected to be missing.

## 5.10 Oral Health and Injuries

	Oral Health Problems					Road Traffic Accidents				Injury			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
<b>Sex</b>													
Female	516	26.6	88.3	3.6	0.0								
Male	497	23.1	89.7	4.1	0.0								
Missing	0												
<b>Residence</b>													
Urban	611	26.0	87.0	4.1	0.0								
Rural	402	23.2	92.3	3.8	0.0								
Missing	0												
<b>Income_Quintile</b>													
Q1(poorest)	153	14.5	93.3	3.7	0.0								
Q2	179	22.5	82.1	6.8	0.0								
Q3	205	30.2	89.7	4.8	0.0								
Q4	220	25.0	90.6	5.2	0.0								
Q5(richest)	220	29.4	89.6	5.6	0.0								
Missing	37												
<b>Age</b>													
18-29	282	27.8	80.9	6.7	0.0								
30-44	279	27.6	97.3	1.8	0.0								
45-59	242	28.1	93.1	3.8	0.0								
60-69	103	19.0	80.3	10.7	0.0								
70-79	71	10.3	93.0	7.2	0.0								
80+	36	4.2	0.0	0.0	0.0								
Missing	0												
Total	1,013	24.8	89.0	2.9	0.0								

M = % Missing of % Covered      SE= Standard Error of % Covered

### Indicators:

#### Oral Health:

"Need" = % of all respondents indicating any problems in the last 12 months with their mouth and/or teeth.

"Coverage" = % of respondents reporting problems with their mouth and teeth who indicated receiving dental or medical care in last 12 months.

#### Road Traffic Accident:

"Need" = % of all respondents indicating having suffered bodily injury from a road traffic accident in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury from a road traffic accident in last 12 months who indicating having received treatment for their injuries on-site or at health facility within 1 hour of accident .

#### Injuries from Non-traffic Accidents:

"Need" = % of all respondents indicating having suffered bodily injury due to non-traffic accidents in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury due to non-traffic accidents in last 12 months who indicating having received treatment for injuries on-site or at health facility within 1 hour of accident

## 5.11 Self Reported Non-Communicable Conditions

		Angina				Osteo-Arthritis				Asthma				Diabetes			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
Sex																	
Female	516	3.5	77.9	9.8		15.9	54.0	5.5	0.6	9.0	68.7	8.2	4.6				
Male	497	6.1	80.0	9.8		11.2	61.7	8.2	0.8	10.3	48.6	7.8	2.3				
Missing	0																
Residence																	
Urban	611	5.4	72.0	11.4		15.3	55.7	5.8	0.5	10.3	56.5	7.6	2.6				
Rural	402	4.0	93.9	4.5		11.0	60.2	8.4	1.1	8.5	61.2	9.9	4.9				
Missing	0																
Income_Quintile																	
Q1(poorest)	153	14.8	87.1	9.7		27.2	59.9	7.9	1.1	9.7	89.1	6.8	2.6				
Q2	179	6.6	85.9	9.5		16.1	60.3	11.4	1.7	7.7	69.4	11.2	3.6				
Q3	205	3.2	47.6	24.4		12.4	78.3	9.9	0.0	7.2	64.8	14.2	0.0				
Q4	220	1.8	71.0	24.1		12.9	41.9	12.5	0.0	9.8	49.3	13.5	5.4				
Q5(richest)	220	1.7	73.3	20.9		4.8	32.6	15.7	0.0	13.6	41.6	10.8	0.0				
Missing	37																
Age																	
18-29	282	0.0	78.0	13.1		1.5	39.5	23.7	0.0	9.5	35.3	9.8	0.0				
30-44	279	0.5	68.2	19.8		5.2	54.3	14.4	0.0	8.8	50.0	13.2	0.0				
45-59	242	6.6	86.2	9.5		15.2	48.7	7.8	0.0	11.0	64.4	11.5	10.5				
60-69	103	12.8	100.0	0.0		28.3	56.9	8.4	0.0	9.2	100.0	0.0	0.0				
70-79	71	18.1				49.1	68.1	9.0	2.8	7.9	72.9	17.4	8.9				
80+	36	15.0				50.5	59.8	16.4	0.0	11.1	100.0	0.0	0.0				
Missing	0																
Total	1,013	4.8	79.2	8.2		13.6	57.1	4.8	0.7	9.6	58.2	6.1	3.4				

**M = % Missing for Covered**

**SE= Standard Error of %Covered**

**Self Reported Non-Communicable Conditions Indicators:**

"Need" = % of total respondents indicating having received a diagnosis of the condition (angina, arthritis, asthma, diabetes) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (angina, arthritis, asthma, diabetes) indicating they have received treatment or screening for the condition in the last 2 weeks.

## 5.12 Mental Health Conditions

	Total	Depression				Schizophrenia			
		Need	Coverage	SE	M	Need	Coverage	SE	M
Sex									
Female	516	5.9	51.2	7.9	0.0				
Male	497	4.6	61.6	10.9	0.0				
Missing	0								
Residence									
Urban	611	5.8	46.1	7.5	0.0				
Rural	402	4.5	74.6	9.1	0.0				
Missing	0								
Income Quintile									
Q1(poorest)	153	8.5	68.7	12.0	0.0				
Q2	179	7.5	64.1	10.7	0.0				
Q3	205	3.9	57.9	19.3	0.0				
Q4	220	5.2	36.0	12.2	0.0				
Q5(richest)	220	2.8	33.4	14.0	0.0				
Missing	37								
Age									
18-29	282	3.2	44.8	20.3	0.0				
30-44	279	5.2	54.1	13.2	0.0				
45-59	242	7.0	58.0	11.3	0.0				
60-69	103	8.1	84.4	9.3	0.0				
70-79	71	5.8	27.4	25.7	0.0				
80+	36	2.0	0.0	0.0	0.0				
Missing	0								
Total	1,013	5.3	55.7	6.8	0.0				

**M = % Missing for Covered**

**SE= Standard Error of %Covered**

**Mental Health Conditions Indicators:**

"Need" = % of total respondents indicating having received a diagnosis of the condition (depression, schizophrenia) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (depression, schizophrenia) indicating they have received treatment or screening for the condition in the last 2 weeks.



**6.01 In general, rate your health today - q2000**

	Very Good	SE	Good	SE	Moderate	SE	Bad	SE	Very Bad	SE	Missing	SE	N
<b>Gender</b>													
Male	36.7	3.0	22.9	2.7	6.8	1.1	2.8	0.9	0.5	0.3	30.2	4.7	497
Female	34.8	3.1	26.6	2.4	10.5	1.4	3.5	0.9	0.2	0.2	24.4	3.7	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	31.2	3.2	25.5	3.0	8.5	1.2	3.7	1.1	0.4	0.2	30.7	5.2	611
Rural	42.6	4.0	23.7	3.2	9.0	1.7	2.2	0.8	0.3	0.3	22.1	5.2	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	24.8	3.9	22.3	4.0	18.4	3.2	6.2	1.9	1.8	1.1	26.5	5.1	153
Q2	23.5	4.0	25.7	4.3	11.0	2.6	2.7	1.5	0.5	0.5	36.7	6.2	179
Q3	39.0	4.1	23.1	3.4	7.5	2.2	1.9	1.1	0.0	0.0	28.6	4.6	205
Q4	34.2	4.2	31.8	4.7	7.4	1.8	4.0	1.5	0.0	0.0	22.7	4.7	220
Q5(richest)	50.6	4.6	20.8	2.9	2.8	1.1	1.7	1.2	0.1	0.2	24.0	4.6	220
M	44.1	8.7	22.0	6.0	6.6	3.6	3.2	3.2	0.0	0.0	24.0	8.5	37
<b>Age</b>													
18-29	54.4	4.6	14.8	2.8	0.9	0.5	1.8	1.0	0.0	0.0	28.1	4.8	282
30-44	37.1	4.0	24.9	4.4	5.5	1.3	0.8	0.5	0.0	0.0	31.7	5.4	279
45-59	27.0	3.4	33.5	3.4	9.5	2.0	4.5	1.5	1.1	0.6	24.2	4.1	242
60-69	26.3	5.3	28.1	4.8	23.8	4.5	2.0	1.4	0.0	0.0	19.8	4.4	103
70-79	11.4	4.0	25.2	5.8	26.0	6.0	11.4	3.9	1.6	1.6	24.4	6.5	71
80+	13.1	5.6	33.0	13.4	11.1	7.0	9.6	6.7	0.0	0.0	33.2	13.8	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	35.8	2.5	24.8	2.2	8.7	1.0	3.1	0.8	0.4	0.2	27.3	3.8	1,013





## 6.03 Difficulty with 'moving around' in last 30 days - q2010

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	88.5	1.7	5.8	1.2	3.7	0.8	1.1	0.5	0.0	0.0	0.9	0.7	497
Female	84.4	1.9	6.7	1.4	7.2	1.5	1.5	0.5	0.0	0.0	0.2	0.2	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	86.4	2.0	6.2	1.4	5.8	1.2	1.5	0.5	0.0	0.0	0.1	0.1	611
Rural	86.4	2.1	6.3	1.2	5.1	1.5	0.9	0.5	0.0	0.0	1.3	0.9	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	73.3	4.1	10.8	2.6	11.8	2.7	3.7	1.5	0.0	0.0	0.4	0.4	153
Q2	84.1	3.4	7.3	2.0	7.8	2.2	0.5	0.5	0.0	0.0	0.4	0.4	179
Q3	87.2	2.8	5.9	2.1	4.2	1.6	1.5	1.0	0.0	0.0	1.1	1.1	205
Q4	88.4	3.0	6.6	2.4	4.1	1.6	0.3	0.3	0.0	0.0	0.7	0.5	220
Q5(richest)	94.2	2.6	2.5	1.0	2.4	2.2	0.5	0.4	0.0	0.0	0.4	0.4	220
M	88.7	5.5	4.0	3.8	2.7	2.3	4.6	3.5	0.0	0.0	0.0	0.0	37
<b>Age</b>													
18-29	95.9	2.2	0.8	0.4	1.5	1.5	0.6	0.4	0.0	0.0	1.2	1.2	282
30-44	94.0	1.8	3.3	1.3	2.0	0.9	0.5	0.3	0.0	0.0	0.2	0.2	279
45-59	87.3	2.4	5.4	1.5	4.7	1.6	2.1	1.0	0.0	0.0	0.5	0.4	242
60-69	73.2	4.7	11.2	3.2	13.9	3.6	1.6	1.2	0.0	0.0	0.0	0.0	103
70-79	56.2	6.6	25.0	5.6	15.1	4.6	2.8	1.9	0.0	0.0	0.9	0.9	71
80+	44.3	15.3	25.8	13.2	26.7	10.7	3.2	3.2	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	86.4	1.4	6.2	1.0	5.5	1.0	1.3	0.4	0.6	0.4			1,013



## 6.05 Difficulty with 'self-care' in last 30 days - q2020

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	93.4	1.4	3.2	0.8	1.6	0.8	0.4	0.4	0.0	0.0	1.5	0.8	497
Female	93.8	1.2	3.6	0.9	2.0	0.6	0.5	0.3	0.0	0.0	0.1	0.1	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	93.8	1.3	3.9	1.0	1.8	0.7	0.3	0.2	0.0	0.0	0.2	0.2	611
Rural	93.3	1.7	2.6	0.9	1.7	0.9	0.7	0.5	0.0	0.0	1.6	0.9	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	83.7	3.2	9.5	2.4	4.8	1.8	0.7	0.7	0.0	0.0	1.3	1.0	153
Q2	93.5	2.4	4.1	1.6	2.0	1.5	0.0	0.0	0.0	0.0	0.4	0.4	179
Q3	94.1	1.9	3.3	1.4	0.3	0.2	1.2	1.0	0.0	0.0	1.1	1.1	205
Q4	96.4	1.4	0.8	0.6	2.1	1.2	0.0	0.0	0.0	0.0	0.7	0.5	220
Q5(richest)	96.8	1.3	1.2	0.9	0.8	0.4	0.5	0.4	0.0	0.0	0.8	0.6	220
M	96.8	3.2	3.2	3.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	37
<b>Age</b>													
18-29	96.7	1.5	1.6	0.8	0.0	0.0	0.0	0.0	0.0	0.0	1.7	1.3	282
30-44	98.1	1.1	0.9	0.7	0.5	0.5	0.2	0.2	0.0	0.0	0.2	0.2	279
45-59	93.4	1.8	3.4	1.2	1.5	0.8	1.2	0.8	0.0	0.0	0.5	0.4	242
60-69	87.5	3.4	5.8	2.3	5.3	2.3	0.0	0.0	0.0	0.0	1.3	1.3	103
70-79	82.3	4.4	13.3	3.8	2.9	2.1	1.5	1.5	0.0	0.0	0.0	0.0	71
80+	76.3	10.4	9.2	6.7	14.5	8.1	0.0	0.0	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	93.6	1.0	3.4	0.7	1.8	0.5	0.4	0.2	0.8	0.4			1,013



**6.07 Amount of 'bodily aches or pains' in last 30 days - q2030**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	76.6	2.4	10.6	1.8	7.1	1.5	4.0	0.9	0.5	0.5	1.2	0.7	497
Female	71.2	2.3	14.7	1.6	9.9	1.7	3.6	0.9	0.2	0.2	0.4	0.2	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	72.7	2.5	14.8	1.7	7.9	1.5	3.9	0.9	0.5	0.4	0.2	0.2	611
Rural	75.7	2.3	9.5	1.4	9.5	1.9	3.6	1.0	0.1	0.1	1.6	0.9	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	60.5	4.5	15.5	3.1	15.6	3.7	8.5	1.9	0.0	0.0	0.0	0.0	153
Q2	68.2	4.4	17.5	3.2	11.3	2.8	1.4	0.9	0.5	0.5	1.1	0.6	179
Q3	75.8	3.4	13.7	2.8	6.4	1.9	3.0	1.2	0.0	0.0	1.1	1.1	205
Q4	74.7	3.6	10.3	2.6	11.2	2.6	2.6	1.2	0.0	0.0	1.2	0.8	220
Q5(richest)	82.3	3.0	9.7	2.0	2.1	0.8	4.3	1.8	1.2	1.1	0.4	0.4	220
M	90.7	4.4	4.1	3.0	0.6	0.6	4.6	3.5	0.0	0.0	0.0	0.0	37
<b>Age</b>													
18-29	90.3	2.6	4.1	1.7	2.3	1.0	2.0	1.0	0.0	0.0	1.2	1.2	282
30-44	84.7	2.7	8.9	2.3	4.4	1.2	1.7	0.8	0.0	0.0	0.2	0.2	279
45-59	67.8	3.6	15.3	2.3	7.2	1.7	6.6	1.7	1.5	1.0	1.6	0.8	242
60-69	53.5	5.1	21.2	4.3	21.8	4.7	3.5	1.5	0.0	0.0	0.0	0.0	103
70-79	38.2	6.5	30.2	6.5	21.5	5.0	10.1	3.6	0.0	0.0	0.0	0.0	71
80+	30.7	11.8	31.5	10.2	34.6	13.6	3.2	3.2	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	73.9	1.7	12.7	1.2	8.5	1.2	3.8	0.7	0.3	0.2	0.8	0.4	1,013



**6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	86.5	2.2	8.4	1.6	3.4	1.2	0.2	0.2	0.0	0.0	1.5	0.9	497
Female	81.7	1.8	11.7	1.4	5.3	1.1	1.2	0.5	0.0	0.0	0.1	0.1	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	84.4	2.2	10.0	1.4	4.8	1.3	0.9	0.4	0.0	0.0	0.0	0.0	611
Rural	83.5	2.4	10.2	1.9	3.7	1.2	0.5	0.4	0.0	0.0	2.0	1.1	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	72.3	4.0	20.5	3.7	7.1	2.3	0.0	0.0	0.0	0.0	0.0	0.0	153
Q2	82.1	3.5	10.1	2.4	5.9	2.1	1.5	1.2	0.0	0.0	0.4	0.4	179
Q3	89.0	2.4	8.6	2.2	0.8	0.5	0.6	0.4	0.0	0.0	1.1	1.1	205
Q4	84.0	3.3	7.1	2.2	6.8	2.4	0.8	0.7	0.0	0.0	1.2	0.8	220
Q5(richest)	88.5	2.5	7.5	1.7	2.7	1.9	0.3	0.2	0.0	0.0	1.1	0.8	220
M	88.6	5.4	8.2	4.5	0.0	0.0	3.2	3.2	0.0	0.0	0.0	0.0	37
<b>Age</b>													
18-29	92.7	2.6	2.9	1.4	1.5	1.5	1.1	0.8	0.0	0.0	1.7	1.3	282
30-44	90.4	2.3	6.2	1.6	2.5	1.5	0.7	0.5	0.0	0.0	0.2	0.2	279
45-59	81.1	2.8	13.2	2.2	4.0	1.4	0.8	0.4	0.0	0.0	1.0	0.7	242
60-69	73.3	4.8	16.8	4.1	9.5	3.2	0.4	0.4	0.0	0.0	0.0	0.0	103
70-79	67.9	6.2	19.5	5.3	12.7	4.5	0.0	0.0	0.0	0.0	0.0	0.0	71
80+	49.6	11.6	38.3	10.4	12.2	7.3	0.0	0.0	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	84.0	1.6	10.1	1.1	4.3	0.9	0.7	0.3	0.8	0.4			1,013





**6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	91.9	1.7	4.1	1.0	2.1	0.7	0.0	0.0	0.2	0.2	1.6	0.8	497
Female	92.1	1.3	5.4	1.2	1.4	0.6	0.3	0.2	0.0	0.0	0.8	0.5	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	91.7	1.7	5.5	1.2	1.8	0.7	0.1	0.1	0.0	0.0	0.9	0.5	611
Rural	92.6	1.8	3.7	1.2	1.6	0.7	0.2	0.1	0.3	0.3	1.7	1.0	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	86.8	3.2	7.6	2.1	1.8	0.9	0.0	0.0	0.7	0.7	3.1	1.9	153
Q2	92.2	2.4	5.5	1.9	2.0	1.0	0.0	0.0	0.0	0.0	0.4	0.4	179
Q3	92.9	2.0	4.7	1.8	1.1	0.8	0.3	0.3	0.0	0.0	1.1	1.1	205
Q4	93.8	1.9	4.1	1.6	1.2	0.8	0.2	0.2	0.0	0.0	0.7	0.5	220
Q5(richest)	93.0	2.6	3.0	1.2	2.4	1.9	0.1	0.1	0.0	0.0	1.5	0.9	220
M	91.9	4.3	4.9	3.0	3.2	3.2	0.0	0.0	0.0	0.0	0.0	0.0	37
<b>Age</b>													
18-29	93.5	2.4	2.5	1.2	1.9	1.5	0.0	0.0	0.0	0.0	2.1	1.3	282
30-44	95.8	1.7	2.9	1.2	0.8	0.6	0.0	0.0	0.0	0.0	0.5	0.3	279
45-59	90.8	2.0	5.7	1.5	1.8	0.9	0.4	0.3	0.5	0.5	0.9	0.6	242
60-69	87.2	3.2	8.8	2.9	3.5	1.7	0.4	0.4	0.0	0.0	0.0	0.0	103
70-79	84.1	4.7	9.4	4.1	2.4	1.9	0.0	0.0	0.0	0.0	4.1	2.1	71
80+	89.6	6.5	10.4	6.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	92.0	1.2	4.8	0.9	1.7	0.5	0.1	0.1	0.1	0.1	1.2	0.5	1,013



### 6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	87.7	1.9	7.4	1.4	2.2	0.7	0.6	0.4	0.0	0.0	2.1	0.9	497
Female	85.6	2.0	6.9	1.2	4.7	1.0	0.9	0.5	0.2	0.2	1.8	0.8	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	87.4	1.9	7.3	1.4	3.2	0.8	1.0	0.5	0.2	0.2	1.0	0.6	611
Rural	85.5	2.5	7.0	1.6	3.8	1.0	0.4	0.4	0.0	0.0	3.3	1.2	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	77.3	3.7	12.7	3.0	7.2	2.1	0.6	0.4	0.7	0.7	1.5	1.0	153
Q2	87.1	2.9	6.2	2.2	6.6	2.0	0.1	0.1	0.0	0.0	0.0	0.0	179
Q3	84.4	3.2	7.8	2.3	1.2	0.8	2.2	1.3	0.0	0.0	4.4	1.8	205
Q4	91.3	2.4	4.1	1.5	3.0	1.4	0.4	0.4	0.0	0.0	1.2	0.6	220
Q5(richest)	89.7	2.2	6.7	1.9	1.0	0.6	0.1	0.1	0.0	0.0	2.5	1.3	220
M	89.5	5.4	5.7	4.0	2.6	2.7	2.2	2.2	0.0	0.0	0.0	0.0	37
<b>Age</b>													
18-29	93.2	1.9	2.9	1.3	1.1	0.4	0.4	0.4	0.0	0.0	2.5	1.5	282
30-44	92.8	2.0	2.9	1.0	0.9	0.6	0.1	0.1	0.0	0.0	3.2	1.5	279
45-59	85.7	2.9	9.1	2.4	3.0	1.0	1.5	1.0	0.0	0.0	0.6	0.4	242
60-69	70.8	5.2	19.1	4.7	6.7	2.4	1.4	1.4	1.1	1.1	0.9	0.9	103
70-79	65.0	6.8	16.3	5.0	16.7	4.6	0.7	0.7	0.0	0.0	1.4	1.4	71
80+	81.5	9.1	7.4	5.5	9.0	6.4	2.2	2.2	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	86.6	1.5	7.1	1.0	3.4	0.6	0.8	0.3	0.1	0.1	1.9	0.6	1,013



## 6.15 Problem with 'sleeping' in last 30 days - q2080

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	83.1	2.2	9.1	1.5	4.6	1.0	1.6	0.7	0.2	0.2	1.4	0.8	497
Female	74.8	2.7	11.8	1.8	8.5	1.6	3.1	0.8	0.9	0.4	0.9	0.4	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	78.3	2.8	10.3	1.5	7.6	1.5	2.2	0.7	0.7	0.3	0.9	0.5	611
Rural	79.7	2.5	10.7	1.9	5.1	1.2	2.7	0.9	0.4	0.3	1.4	0.9	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	66.8	4.7	15.7	3.6	12.7	3.3	4.0	1.5	0.3	0.3	0.5	0.4	153
Q2	76.2	4.0	12.8	2.8	7.5	2.4	2.2	1.2	1.2	0.7	0.0	0.0	179
Q3	83.2	2.9	6.9	1.8	5.1	1.8	2.4	1.2	0.0	0.0	2.4	1.3	205
Q4	81.9	2.7	6.8	1.7	5.8	1.6	3.1	1.2	0.4	0.4	1.9	1.0	220
Q5(richest)	81.2	3.5	11.7	2.8	4.8	1.9	1.1	0.6	0.6	0.5	0.6	0.4	220
M	85.1	6.7	11.7	6.2	0.0	0.0	0.0	0.0	3.2	3.2	0.0	0.0	37
<b>Age</b>													
18-29	89.2	2.7	5.3	1.6	3.1	1.7	0.3	0.3	0.4	0.4	1.7	1.2	282
30-44	85.0	2.6	8.6	2.0	3.4	1.0	1.9	0.8	0.0	0.0	1.0	0.7	279
45-59	77.0	3.4	7.5	1.8	7.8	1.9	4.5	1.4	1.7	0.7	1.3	0.8	242
60-69	61.4	5.2	22.5	4.1	12.0	3.6	3.6	1.7	0.0	0.0	0.4	0.4	103
70-79	54.3	5.8	23.2	5.4	17.9	4.6	3.7	2.2	0.9	0.9	0.0	0.0	71
80+	60.5	13.1	25.5	11.0	12.0	7.1	2.0	2.1	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	78.9	1.9	10.5	1.2	6.6	1.0	2.4	0.5	0.6	0.2	1.1	0.5	1,013



## 6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	84.0	2.0	10.3	1.5	3.0	1.0	1.1	0.4	0.2	0.2	1.4	0.8	497
Female	77.6	2.5	11.8	1.6	8.0	1.8	1.4	0.5	0.2	0.2	1.0	0.4	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	81.9	2.7	9.7	1.5	6.0	1.8	1.3	0.4	0.1	0.1	1.1	0.5	611
Rural	79.0	2.5	13.1	2.0	4.9	1.2	1.2	0.5	0.4	0.3	1.4	0.9	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	74.6	4.0	14.4	3.3	8.3	2.3	1.2	0.8	0.7	0.7	0.8	0.6	153
Q2	76.9	3.9	13.9	3.2	6.1	2.0	1.9	0.9	0.5	0.5	0.7	0.7	179
Q3	84.3	2.8	6.3	1.7	6.6	2.2	0.5	0.5	0.0	0.0	2.4	1.3	205
Q4	77.5	3.1	15.5	2.5	4.7	1.6	0.9	0.6	0.2	0.2	1.2	0.6	220
Q5(richest)	87.5	3.9	6.7	1.8	3.5	3.0	1.4	0.7	0.0	0.0	1.0	0.5	220
M	83.3	6.7	10.1	4.7	2.5	2.3	4.1	3.2	0.0	0.0	0.0	0.0	37
<b>Age</b>													
18-29	87.8	3.3	5.7	1.6	4.0	2.6	0.8	0.6	0.0	0.0	1.7	1.2	282
30-44	87.1	2.3	8.1	1.7	2.9	1.0	0.2	0.2	0.0	0.0	1.7	0.9	279
45-59	74.7	3.5	15.9	2.8	5.4	1.7	2.4	0.9	0.8	0.6	0.8	0.5	242
60-69	73.1	5.1	13.3	3.3	9.3	3.6	3.0	1.6	0.4	0.4	0.8	0.9	103
70-79	66.0	6.8	12.8	3.9	19.8	5.1	1.4	1.4	0.0	0.0	0.0	0.0	71
80+	66.9	12.7	33.1	12.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	80.7	1.9	11.1	1.2	5.5	1.2	1.3	0.3	0.2	0.1	1.2	0.4	1,013





## 7.1 Self-reported utilization of health services and perceived unmet need

	<u>Ambulatory Adult</u>		<u>Ambulatory Child</u>		<u>Inpatient Adult</u>		<u>Inpatient Child</u>		<u>Informal Care</u>		<u>Unmet Care</u>		<u>Non use</u>	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>														
Female	48.2	2.9	12.2	1.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	39.5	3.1
Male	44.9	3.4	6.2	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	48.9	3.5
<b>Area of Residence</b>														
Urban	48.8	3.3	8.2	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	43.0	3.4
Rural	43.6	4.2	11.3	2.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	45.1	4.4
<b>Age Group</b>														
18-29y	43.6	3.8	7.1	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	49.4	4.0
30-44y	37.1	4.1	24.1	3.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	38.8	4.3
45-59y	51.1	3.7	3.0	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	45.9	3.6
60-69y	57.5	5.4	1.2	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	41.3	5.5
70+y	62.6	6.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	37.4	6.0
<b>Education Years</b>														
0														
1-5	54.3	9.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	45.7	9.2
6-11	62.1	4.8	4.2	1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	33.8	4.8
12+	42.8	2.9	13.0	1.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	44.2	3.2
<b>Income Quintiles</b>														
Q1	54.2	4.7	5.7	1.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	40.1	4.6
Q2	54.7	4.7	6.8	1.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	38.4	4.5
Q3	43.2	4.3	12.5	2.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	44.3	4.7
Q4	41.3	3.6	10.6	2.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	48.1	3.9
Q5	46.5	4.7	11.0	2.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	42.6	4.4
<b>Ownership</b>														
Government	65.1	3.6	18.7	3.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	16.1	2.8
Private	66.2	4.3	5.5	1.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	28.3	4.5
NGO	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other	49.4	35.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	50.6	35.4
TOTAL	46.7	2.6	9.5	1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	43.9	2.7

This table presents the percent of people reporting utilization of health services when they last needed care. It combines information from Q7000, Q7001 and Q7016. Utilization was recorded in the WHS in three separate categories: ambulatory care, hospital inpatient care, and informal care. Time cut-offs for utilization recall used here are 1 year for ambulatory care and informal care, and 3 years for hospital inpatient care. Utilization of ambulatory and inpatient care services

have been further sub-divided into "adult" and "child" categories. In addition to utilization, two other related aspects are being reported here - unmet need and non-use (regardless of expressed need) of health services. Unmet need was estimated on the basis of responses to Q7000 and Q7004. Non-use of health services is simply the percent of respondents who reported no need in Q7000 or did not utilize services in the time cut-offs mentioned above.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, and ownership (of health facility). Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership was constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services.

## 7.2 Patient assessed characteristics of ambulatory health services

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	Skills		Equipment		Drug supplies		Minutes	SE	%	SE	mean cost	SE	%	SE
Sex																
1-5	100.0	0.0							11.8	4.5						
12+	86.7	3.4							15.5	1.7						
Area of Residence																
18-29y	84.0	5.7							15.7	2.4						
30-44y	87.6	5.5							14.7	1.8						
Age Group																
45-59y	93.7	3.0							14.4	1.8						
6-11	93.9	3.0							15.2	1.8						
60-69y	87.4	8.4							19.0	4.0						
70+y	93.2	6.8							36.7	20.1						
0																
Education Years																
Female	89.3	3.1							15.1	1.8						
Male	87.6	4.6							18.2	3.0						
Q1	94.9	4.5							15.2	1.4						
Q2	86.5	7.2							14.8	2.1						
Income Quintiles																
Q3	96.7	2.6							15.6	2.5						
Q4	82.7	6.4							20.8	6.2						
Q5	84.7	4.7							15.1	2.2						
Rural	89.5	4.9							19.6	3.7						
Urban	88.0	3.2							14.4	1.2						
Condition																
Child-Visit	64.6	14.8							13.6	1.6						
Preventive	18.8	8.8							17.4	3.6						
Childbirth	84.4	14.3							19.8	6.7						
Acute	100.0	0.0							17.0	6.6						
Injury	100.0	0.0							15.5	0.5						
Chronic	100.0	0.0							14.4	1.2						

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	Skills		Equipment		Drug supplies		Minutes	SE	%	SE	mean cost	SE	%	SE
Other	96.6	1.7							16.6	2.7						
<b>Ownership</b>																
Government	83.2	5.6							15.8	1.8						
Private	88.7	3.4							16.6	2.6						
NGO																
Other																
<b>Self-Assessed Health</b>																
Very-Good	79.4	6.1							17.2	2.4						
Good	88.7	4.6							20.5	5.3						
Moderate	93.2	6.9							13.5	2.8						
Bad	100.0	0.0							16.4	5.1						
Very-Bad	100.0	0.0							10.5	0.5						
TOTAL	88.5	2.7							16.5	1.9						

This table looks at various characteristics of a visit to an ambulatory care facility, for those who utilized such services in the previous 12 months (Q7204 and Q7205). Characteristics of a visit reported here include:

- (1) The percent of patients visiting a medical doctor (of those visiting any ambulatory provider) (Q7302).
- (2) Patients' rating of *skills*, *equipment*, and *drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7304, Q7305 and Q7306, respectively.
- (3) Barriers or facilitators to care access have been reported here for those who utilized ambulatory care services in the last 12 months. They include:
  - (a) *Mean time taken to reach facility*: patient reported times from Q7307 were used to calculate averages across sample sub-groups.
  - (b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7308.
  - (c) *Mean cost of visit*: patient reported costs in Q7309-Q7313 were summed up for an individual and then averaged across sample sub-groups.
  - (d) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination listed in Q7328-Q7334.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include: sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents could select one of the

following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

### 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention		Dignity		Privacy		Communication	
	Travel Time	Wait Time	Talked Respectfully				Clear Explanation	Time for Questions
	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>								
Female		32.2	5.4	15.2	4.9		18.6	4.9
Male		21.6	5.8	13.4	5.2		14.7	5.2
<b>Area of Residence</b>								
Urban		30.9	6.3	18.9	5.8		21.3	5.7
Rural		21.0	6.4	6.5	3.1		9.0	3.6
<b>Age Group</b>								
18-29y		21.7	5.0	9.1	3.8		9.6	3.5
30-44y		17.9	6.1	6.9	4.2		9.6	4.8
45-59y		39.4	8.3	23.0	8.2		25.3	8.1
60-69y		37.4	12.2	28.1	11.5		33.9	11.8
70+y		24.7	11.0	15.7	8.4		24.7	11.0
<b>Education Years</b>								
0								
1-5		0.0	0.0	0.0	0.0		0.0	0.0
6-11		18.3	5.8	6.9	4.0		13.6	5.0
12+		32.8	5.6	18.4	5.3		20.1	5.2
<b>Income Quintiles</b>								
Q1		17.5	6.2	10.8	5.7		16.5	6.7
Q2		26.4	8.8	8.6	4.2		14.1	5.4
Q3		30.4	8.2	13.8	6.5		13.8	6.5
Q4		22.9	6.6	12.1	5.4		12.1	5.4
Q5		30.6	7.2	20.9	6.6		23.5	6.1
<b>Condition</b>								
Child-Visit		13.3	8.6	8.0	7.8		8.0	7.8
Preventive		27.4	14.9	10.7	7.4		10.7	7.4
Childbirth		51.5	18.5	14.0	13.1		60.6	17.2
Acute		26.9	13.6	19.4	12.6		19.4	12.6
Injury		0.0	0.0	0.0	0.0		0.0	0.0
Chronic		27.0	10.6	5.9	6.7		27.0	10.6
Other		28.7	6.0	16.4	5.3		16.6	5.3
<b>Ownership</b>								
Government		24.4	6.7	12.7	4.8		14.7	4.6
Private		18.7	4.4	2.8	1.5		6.0	1.8
NGO								
Other								
<b>Self-Assessed Health</b>								
Very-Good		30.4	6.5	11.1	5.0		13.0	5.2
Good		41.7	8.0	23.6	7.6		26.9	7.1
Moderate		44.2	13.7	28.7	13.8		34.4	14.1
Bad		49.2	25.1	27.7	23.1		73.0	22.8
Very-Bad		100.0	0.0	0.0	0.0		0.0	0.0
TOTAL		27.3	4.6	14.4	3.9		16.8	3.9

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

### 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Autonomy		Confidentiality				Choice of		Basic Amenities					
	Treatment Information		Involvement		Talk Privately		Confidentiality of Records		Healthcare Provider		Cleanliness		Space	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female			18.8	5.6	18.6	5.0			20.2	5.2	16.2	4.9		
Male			16.8	5.8	16.1	5.6			18.7	5.9	12.1	4.9		
Area of Residence														
Urban			21.2	6.4	21.0	5.9			23.2	6.1	19.2	5.8		
Rural			12.2	4.7	11.3	4.4			13.1	4.4	5.8	2.9		
Age Group														
18-29y			14.8	5.8	13.9	5.1			16.5	6.5	5.9	2.9		
30-44y			9.5	4.8	11.0	5.2			12.3	5.0	9.7	4.6		
45-59y			23.1	8.2	24.8	8.3			26.0	8.1	24.2	8.3		
60-69y			33.9	11.8	26.9	11.5			33.9	11.8	26.9	11.5		
70+y			24.7	11.0	15.7	8.4			15.7	8.4	15.7	8.4		
Education Years														
0														
1-5			8.5	9.1	0.0	0.0			0.0	0.0	0.0	0.0		
6-11			13.6	5.6	6.5	3.9			15.1	5.0	7.3	3.8		
12+			19.3	5.6	21.2	5.4			21.6	5.5	18.1	5.3		
Income Quintiles														
Q1			14.8	6.9	11.6	5.5			18.2	6.5	11.6	5.4		
Q2			14.1	5.4	8.6	4.2			14.3	5.3	11.4	4.8		
Q3			21.3	8.2	21.3	8.1			24.8	8.5	14.8	6.5		
Q4			8.3	4.1	12.1	5.4			13.3	5.3	9.8	4.3		
Q5			23.6	7.6	25.8	6.9			21.5	6.4	19.5	6.4		
Condition														
Child-Visit			5.3	5.4	13.3	8.6			10.4	7.9	0.0	0.0		
Preventive			4.3	4.3	4.3	4.3			4.3	4.3	6.6	4.7		
Childbirth			48.6	18.6	20.7	14.2			48.4	18.5	14.0	13.1		
Acute			7.6	7.5	19.4	12.6			15.2	10.2	7.6	7.5		
Injury			0.0	0.0	0.0	0.0			0.0	0.0	0.0	0.0		
Chronic			5.9	6.7	27.0	10.6			27.0	10.6	5.9	6.7		
Other			21.6	5.7	20.3	5.6			22.9	5.6	16.2	5.2		
Ownership														
Government			20.4	5.9	18.3	5.4			27.3	6.6	13.1	4.3		
Private			5.2	2.6	5.8	2.4			5.4	2.6	2.1	1.0		
NGO														
Other														
Self-Assessed Health														
Very-Good			17.3	6.6	18.2	6.3			20.6	6.5	13.4	5.4		
Good			27.3	8.5	28.3	7.8			28.7	7.1	20.7	6.8		
Moderate			28.7	13.8	28.7	13.8			34.4	14.1	28.7	13.8		
Bad			100.0	0.0	49.2	25.1			78.4	19.5	49.2	25.1		
Very-Bad			0.0	0.0	0.0	0.0			100.0	0.0	0.0	0.0		
TOTAL			17.9	4.4	17.4	4.2			19.5	4.3	14.3	3.9		



The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.4 Patient assessed characteristics of hospital inpatient health services

	Patients staying for more than 6 days on last hospital visit		Patients rating characteristics of health care provider as adequate / more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Patients waiting more than a month to be admitted		Mean number of patients sleeping in the same room		Mean cost for a hospital visit		Percent of population perceiving discrimination	
	%	SE	Skills		Equipment		Drug supplies		Minutes	SE	%	SE	%	SE	No	SE	mean cost	SE	%	SE
Sex																				
Female	32.5	5.0							45.7	6.9			15.6	3.6	4.1	0.8				
Male	50.9	6.0							46.7	5.6			15.0	4.9	4.5	0.9				
Area of Residence																				
Urban	38.5	4.9							36.5	6.0			14.8	4.1	4.5	0.8				
Rural	40.3	6.3							56.3	7.9			16.0	4.7	4.0	0.9				
Age Group																				
18-29y	35.9	8.8							43.3	8.1			11.3	6.8	4.6	1.1				
30-44y	17.4	4.6							39.3	4.0			9.1	3.8	3.9	0.9				
45-59y	51.2	8.4							50.3	10.7			22.3	6.9	4.5	0.9				
60-69y	68.5	10.1							61.9	18.5			16.8	8.4	3.6	1.0				
70+y	57.9	11.6							44.3	8.9			24.9	8.9	5.5	1.1				
Education Years																				
0																				
1-5	48.2	8.0							47.5	8.2			38.9	10.6	4.4	1.2				
6-11	61.9	9.4							36.2	8.0			24.2	7.5	5.1	1.0				
12+	30.8	4.6							41.6	3.6			12.1	3.1	4.1	0.8				
Income Quintiles																				
Q1	56.1	8.9							37.9	5.7			14.4	6.0	4.9	0.8				
Q2	52.5	9.3							33.0	4.5			9.2	5.2	5.4	1.2				
Q3	42.1	8.6							45.9	9.6			18.9	8.2	5.1	0.9				
Q4	22.5	7.5							61.2	15.6			2.1	2.2	3.0	0.9				
Q5	30.7	9.3							45.4	6.2			25.9	6.8	3.5	1.0				
Condition																				
Child-Visit	16.0	7.8							39.2	6.0			13.5	7.4	3.9	1.0				
Fever	17.8	9.2							30.9	7.5			3.2	3.3	3.0	1.0				
Childbirth	1.9	2.0							29.1	5.2			0.0	0.0	3.8	0.8				
Arthritis	90.8	9.0							53.8	16.9			65.7	15.5	3.8	1.3				
Asthma	27.8	11.4							34.4	5.8			0.0	0.0	4.4	1.1				
Heart-Disease	89.5	10.0							71.6	15.9			20.1	17.0	5.1	1.2				
Injury	63.8	14.0							46.5	13.1			6.4	6.4	5.9	1.5				
Minor-Surgery	25.2	8.7							64.3	13.7			30.9	9.6	3.7	1.1				
Other	49.0	6.3							42.3	7.7			12.1	3.9	4.4	0.9				
Ownership																				
Government	40.3	4.2							45.7	5.4			14.1	3.3	4.5	0.8				
Private	25.0	10.7							44.8	8.7			22.6	9.5	2.4	1.2				
NGO																				
Other																				
Self-Assessed Health																				
Very-Good	24.9	6.1							56.2	13.2			7.8	4.2	3.4	0.9				
Good	36.6	5.7							45.1	5.2			25.7	7.2	4.0	0.9				
Moderate	56.0	9.8							50.0	7.2			23.8	8.3	4.2	0.9				
Bad	56.0	13.8							40.0	10.6			10.2	9.5	7.0	1.7				
Very-Bad	24.7	26.3							50.6	11.0			24.7	26.3	5.5	0.5				
TOTAL	39.4	4.0							46.1	5.1			15.4	3.1	4.3	0.8				

This table looks at various characteristics of a visit to a hospital inpatient care facility, for those who utilized such services in the past 3 years (Q7200, Q7201 and Q7402). Characteristics of a visit reported here include:

(1) The percent of patients who stayed 6 days or more at an inpatient facility during their last visit (Q7404).

(2) Patients' rating of *skills, equipment, and drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7405, Q7406 and Q7407, respectively.

(3) Barriers or facilitators to care access have been reported here for those who utilized inpatient care services in the last 3 years. They include:

(a) *Mean time taken to reach facility*: patient reported times from Q7408 were used to calculate averages across sample sub-groups.

(b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7409.

(c) *Patients waiting more than a month to be admitted*: percent of patients who reported waiting more than a month to be admitted in Q7410.

(d) *Mean number of patients sleeping in same room*: averages across sample sub-groups have been constructed for numbers reported in Q7417.

(e) *Mean cost of visit*: patient reported costs in Q7411-Q7415 were summed up for an individual and then averaged across sample sub-groups.

(f) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination asked in Q7433-Q7439.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

### 7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication				Autonomy			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions		Treatment Information		Involvement	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex																
Female			22.3	4.1	12.9	3.6			9.7	3.2					18.8	3.9
Male			23.5	5.8	9.2	4.0			14.6	4.7					18.9	5.7
Area of Residence																
Urban			21.8	4.6	11.4	3.2			8.6	2.8					16.2	3.3
Rural			23.8	5.2	11.6	4.6			14.7	4.9					21.7	6.8
Age Group																
18-29y			25.6	9.1	10.0	7.2			10.0	7.2					21.8	10.9
30-44y			21.9	5.4	17.5	6.0			17.3	5.9					22.2	5.0
45-59y			21.8	6.3	8.3	3.8			8.1	3.8					13.5	4.5
60-69y			20.2	8.5	7.3	5.2			7.0	5.0					14.6	7.2
70+y			26.9	9.1	5.6	4.6			7.8	5.0					21.9	8.4
Education Years																
0																
1-5			0.0	0.0	0.0	0.0			0.0	0.0					0.0	0.0
6-11			26.2	7.1	11.6	4.8			15.9	5.8					18.9	5.9
12+			21.8	4.8	12.8	3.9			13.5	4.2					22.0	5.8
Income Quintiles																
Q1			24.6	6.9	10.4	4.4			12.2	5.4					21.7	7.4
Q2			20.2	7.1	4.1	3.1			6.8	4.1					23.8	7.4
Q3			20.2	8.2	18.0	7.8			23.0	7.9					19.5	7.6
Q4			16.7	6.1	9.3	5.1			10.1	4.6					19.4	8.3
Q5			29.9	7.5	11.3	4.8			6.3	3.8					14.1	5.6
Condition																
Child-Visit			30.6	10.4	22.0	10.4			19.4	9.4					33.9	10.0
Fever			11.1	8.0	0.0	0.0			3.3	3.2					6.6	6.5
Childbirth			6.4	4.9	16.3	8.5			16.3	8.5					23.7	8.5
Arthritis			36.3	13.2	3.1	3.0			20.2	15.4					9.4	7.0
Asthma			31.3	13.7	27.3	12.8			13.7	9.8					24.7	14.0
Heart-Disease			0.0	0.0	8.1	8.0			0.0	0.0					18.6	12.3
Injury			26.9	11.9	15.8	11.6			3.7	3.7					6.7	5.0
Minor-Surgery			27.9	9.1	20.8	8.2			24.4	8.5					26.5	9.2
Other			26.8	5.4	6.7	3.1			9.0	3.7					20.3	5.0
Ownership																
Government			24.7	3.9	12.1	3.1			12.2	3.2					20.2	4.3
Private			0.0	0.0	0.0	0.0			0.0	0.0					4.4	4.2
NGO																
Other																
Self-Assessed Health																
Very-Good			20.1	6.6	9.7	5.0			11.3	5.4					16.8	5.9
Good			22.4	6.5	13.3	6.0			11.4	5.5					21.8	4.6
Moderate			35.7	8.3	20.0	7.9			17.4	6.6					25.7	9.2
Bad			30.3	12.8	11.7	8.1			11.7	8.1					30.5	12.6
Very-Bad			75.3	26.3	0.0	0.0			75.3	26.3					75.3	26.3
TOTAL			22.7	3.5	11.5	2.8			11.6	2.8					18.8	3.7

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Confidentiality		Choice of health care provider		Basic Amenities		Social Support	
	Talk Privately	Confidentiality of records			Cleanliness	Space	Family Visit	External Contact
	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>								
Female	15.1	3.4			13.1	3.3	7.0	2.5
Male	11.7	5.0			11.1	3.8	7.3	3.0
<b>Area of Residence</b>								
Urban	14.3	3.3			13.6	3.4	9.4	2.9
Rural	13.3	5.1			11.1	3.9	4.6	2.3
<b>Age Group</b>								
18-29y	13.6	10.3			5.6	4.0	2.8	2.8
30-44y	16.6	4.4			18.9	5.0	10.8	4.2
45-59y	10.4	3.8			11.3	4.0	6.0	3.0
60-69y	18.1	7.9			11.4	6.2	3.5	3.5
70+y	7.9	5.1			2.7	2.6	7.3	5.3
<b>Education Years</b>								
0								
1-5	0.0	0.0			9.3	11.2	0.0	0.0
6-11	17.3	6.0			8.1	4.6	4.9	3.0
12+	15.0	4.6			14.6	3.3	7.7	2.6
<b>Income Quintiles</b>								
Q1	9.9	4.2			6.1	3.0	6.0	3.6
Q2	20.9	7.6			11.2	5.2	9.0	5.3
Q3	16.7	7.0			8.7	4.9	10.2	5.6
Q4	13.7	5.4			12.2	4.8	4.1	3.0
Q5	11.3	4.8			19.7	7.1	7.4	3.8
<b>Condition</b>								
Child-Visit	32.3	10.8			11.3	6.2	26.2	10.2
Fever	3.3	3.2			0.0	0.0	0.0	0.0
Childbirth	16.3	8.5			16.0	7.5	16.3	8.5
Arthritis	9.4	7.0			0.0	0.0	0.0	0.0
Asthma	38.3	14.2			18.1	10.7	13.7	9.8
Heart-Disease	15.2	10.8			4.8	4.8	0.0	0.0
Injury	6.7	5.0			3.7	3.7	5.7	5.6
Minor-Surgery	22.9	8.2			26.7	8.2	7.1	4.1
Other	10.4	3.6			10.3	3.5	6.4	2.7
<b>Ownership</b>								
Government	14.8	3.4			12.9	2.7	7.0	2.0
Private	0.0	0.0			2.4	2.3	0.0	0.0
NGO								
Other								
<b>Self-Assessed Health</b>								
Very-Good	13.7	5.3			6.5	3.5	6.0	3.6
Good	14.2	4.2			17.9	3.8	8.1	3.6
Moderate	24.7	6.6			21.8	8.2	4.1	2.8
Bad	5.8	5.6			11.8	8.1	23.0	11.1
Very-Bad	0.0	0.0			0.0	0.0	0.0	0.0
TOTAL	13.8	3.0			12.4	2.6	7.1	1.9

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"

	Prompt Attention		Dignity		Communication		Autonomy		Confidentiality		Choice		Basic Amenities		Social Support	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>																
Female																
Male																
<b>Area of Residence</b>																
Urban																
Rural																
<b>Age Group</b>																
18-29y																
30-44y																
45-59y																
60-69y																
70+y																
<b>Education Years</b>																
0																
1-5																
6-11																
12+																
<b>Income Quintiles</b>																
Q1																
Q2																
Q3																
Q4																
Q5																
<b>Ownership</b>																
Government																
Private																
NGO																
Other																
<b>Self-Assessed Health</b>																
Very-Good																
Good																
Moderate																
Bad																
Very-Bad																
<b>Type of Care</b>																
Ambulatory-Adul																
Ambulatory-Chil																
Inpatient-Adult																
Inpatient-Child																
Non-Users																
TOTAL																

This table shows the percent of people who rated a responsiveness domain as "Extremely Important". All survey respondents were queried on their perception of the importance of each of the eight responsiveness domains in section 7100 of the WHS where each domain was described in detail (Q7100-Q7107). Respondents were asked to say whether the domain was "Extremely Important", "Very Important", "Moderately Important", "Slightly Important" or "Not



Important".

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, ownership (of health facility), self-assessed health, and type of care. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, self-assessed health and type of care sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad". Type of care is a constructed sample sub-group combining information from Q7000, Q7001 and Q7016. There are five categories of type of care: "Ambulatory adult", "Ambulatory child", "Inpatient adult", "Inpatient child" and "Non users".

---

## 8.0 Expenditure Background Indicators (2002)

---

GDP per capita (\$)	31,033
Total health expenditure per capita (\$)	2,255
Total population (000)	3,911
Life expectancy at birth (Both sexes)	77.1
Life expectancy at birth (male)	74.4
Life expectancy at birth (female)	79.8
Healthy life expectancy at birth (Both sexes)	69.8
Healthy life expectancy at birth (Male)	68.1
Healthy life expectancy at birth (Female)	71.5
Government funded health expenditure as a percentage of total health expenditure	75.2
Social security funded health expenditure as a percentage of total health expenditure	0.6
Private health insurance premiums as a percentage of total health expenditure	5.4
Out-of-pocket payment as a percentage of total health expenditure	13.2









## 10.1 Inability to control the important things in life - q8000

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
<b>Gender</b>													
Male	33.5	3.8	20.3	2.6	31.9	3.3	11.7	2.4	1.0	0.4	1.6	0.7	497
Female	36.8	3.6	24.0	3.4	29.6	2.8	4.7	1.1	3.4	1.1	1.5	0.9	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	33.7	3.9	20.2	3.0	32.0	3.3	9.6	2.3	2.4	0.8	2.0	0.9	611
Rural	37.3	5.0	25.1	3.8	28.9	4.3	5.9	1.5	1.9	0.8	0.8	0.5	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	32.0	4.6	18.2	4.2	32.0	3.8	12.8	2.9	2.9	1.2	2.0	0.9	153
Q2	28.7	4.3	21.1	4.0	33.5	4.6	11.3	3.3	3.6	1.8	1.8	1.0	179
Q3	35.2	4.7	21.5	3.7	31.5	4.1	7.7	2.3	2.2	1.2	2.0	1.3	205
Q4	40.3	4.6	21.1	3.6	31.6	4.8	5.4	1.8	1.6	1.0	0.0	0.0	220
Q5(richest)	37.8	4.6	28.3	3.9	25.0	4.2	5.9	2.0	1.1	0.6	1.8	1.2	220
M	33.0	8.7	16.4	6.7	37.3	9.5	6.2	5.2	3.2	3.2	4.0	3.6	37
<b>Age</b>													
18-29	42.3	4.4	21.6	3.5	26.0	3.4	7.0	2.2	1.1	0.6	2.0	1.1	282
30-44	34.7	4.6	20.9	3.2	31.3	3.9	8.6	2.5	2.9	1.2	1.7	0.9	279
45-59	29.7	4.0	22.7	3.0	34.3	3.8	8.7	2.2	3.4	1.3	1.3	0.9	242
60-69	32.9	6.1	23.9	5.0	30.8	4.9	9.7	3.1	0.9	0.6	1.8	1.2	103
70-79	35.4	6.0	22.6	5.5	30.4	5.2	7.9	3.5	3.2	3.1	0.7	0.7	71
80+	25.7	11.2	27.0	16.4	40.9	14.6	6.4	6.0	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	35.2	3.0	22.2	2.4	30.8	2.6	8.2	1.5	2.2	0.6	1.5	0.6	1,013

## 10.2 Inability to cope - q8001

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
<b>Gender</b>													
Male	31.6	3.4	20.8	2.9	30.6	3.2	13.1	3.0	2.4	0.9	1.5	0.7	497
Female	29.2	3.4	23.5	3.2	34.3	2.9	6.6	1.4	5.4	1.5	1.0	0.7	516
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	27.6	3.4	21.3	3.2	33.5	3.0	11.1	3.0	5.0	1.3	1.5	0.7	611
Rural	34.7	4.8	23.5	4.0	31.0	4.2	7.8	2.2	2.2	0.9	0.8	0.5	402
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	27.7	4.8	16.8	3.8	34.2	4.3	11.9	2.9	7.6	2.2	1.7	0.8	153
Q2	25.9	4.1	18.2	4.1	33.7	4.3	15.3	5.0	5.2	2.2	1.8	1.0	179
Q3	31.1	4.4	22.5	3.6	29.6	4.0	10.8	2.5	5.2	1.8	0.8	0.6	205
Q4	30.5	4.1	28.6	4.2	33.5	4.7	4.6	1.5	2.7	1.6	0.0	0.0	220
Q5(richest)	35.5	4.2	24.8	4.1	29.7	3.8	7.7	2.7	0.5	0.4	1.8	1.2	220
M	28.3	8.7	6.6	3.5	46.0	9.4	11.9	6.1	3.2	3.2	4.0	3.6	37
<b>Age</b>													
18-29	38.0	4.3	22.4	3.7	28.5	3.3	7.6	2.3	2.4	1.3	1.1	0.6	282
30-44	28.2	4.0	21.1	4.0	32.4	4.0	12.3	3.5	4.3	1.5	1.7	0.9	279
45-59	24.9	3.7	22.8	3.0	36.3	4.2	9.5	2.6	5.4	1.7	1.1	0.8	242
60-69	30.6	6.4	23.3	4.9	30.7	5.3	10.9	3.4	2.7	1.5	1.8	1.2	103
70-79	29.8	6.1	19.5	4.5	33.8	6.3	9.2	3.7	7.0	3.7	0.7	0.7	71
80+	25.5	10.8	25.0	16.6	41.1	14.7	8.4	6.3	0.0	0.0	0.0	0.0	36
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	30.4	2.8	22.1	2.5	32.5	2.5	9.8	2.0	3.9	0.8	1.3	0.5	1,013







