

# **REPORT OF URUGUAY**

## TABLE OF CONTENTS

1	Introduction
2	Methodology
2.1	Sampling
2.2.1	Geographic Component of the Survey
2.2.2	The map
2.3	Survey Metrics
2.3.1	Sample deviation index
2.3.2	HH Sample deviation index
2.3.3	Response rate
2.3.4	Missing data
2.3.5	Reliability
3	Socio-Demographic Profile
3.1	Household composition by residence and age groups according to income quintile
3.2	Selected age distributions by income quintile
3.3	Characteristics of head of household by income quintile
3.4	Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex
3.5	Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education
4	Risk Factors
4.1	Prevalence of smoking and average daily tobacco consumption
4.2	Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers
4.3	Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

- 4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water
- 4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility
- 4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used
- 5 Coverage
  - 5.01 Tuberculosis
  - 5.02 HIV Prevention and Screening
  - 5.03 Antenatal Care and Mother to Child Transmission of HIV
  - 5.04 Birth Delivery Care
  - 5.05 Child's Health - Common Illness
  - 5.06 Child's Health - Danger Signs
  - 5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card
  - 5.08 Vaccination Status of Children 12-23 Months
  - 5.09 Women's Cancer Screening
  - 5.10 Oral Health and Injuries
  - 5.11 Self Reported Non-Communicable Conditions
  - 5.12 Mental Health Conditions
  - 5.13 Cataracts
- 6 Health State Descriptions
  - 6.01 In general, rate your health today - q2000
  - 6.02 Difficulty with 'work and household activities' in last 30 days - q2001
  - 6.03 Difficulty with 'moving around' in last 30 days - q2010
  - 6.04 Difficulty with 'vigorous activities' in last 30 days - q2011
  - 6.05 Difficulty with 'self-care' in last 30 days - q2020
  - 6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021

- 6.07 Amount of 'bodily aches or pains' in last 30 days - q2030
- 6.08 Amount of 'bodily discomfort' in last 30 days - q2031
- 6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050
- 6.10 Difficulty with 'learning a new task' in last 30 days - q2051
- 6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060
- 6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061
- 6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071
- 6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072
- 6.15 Problem with 'sleeping' in last 30 days - q2080
- 6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081
- 6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090
- 6.18 Problem with 'worry or anxiety' in last 30 days - q2091
- 6.19 Adjusted for vignette responses
- 7 Health System Responsiveness
  - 7.1 Self-reported utilization of health services and perceived unmet need
  - 7.2 Patient assessed characteristics of ambulatory health services
    - 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1
    - 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2
  - 7.4 Patient assessed characteristics of hospital inpatient health services
    - 7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1
    - 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2
  - 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"
- 8 Health Expenditure, Insurance and Human Resources for Health

- 8.0 Expenditure Background Indicators (2002)
- 8.1 Distribution of health payments across different population groups
- 8.2 Percent distribution of structure of out-of-pocket health payments
- 8.3 Financial sources used by households for paying for health services
- 8.4 Insurance Coverage
- 8.5 Human resources for health
- 9.1 Adult Mortality Chart 1
- 9.2 Adult Mortality Chart 2
- 9.3 Adult Mortality Chart 3
- 9.4 Adult Mortality Chart 4
- 10 Stress and Social Capital
- 10.1 Inability to control the important things in life - q8000
- 10.2 Inability to cope - q8001
- 10.3 Satisfaction with your health - q8002
- 10.4 Feeling safe when alone at home - q8011
- 10.5 Feeling safe when walking down the street alone after dark - q8012

# 1. Introduction

The World Health Organization initiated the World Health Survey (WHS) as a *data collection platform* to obtain information on the health of populations and on health systems. The WHS responds to the need of countries for a detailed and sustainable health information system and gathers data through surveys to measure essential population health parameters. The WHS uses standard survey procedures and instruments for general population surveys to gather comparable data across WHO Member States. These methods and instruments have been refined through scientific review of literature, extensive consultations with international experts and large-scale pilot tests. WHS is designed to evolve through its implementation by continuous input from collaborators, including policy makers, survey institutions, scientists and other interested parties. The data is jointly owned by the countries and the WHO, and there is a commitment for a long-term data collection, building local capacity and using this information for health policy.

The objectives of the World Health Survey were to:

- develop a **means** of providing:
  - **valid, relevant, reliable** and **comparable** information on **outcomes, functions** and **inputs** to health systems
  - to **supplement** the information provided by routine **HIS**
- build the **evidence base** necessary for policy-makers:
  - to **routinely monitor** if health systems are achieving the **desired goals**
  - and in some countries, to assess if the **increase in health resources** is having the desired effect

The introductory chapter of each country report will provide details on the current sources of data from routine health information systems and ad hoc surveys. It will describe the health situation in the country and identify gaps. It will also present a brief overview on the health system organization within the country.

## 2. Sampling

The WHO sampling guidelines emphasized scientific principles of survey sampling as explicit standards for quality, included examples of good sampling plans, and identified quality assurance standards for countries to adhere to. WHO and technical advisers provided technical support to countries when needed. The following principles were used for sampling in the WHS:

1. The sample should target the *de facto* population (i.e. all people living in that country including guest workers, immigrants and refugees) and not *de jure* population (i.e. the citizens of that country alone) with complete geographic coverage of the country, if possible, or a representative geographic segment of the country.
2. A *sample frame* with 90% coverage that included all key subgroups of interest. Enumeration or listing of households to update the frame was recommended.
3. The WHS sample identified all adult members of the general population aged 18 years or older as its sample. Households are selected using a random, stratified sampling procedure. One individual per household is then selected through a random selection procedure using the Kish table method.
4. The WHS used a sampling strategy which encompasses a known non-zero selection probability for any individual included in the study. This makes it possible to extrapolate the data to the whole population. Households and individuals are selected with a known probability in order to provide selection probabilities for each household and individual in the data set.
5. The WHS sampling strategy was without replacement and categories of non-respondents were clearly defined.
6. The WHS used GPS devices in certain countries and digitized maps to geocode the data within specified guidelines.

Each national report will present a detailed description of the sampling strategy and the allocation tables

---

## 2.2.1 Geographic Component of the Survey

---

Geographic Information refers to any information associated with a location on the Earth's surface. This location can be associated to a coordinate (latitude/longitude), a place's name, a postal address or any other geographic object.

Household Surveys represent an important source of information. If the geographic location of the surveyed clusters is collected at the same time that the interview is conducted users are then able to integrate the spatial dimension in their analysis

The geographic component of the WHO WHS is stored in two sections of the questionnaire:

- 0100 Sampling Information allowing the link between the location of each cluster and the sampling frame used for the survey. This section has also been used in order to collect the type of setting in which each household was located at the time of the survey. This section of the questionnaire has been used for all the countries part of the WHO WHS
- 0200 Geocoding Information for the collection of the latitude and longitude of each household. This section has only been used in 27 countries where GPS devices were available or have been provided. The choice of these countries has been done on the basis of the availability of digital maps delimiting the surveyed clusters, the field data collection planning and their distribution within the different WHO regions. Specific training and field material have been prepared and used by the interviewers or field supervisors in these countries in order to collect the latitude and longitude of each of the surveyed household.


As the sampling design may not always include the administrative structure of the country and as the codes entered in the data set for the sampling level needed to be labelled, a key correspondence table containing this information has been collected after the survey itself.

Once all the survey data and the key correspondence table available a specific protocol has been applied to the two sections mentioned earlier in order to clean eventual data collection or data entry mistakes as well as to fill the gaps in close collaboration with the survey institution.

The questionnaire as well as the training material and data cleaning protocols used can be downloaded from the WHO WHS web site (<http://www.who.int/healthinfo/survey/en/index.html>) in the "Survey Instrument and Related Documents" section.




2.3.1 Sample deviation index



SDI

2.3.2 HH Sample deviation index



HDI

---

### 2.3.3 Response rate

---

---

#### Household

Selected	3,000
----------	-------

Interviewed	2,982
-------------	-------

Percentages	99
-------------	----

#### Individual


Selected	2,984
----------	-------

Interviewed	2,976
-------------	-------


Percentages	100
-------------	-----

---

### 2.3.4 Missing data



### 2.3.5 Reliability



### 3.1 Household composition by residence and age groups according to income quintile

	Percent Rural	SE	Mean Number of									
			Household members	SE	Children < 2 years	SE	Children < 5years	SE	Children 5 -17 years	SE	Persons 18+ years	SE
Q1(poorest)	20.6	5.43	2.5	0.09	0.1	0.01	0.2	0.02	0.6	0.05	1.8	0.05
Q2	10.3	3.45	2.6	0.12	0.0	0.01	0.2	0.01	0.5	0.07	1.9	0.07
Q3	6.2	2.87	2.6	0.14	0.0	0.01	0.1	0.02	0.5	0.06	2.0	0.07
Q4	4.1	2.69	2.9	0.15	0.0	0.01	0.2	0.03	0.6	0.06	2.1	0.08
Q5(richest)	1.7	1.21	3.2	0.04	0.1	0.01	0.2	0.02	0.6	0.03	2.4	0.05
All	7.8	3.62	2.8	0.04	0.1	0.01	0.2	0.01	0.6	0.03	2.1	0.03

Note: table based on de jure members of household i.e. usual residents

---

### 3.2 Selected age distributions by income quintile

---

	<b>Females Age 18–49</b>	<b>Persons Age 50+</b>	<b>Persons Age 60+</b>
Q1(poorest)	242	352	257
Q2	324	454	283
Q3	398	472	300
Q4	453	538	288
Q5(richest)	527	533	233
All	1,943	2,349	1,362

Note: table based on de jure members of household i.e. usual residents

---

### 3.3 Characteristics of head of household by income quintile

---

	Mean Age of Head	SE	Median Education of Head	Percent Female Head	SE
Q1 (poorest)	51	1.51	3	40.7	3.94
Q2	50	0.87	3	37.3	4.73
Q3	50	0.95	3	37.1	6.66
Q4	49	0.47	4	35.7	2.04
Q5 (richest)	48	0.30	6	27.4	0.93
All	50	0.33	4	35.4	2.54

#### Education

1=no formal education

2=less than primary school

3=primary school completed

4=secondary school completed

5=high school (or equivalent) completed

6=college/pre-university/university completed

7=post graduate degree completed

Note: table based on de jure members of household i.e. usual residents



### 3.4 Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex

	Male	SE	Female	SE	Missing	Total	SE
<b>Age</b>							
0-4	6.3	0.30	5.1	0.50		5.70	0.30
5-14	16.0	0.60	14.0	1.20		15.00	0.70
15-29	24.6	0.80	23.4	0.70		24.00	0.30
30-44	19.8	0.40	20.4	0.50		20.10	0.40
45-59	17.9	0.70	19.2	0.80		18.60	0.40
60-69	9.0	0.40	8.7	0.50		8.80	0.30
70-79	4.8	0.40	6.3	0.70		5.60	0.50
80+	1.6	0.20	2.9	0.30		2.30	0.20
Missing	0.1	0.00	0.0	0.00		0.00	0.00
<b>Residence</b>							
Urban	91.4	3.80	91.4	4.00		91.40	3.90
Rural	8.6	3.80	8.6	4.00		8.60	3.90
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Marital_Status</b>							
Never	46.9	0.80	41.2	1.60		44.00	0.70
Currently	36.4	0.90	33.3	0.70		34.80	0.70
Separated	4.1	0.40	3.3	0.40		3.70	0.20
Divorced	3.6	0.40	7.2	0.80		5.50	0.40
Widowed	3.0	0.40	9.6	0.80		6.40	0.40
CoHabiting	5.8	0.50	5.3	0.40		5.60	0.40
Missing	0.1	0.10	0.1	0.00		0.10	0.10
<b>Education</b>							
NoFormal	7.5	0.60	6.7	0.70		7.10	0.60
LessthanPrimary	16.4	1.30	13.9	1.40		15.10	1.30
Primary	33.9	5.20	31.9	4.10		32.90	4.60
Secondary	21.9	2.70	22.2	2.30		22.00	2.50
HighSchool	4.6	0.40	4.8	0.50		4.70	0.40
College	14.7	3.80	19.2	3.60		17.10	3.70
PostGraduate	0.8	0.30	1.3	0.40		1.00	0.30
Missing	0.2	0.10	0.1	0.00		0.20	0.10
Total	100.0		100.0			100.00	
Number	4,025.0		4,354.0			8,379.00	
Number of Households						2,982.00	

### 3.5 Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education

	Male	SE	Female	SE	Missing	Total	SE
<b>Age</b>							
18-29	28.9	1.70	24.6	1.10		26.70	0.90
30-44	28.4	1.10	26.2	1.20		27.20	1.00
45-59	21.9	0.90	20.8	1.10		21.40	0.80
60-69	10.3	0.60	12.2	0.80		11.30	0.60
70-79	7.5	0.70	10.3	1.10		9.00	0.80
80+	2.9	0.50	5.9	0.50		4.50	0.40
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Residence</b>							
Urban	92.7	3.10	92.9	3.80		92.80	3.40
Rural	7.3	3.10	7.1	3.80		7.20	3.40
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Marital_Status</b>							
Never	30.0	1.70	23.1	0.70		26.40	0.90
Currently	49.9	1.20	41.6	1.70		45.50	1.30
Separated	4.6	0.40	4.1	0.40		4.30	0.30
Divorced	3.9	0.50	10.2	0.90		7.20	0.60
Widowed	3.7	0.50	14.0	0.70		9.10	0.60
CoHabiting	7.9	0.80	7.1	0.80		7.50	0.70
Missing	0.0	0.00	0.0	0.00		0.00	0.00
<b>Education</b>							
NoFormal	1.1	0.40	1.1	0.50		1.10	0.40
LessthanPrimary	9.9	2.60	8.8	1.20		9.30	1.80
Primary	35.2	5.60	28.7	3.90		31.80	4.80
Secondary	24.5	2.80	26.9	1.90		25.80	2.30
HighSchool	8.0	1.10	7.6	1.30		7.80	0.90
College	20.0	4.60	24.4	4.20		22.30	4.40
PostGraduate	1.4	0.50	2.5	0.40		1.90	0.40
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Total	100.0		100.0			100.00	
Number	1,414.0		1,563.0			2,977.00	

#### 4.1 Prevalence of smoking and average daily tobacco consumption

	Current daily smoker	SE	Smoker, not daily	SE	Non-smoker	SE	Missing	Average daily tobacco consumption*	SE	Number of respondents
<b>Sex</b>										
Male	33.3	1.20	5.0	0.70	61.1	1.30	0.6	17.4	0.90	1,414
Female	23.1	1.70	5.0	0.50	71.5	2.00	0.3	14.0	0.30	1,563
Missing										0
<b>Residence</b>										
Urban	28.1	1.10	5.2	0.40	66.2	1.20	0.5	15.9	0.50	2,763
Rural	26.0	2.30	2.3	0.80	71.4	2.60	0.2	15.6	0.90	214
Missing										0
<b>Income Quintile</b>										
Q1	34.5	2.90	3.6	1.20	61.7	3.50	0.2	16.0	1.30	417
Q2	32.2	2.40	3.3	0.70	63.4	2.30	1.1	14.5	0.70	528
Q3	27.7	2.40	3.5	1.00	68.7	2.70	0.1	16.5	0.70	601
Q4	26.9	3.00	5.0	1.50	67.7	2.30	0.4	16.2	0.50	651
Q5	23.2	1.30	8.4	0.60	68.3	1.10	0.1	16.3	0.30	750
Missing										31
<b>Age</b>										
18-29	36.2	1.50	8.6	1.80	54.6	2.30	0.6	14.0	0.50	793
30-44	32.6	1.80	5.1	0.50	61.9	1.90	0.5	16.5	0.70	811
45-59	30.0	2.30	3.0	0.60	66.6	2.70	0.4	18.0	1.30	636
60-69	16.6	1.70	3.8	0.80	78.9	1.70	0.7	16.9	1.30	337
70-79	12.4	1.90	3.1	1.50	84.5	2.00	0.0	13.6	1.10	268
80+	0.8	0.40	0.0	0.00	99.2	0.40	0.0	3.0	0.00	133
Missing										0
All	28.0	1.10	5.0	0.50	66.6	1.30	0.5	15.9	0.40	2,977

\* Denominator used is current daily smoker

## 4.2 Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers

	Life time abstainers	SE	Non-heavy drinkers	SE	Infrequent Heavy Drinkers*	SE	Frequent Heavy drinkers**	SE	Missing	Number of respondents
<b>Sex</b>										
Male	25.1	3.80	68.0	3.80	5.5	0.70	1.3	0.30	0.1	1,414
Female	45.7	6.70	52.6	6.50	1.6	0.30	0.1	0.00	0.1	1,563
Missing										0
<b>Residence</b>										
Urban	35.1	4.70	60.6	4.70	3.5	0.40	0.7	0.10	0.1	2,763
Rural	46.2	5.40	51.4	5.20	2.2	0.80	0.3	0.20	0.0	214
Missing										0
<b>Income Quintile</b>										
Q1	44.4	6.20	52.5	6.50	1.8	0.60	1.3	0.70	0.0	417
Q2	37.8	4.70	57.6	5.10	3.7	1.70	0.9	0.40	0.0	528
Q3	39.9	5.50	58.0	5.00	1.9	0.60	0.3	0.10	0.0	601
Q4	34.0	2.90	61.8	4.10	3.9	1.60	0.3	0.10	0.0	651
Q5	28.3	3.60	65.9	2.80	4.9	1.00	0.8	0.10	0.0	750
Missing										31
<b>Age</b>										
18-29	23.0	6.00	67.9	5.30	7.8	1.30	1.2	0.60	0.2	793
30-44	37.4	5.20	58.1	5.20	3.7	0.50	0.7	0.30	0.1	811
45-59	36.9	5.90	61.7	6.00	1.0	0.40	0.4	0.20	0.0	636
60-69	41.8	4.40	56.9	4.80	0.6	0.50	0.7	0.30	0.0	337
70-79	48.6	3.30	51.4	3.30	0.0	0.00	0.0	0.00	0.0	268
80+	58.7	5.00	40.3	4.90	1.0	1.10	0.0	0.00	0.0	133
Missing										0
All	35.9	4.70	59.9	4.60	3.4	0.40	0.7	0.10	0.1	2,977

\*Infrequent heavy drinkers: 1-2 days with 5 or more standard drinks per week(last 7 days)

\*\*Frequent heavy drinkers: 3 or more days with 5 or more standard drinks per week(last 7 days)

### 4.3 Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

	Fruit and vegetable intake					Physical Activity					Number of respondents
	*Insufficient	SE	Sufficient	SE	Missing	**Insufficient	SE	Sufficient	SE	Missing	
Sex											
Male	75.2	1.40	24.8	1.40	0.0	20.4	2.20	79.6	2.20	0.1	1,414
Female	70.0	2.10	30.0	2.10	0.0	25.9	2.30	74.0	2.30	0.1	1,563
Missing											0
Residence											
Urban	72.7	1.10	27.3	1.10	0.0	23.2	2.20	76.7	2.20	0.1	2,763
Rural	69.4	4.40	30.6	4.40	0.0	23.8	5.40	76.2	5.40	0.0	214
Missing											0
Income Quintile											
Q1	78.4	3.00	21.6	3.00	0.0	21.8	4.40	78.2	4.40	0.0	417
Q2	72.5	2.70	27.5	2.70	0.0	22.8	3.90	77.2	3.90	0.0	528
Q3	75.2	2.20	24.8	2.20	0.0	27.9	3.50	72.1	3.50	0.0	601
Q4	68.5	2.30	31.5	2.30	0.0	24.2	2.90	75.8	2.90	0.0	651
Q5	70.3	1.30	29.7	1.30	0.0	19.8	1.50	80.2	1.50	0.0	750
Missing											31
Age											
18-29	78.1	2.60	21.9	2.60	0.0	15.8	1.80	84.0	1.80	0.2	793
30-44	72.1	2.00	27.9	2.00	0.0	22.9	2.60	77.0	2.60	0.1	811
45-59	68.4	1.30	31.6	1.30	0.0	22.4	2.60	77.6	2.60	0.0	636
60-69	75.6	1.90	24.4	1.90	0.0	30.5	2.70	69.5	2.70	0.0	337
70-79	67.5	2.90	32.5	2.90	0.0	31.8	4.90	68.2	4.90	0.0	268
80+	63.4	7.80	36.6	7.80	0.0	39.0	6.00	61.0	6.00	0.0	133
Missing											0
All	72.5	1.10	27.5	1.10	0.0	23.3	2.00	76.6	2.10	0.1	2,977

\*Insufficient intake of fruit/vegetables: less than 5 servings of fruit/vegetables per one typical day

\*\*Insufficient physical activity: less than 150 minutes per week(last 7 days) spent on walking/moderate activity/vigorous activity

#### 4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water

	Piped into households	SE	other improved sources	SE	Total improved sources	SE	No improved sources	SE	Missing	Number of households
<b>Residence</b>										
Urban	96.9	0.50	2.4	0.50	99.4	0.10	0.1	0.10	0.6	2,748
Rural	84.1	4.10	15.4	4.10	99.5	0.40	0.5	0.40	0.0	234
Missing										0
<b>Income_Quintile</b>										
Q1(poorest)	85.2	2.30	13.3	2.20	98.5	0.50	0.5	0.30	1.0	477
Q2	96.1	1.10	3.3	1.10	99.5	0.40	0.1	0.10	0.4	566
Q3	96.9	0.60	2.6	0.60	99.5	0.30	0.0	0.00	0.5	622
Q4	99.0	0.40	0.6	0.50	99.6	0.20	0.0	0.00	0.4	638
Q5(richest)	99.5	0.20	0.0	0.00	99.5	0.20	0.0	0.00	0.5	643
Missing										36
Total	95.9	0.80	3.4	0.80	99.4	0.10	0.1	0.10	0.5	2,982

other improved sources: public standpipe/protected tube well/borehole/protected dug well/protected spring/rainwater collection

no improved sources: unprotected well/unprotected spring/water taken directly from a pond or stream/tanker truck water/vendor provided water

#### 4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility

	Flush toilet to sewage system	SE	Other improved toilet facilities	SE	Total improved sanitation	SE	No improved sanitation	SE	Missing	Number of households
<b>Residence</b>										
Urban	80.5	4.50	19.0	4.50	99.4	0.10	0.5	0.10	0.1	2,748
Rural	41.8	8.60	56.4	8.40	98.3	0.60	1.6	0.70	0.1	234
Missing										0
<b>Income_Quintile</b>										
Q1(poorest)	42.8	4.80	55.6	4.60	98.3	0.40	1.6	0.50	0.0	477
Q2	67.6	3.90	31.7	3.80	99.3	0.30	0.7	0.30	0.0	566
Q3	85.3	3.30	14.3	3.40	99.6	0.20	0.4	0.20	0.0	622
Q4	88.2	4.00	11.3	4.00	99.5	0.20	0.3	0.20	0.2	638
Q5(richest)	93.8	2.20	6.0	2.10	99.7	0.10	0.1	0.10	0.2	643
Missing										36
Total	77.5	5.40	21.9	5.40	99.4	0.10	0.6	0.10	0.1	2,982

other improved sanitation facilities: connection to septic system/pour-flush latrine/covered dry latrine(with privacy) [provided facilities not shared]  
no improved sanitation: uncovered dry latrine(without privacy)/bucket latrine/no facilities(open defecation)

#### 4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used

	Electricity, Gas	SE	Kerosene	SE	Solid fuels on improved stove	SE	Solid fuels on traditional stove	SE	Total solid fuel use	SE	Missing	Number of households
<b>Residence</b>												
Urban	98.8	0.10	0.4	0.10	0.0	0.00	0.4	0.10	0.4	0.10	0.3	2,748
Rural	92.5	2.60	0.3	0.30	2.3	1.20	4.9	2.20	7.2	2.60	0.0	234
Missing												0
<b>Income_ Quintile</b>												
Q1(poorest)	91.8	1.40	2.5	1.00	1.1	0.60	4.1	1.20	5.3	1.50	0.5	477
Q2	98.8	0.50	0.0	0.00	0.2	0.10	0.3	0.20	0.5	0.30	0.7	566
Q3	100.0	0.00	0.0	0.00	0.0	0.00	0.0	0.00	0.0	0.00	0.0	622
Q4	99.8	0.10	0.0	0.00	0.0	0.00	0.0	0.00	0.0	0.00	0.2	638
Q5(richest)	99.6	0.10	0.0	0.00	0.0	0.00	0.0	0.00	0.0	0.00	0.4	643
Missing												36
Total	98.3	0.30	0.4	0.10	0.2	0.10	0.7	0.30	0.9	0.30	0.3	2,982



## 5.01 Tuberculosis

		Tuberculosis			
	Total	Need	Coverage	SE	M
Sex					
Female	1,563	4.0	20.2	3.3	0.0
Male	1,414	2.3	17.6	9.3	1.3
Missing	0				
Residence					
Urban	2,763	3.2	19.0	3.5	0.5
Rural	214	3.9	22.5	10.0	0.0
Missing	0				
Income_Quintile					
Q1(poorest)	417	4.3	31.7	11.3	0.0
Q2	528	3.0	23.9	12.9	0.0
Q3	601	2.0	10.4	5.3	0.0
Q4	651	4.1	19.8	7.8	0.0
Q5(richest)	750	3.1	10.9	3.2	1.9
Missing	31				
Age					
18-29	793	2.5	11.5	5.7	0.0
30-44	811	2.9	9.3	6.7	0.0
45-59	636	2.6	46.3	8.9	0.0
60-69	337	6.5	11.8	10.6	2.0
70-79	268	3.8	36.3	11.6	0.0
80+	133	3.0	0.0	0.0	0.0
Missing	0				
Total	2,977	3.2	19.3	3.3	0.5

**M = % Missing of Covered      SE= Standard Error of % Covered**

**Tuberculosis Indicators:**

**"Need" = % of all respondents with self-reported continuous cough lasting 3 weeks, OR had blood in phlegm, OR had coughed blood in the last 12 months**

**"Coverage" = % of respondents with self reported symptoms who indicated receiving a TB screening test in the last 12 months.**

## 5.02 HIV Prevention and Screening

	Total (persons of reproductive age 18-49)	Voluntary Counselling and Testing for HIV/AIDS				Condom Usage			
		Need	Coverage	SE	M	Need	Coverage	SE	M
Sex									
Females	912	82.6	19.6	3.4	13.6	27.4	58.7	2.0	2.5
Males	926	95.0	6.0	2.1	65.8	42.1	78.5	3.0	1.2
Missing									
Residence									
Urban	1,704	88.6	12.9	2.9	41.5	35.4	70.6	2.2	1.8
Rural	134	92.6	4.7	1.6	43.7	27.5	72.7	7.0	0.0
Missing									
Income_Quintile									
Q1(poorest)	233	89.2	10.1	2.2	46.9	33.7	70.0	5.6	1.0
Q2	309	87.7	10.4	2.9	49.1	41.5	76.4	4.0	0.6
Q3	366	87.7	10.7	2.9	39.2	33.4	77.7	3.9	1.3
Q4	401	90.9	12.9	4.0	41.1	31.8	68.0	3.4	3.8
Q5(richest)	514	89.9	14.5	2.1	37.6	34.9	65.2	2.4	1.7
Missing	16								
Total	1,838	88.9	12.3	3.0	41.7	34.8	70.7	2.2	1.7

**M = % Missing of Covered      SE= Standard Error of % Covered**

**HIV Prevention and Screening Indicators:**

**"Total" = number of respondents of reproductive age.**

**"Voluntary Counselling and Testing for HIV/AIDS Need" = % of respondents of reproductive age who are sexually active, i.e. have had sexual intercourse in the last 12 months.**

**"Voluntary Counselling and Testing for HIV/AIDS Coverage" = % of respondents of reproductive age who are sexually active who indicated having taken an HIV test and receiving the results of test.**

**"Condom Usage Need" = % of respondents of reproductive age whose last sexual encounter during the last 12 months is classified as risky sexual behavior, i.e. having had sexual intercourse with a non-cohabitating partner .**

**"Condom Usage Coverage" = % of respondents of reproductive age who used a condom last time had risky sex during the last 12 months (with non-cohabitating partner).**

### 5.03 Antenatal Care and Mother to Child Transmission of HIV

	Antenatal Care											Maternal to Child HIV Trans.				
	Women 18-49	Birth in last 5 years	Cov	SE	Visit 1x	Visit 2x	Visit 3x	Blood pressure checked	Blood sample taken	Told Signs of Pregnancy Complications	M	Birth in last 2 years	Cov	SE	M	
Residence																
Urban	842	25.3	75.9	2.9	83.2	82.9	82.9	91.6	91.8	83.8	11.6	10.5	63.0	8.8	4.3	
Rural	71	24.3	86.9	5.5	92.1	92.1	92.1	100.0	99.1	93.8	3.9	8.4	44.8	10.5	0.0	
Missing																
Income_Quintile																
Q1(poorest)	105	38.0	53.5	8.3	67.6	66.1	66.1	82.5	81.6	72.9	12.8	21.0	60.2	11.1	7.6	
Q2	138	22.6	80.3	7.9	94.7	94.7	94.7	96.7	96.7	82.3	4.2	9.2	77.7	15.8	0.0	
Q3	192	25.3	77.7	6.1	83.1	83.1	83.1	96.2	97.5	86.5	9.3	5.3	60.1	16.7	10.9	
Q4	204	23.2	86.2	4.5	90.7	90.7	90.7	93.9	93.9	89.4	9.3	10.6	59.8	16.0	4.7	
Q5(richest)	260	23.2	81.0	3.9	83.6	83.6	83.6	91.5	91.5	87.3	16.4	10.1	55.2	10.7	0.0	
Missing	14															
Total	912	25.2	76.7	3.0	83.9	83.6	83.6	92.2	92.3	84.5	11.0	10.4	61.9	7.9	4.0	

M = % Missing of Covered SE= Standard Error of %Covered

Antenatal Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least 3 times or more by health care professional, received blood pressure, blood sample taken, and told signs about complications for their last pregnancy (aggregate antenatal care indicator).

"Visit 1x" "Visit 2x" "Visit 3x" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least once, twice or three times or more respectively by a health care professional during their last pregnancy.

"Blood pressure checked" = % of female respondents of reproductive age with live birth in the last 5 years who had blood pressure taken when they saw health care professional during their last pregnancy.

"Blood sample taken" = % of female respondents of reproductive age with live birth in the last 5 years who had a blood sample taken when they saw health care professional during their last pregnancy.

"Told Signs of Pregnancy Complications" = % of female respondents of reproductive age with live birth in the last 5 years who had been told signs of pregnancy complications when they saw health care professional during their last pregnancy.

Maternal to Child HIV Transmission Indicators:

"Birth in last 2 years" = % of female respondents of reproductive age indicating having had a live birth in the last 2 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 2 years who indicated having been counseled and tested and given results for HIV during their last pregnancy.

## 5.04 Birth Delivery Care

	Women 18-49	Delivery Care				Place of Delivery						
		Need	Coverage	SE	M	Public Health Facility	SE	Private Health Facility	SE	Home	SE	M
Residence												
Urban	842	25.3	82.2	3.3	8.3	34.0	2.2	58.3	2.5	7.7	1.3	0.0
Rural	71	24.3	84.1	7.2	0.0	34.5	7.8	65.5	7.8	0.0	0.0	0.0
Missing												
Income Quintile												
Q1(poorest)	105	38.0	65.7	6.2	12.4	74.4	14.1	16.3	9.3	9.3	6.3	0.0
Q2	138	22.6	82.4	10.3	3.3	46.2	14.7	47.3	15.8	6.6	3.6	0.0
Q3	192	25.3	84.8	3.7	5.4	44.9	8.8	49.7	7.8	5.4	4.0	0.0
Q4	204	23.2	89.6	3.1	6.1	16.0	5.7	77.9	7.6	6.1	3.9	0.0
Q5(richest)	260	23.2	85.2	3.8	10.2	7.5	4.0	84.0	5.5	8.5	2.7	0.0
Missing	14											
Total	912	25.2	82.3	3.1	7.7	34.0	2.1	58.9	2.6	7.1	1.3	0.0

M = % Missing of Covered SE= Standard Error of % Covered Birth Delivery Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Coverage" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having received assistance with birth delivery from a health care professional (doctor, nurse or midwife) during their last birth.

"Public Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a public health facility.

"Private Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a private health facility.

"Home" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered home or in the field (etc.)

## 5.05 Child's Health - Common Illness

	Total Children under Age of 5	Fever	More liquid	SE	Same or more food	SE	M	Cough	More liquid	SE	Same or more food	SE	M	Diarrhea	More liquid	SE	Same or more food	SE	M
<b>Sex</b>																			
Females	190	31.4	45.7	5.6	29.9	8.8	1.0	18.7	52.5	9.1	40.0	31.4	3.0	8.8	16.6	8.3	48.2	8.6	0.0
Males	209	28.5	46.2	5.6	19.5	5.6	0.0	21.8	27.7	9.0	13.6	28.5	0.0	10.5	36.3	11.8	20.3	10.3	4.2
Missing	0																		
<b>Residence</b>																			
Urban	367	30.1	46.3	5.0	23.5	5.2	0.5	20.4	37.5	5.1	22.8	30.1	1.4	9.7	27.4	7.3	34.4	8.1	2.6
Rural	32	27.6	41.2	14.7	40.0	13.3	0.0	19.9	50.7	20.2	52.3	27.6	0.0	10.0	31.9	18.7	11.0	10.4	0.0
Missing																			
<b>Income_Quintile</b>																			
Q1(poorest)	66	21.0	38.7	21.4	16.1	10.4	0.0	13.2	36.0	15.0	18.4	21.0	0.0	6.9	3.6	3.8	54.4	22.1	0.0
Q2	62	19.7	35.9	9.5	37.9	15.9	5.0	22.7	34.2	12.2	60.7	19.7	0.0	5.7	73.9	23.6	66.4	27.5	0.0
Q3	87	33.0	48.9	12.8	31.2	13.9	0.0	23.4	41.9	14.8	1.7	33.0	0.0	23.0	30.3	12.1	38.8	11.9	0.0
Q4	90	35.1	53.5	11.0	31.9	8.0	0.0	29.2	42.7	12.1	38.3	35.1	4.0	2.9	35.1	32.2	0.0	0.0	35.1
Q5(richest)	92	33.3	39.8	5.0	12.0	6.5	0.0	12.6	32.2	14.9	0.0	33.3	0.0	8.2	13.5	4.9	0.0	0.0	0.0
Missing	3																		
Total	399	29.9	46.0	4.7	24.7	5.4	0.5	20.3	38.6	5.0	25.1	29.9	1.3	9.7	27.7	6.9	32.4	7.2	2.4

**M = % Missing of Covered      SE= Standard Error of % Covered**

**Child's Common Illness Indicators:**

**Total = Number of children under the age of 5 years at the time of the interview.**

**"Fever"/"Cough"/"Diarrhea" = % of children under the age of 5 with reported episode of (cough, fever, or diarrhea) in the last 3 months.**

**"More Liquid" = % of children under 5 with either fever, cough, or diarrhea in the last 3 months, that were given more liquids to drink during their illness.**

**"Same or More Food" = % of children under 5 with either fever, cough, or diarrhea in the last 3 months, that were given the same amount or more food to eat during their illness.**

## 5.06 Child's Health - Danger Signs

	Children under 5	Fast Breath	Seeking Treatment						Vomit	Seeking Treatment						No Drink or Eat	Seeking Treatment					
			Yes	SE	Public	Private	Trad	M		Yes	SE	Public	Private	Trad	M		Yes	SE	Public	Private	Trad	M
Sex																						
Females	190	0.1	100.0	0.0	27.8	53.9	18.3	0.0	3.9	100.0	0.0	9.0	68.2	22.9	0.0	4.5	100.0	0.0	54.1	45.9	0.0	0.0
Males	209	0.1	94.7	4.0	26.5	62.7	10.8	0.0	3.5	97.8	2.5	17.5	65.0	17.5	0.0	6.1	77.3	15.0	15.3	66.0	18.7	0.0
Missing	0																					
Residence																						
Urban	367	0.1	96.8	2.3	23.7	62.0	14.2	0.0	3.7	100.0	0.0	8.5	69.8	21.7	0.0	5.7	86.8	9.7	34.1	55.6	10.3	0.0
Rural	32	0.1	100.0	0.0	80.1	7.6	12.3	0.0	3.4	85.3	15.5	79.9	20.1	0.0	0.0	1.8	72.4	28.2	0.0	100.0	0.0	0.0
Missing																						
Income_Quintile																						
Q1(poorest)	66	0.1	65.3	14.6	87.7	0.0	12.3	0.0	3.2	92.2	8.2	100.0	0.0	0.0	0.0	5.9	56.5	9.4	100.0	0.0	0.0	0.0
Q2	62	0.1	100.0	0.0	63.9	36.1	0.0	0.0	3.9	100.0	0.0	0.0	100.0	0.0	0.0	6.5	100.0	0.0	36.2	48.6	15.2	0.0
Q3	87	0.1	100.0	0.0	36.6	63.4	0.0	0.0	5.0	100.0	0.0	0.0	71.2	28.8	0.0	11.2	100.0	0.0	20.9	66.2	12.9	0.0
Q4	90	0.2	100.0	0.0	5.5	59.4	35.1	0.0	2.9	100.0	0.0	0.0	35.1	64.9	0.0	2.4	100.0	0.0	0.0	100.0	0.0	0.0
Q5(richest)	92	0.1	100.0	0.0	0.0	88.6	11.4	0.0	3.6	100.0	0.0	0.0	100.0	0.0	0.0	1.3	0.0	0.0				0.0
Missing	3																					
Total	399	0.1	96.9	2.3	27.1	58.8	14.1	0.0	3.7	98.9	1.1	13.1	66.6	20.3	0.0	5.4	86.4	9.4	33.3	56.7	10.0	0.0

**M = % Missing of those Seeking Treatment      SE= Standard Error of % Seeking Treatment**

**Child Danger Signs Indicators:**

**Total = Number of children under the age of 5 years at the time of the interview.**

**"Fast Breathing"/"Vomiting"/"No Drink or Eat" = % of children under the age of 5 with reported episodes of (fast breathing, vomiting, or no eating or drinking) in the last 3 months.**

**"Seeking Treatment" = % of children under 5 with episodes of fast breathing, vomiting, or no eating or drinking in the last 3 months that were taken for care/treatment at either a private health facility, public health facility, or traditional healer (other).**

**Note: large number of missings are usually due to not having information type of facility (either a no answer or a don't know answer).**

## 5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card

	Children 12-23 months with card	DPT 3 Immunization			Measles		
		%Cov	SE	M	%Cov	SE	M
Sex							
Females	9	72.9	17.1	0.0	42.1	21.7	0.0
Males	14	66.8	17.5	0.0	47.7	8.4	0.0
Missing	0						
Residence							
Urban	22	74.4	12.9	0.0	41.2	10.0	0.0
Rural	2	20.8	20.0	0.0	85.2	15.3	0.0
Missing	0						
Income Quintile							
Q1(poorest)	4	41.3	24.2	0.0	50.4	25.4	0.0
Q2	2	0.0	0.0	0.0	37.0	32.9	0.0
Q3	4	51.4	20.8	0.0	33.1	9.1	0.0
Q4	4	77.4	21.7	0.0	41.7	14.4	0.0
Q5(richest)	8	100.0	0.0	0.0	41.5	25.1	0.0
Missing	2						
Total	24	69.2	13.6	0.0	45.5	8.5	0.0

M = % Missing of Covered      SE= Standard Error of % Covered

Vaccinations Meeting WHO Timeline for Children 12 - 23 months Indicators:

Total = Children 12 - 23 months with a vaccination card seen by the interviewer (does not include children less than 12 months to account for censored observations).

"DPT 3" = % of children 12 - 23 months with a vaccination card seen by the interviewer who received 3 inoculations of DTP not sooner than 6, 10 and 14 weeks respectively and before 12 months of age, and that was verified by the interviewer as being recorded on the card.

"Measles" = % of children 12 - 23 months with a vaccination card seen by the interviewer who received who received 1 inoculation of measles not sooner than 8 months of age and before 12 months of age.

## 5.08 Vaccination Status of Children 12-23 Months

	Children 12-23 months	Card seen			DPT 3 Immunization							Measles						
		Seen	SE	M	Total		Card		No Card		M	Total		Card		No Card		M
					%Cov	SE	%Cov	SE	%Cov	SE		%Cov	SE	%Cov	SE			
Sex																		
Females	46	20.5	8.5	0.0	26.8	8.8	59.2	13.8	40.8	13.8	40.6	59.7	10.8	26.6	10.2	73.4	10.2	11.5
Males	51	28.2	6.1	0.0	32.0	5.9	79.6	14.5	20.4	14.5	38.4	58.6	6.0	38.1	8.3	61.9	8.3	22.6
Missing	0																	
Residence																		
Urban	93	23.3	3.3	0.0	28.1	3.8	70.6	9.5	29.4	9.5	41.3	58.8	5.1	30.2	6.9	69.8	6.9	18.2
Rural	5	51.2	23.5	0.0	59.1	16.1	73.8	23.0	26.2	23.0	0.0	66.7	16.7	76.8	19.4	23.2	19.4	0.0
Missing	0																	
Income_Quintile																		
Q1(poorest)	19	18.5	9.6	0.0	18.6	7.6	70.5	28.8	29.5	28.8	34.9	52.6	15.5	17.7	12.2	82.3	12.2	7.4
Q2	17	13.0	8.1	0.0	13.9	9.4	34.6	32.0	65.4	32.0	38.3	45.8	16.0	10.5	10.2	89.5	10.2	31.1
Q3	16	27.8	9.6	0.0	32.3	8.0	64.7	25.4	35.3	25.4	36.9	38.4	8.0	39.8	17.6	60.2	17.6	29.2
Q4	14	28.4	8.0	0.0	39.5	12.4	72.0	23.7	28.0	23.7	56.4	78.9	7.5	36.0	11.0	64.0	11.0	17.0
Q5(richest)	29	26.8	9.1	0.0	35.1	8.1	76.5	10.9	23.5	10.9	38.6	70.0	6.9	38.4	11.4	61.6	11.4	10.7
Missing	2																	
Total	98	24.5	3.3	0.0	29.5	3.6	70.9	8.8	29.1	8.8	39.4	59.1	5.0	32.6	6.3	67.4	6.3	17.3

**M = % Missing (% without adequate information to estimate Total DPT3 or Measles Coverage)**

**SE= Standard Error of % Total, DPT3 or Measles Coverage**

**Explanation of Indicators for Vaccination Status of Children 12 - 23 Months:**

**"Card Seen" = % of children 12 - 23 months of age who had a vaccination card seen by the interviewer.**

**"DPT 3 Immunization": does not account for WHO timeline recommendations.**

**"Total" = % of all children 12 - 23 months of age with DPT3 vaccination either verified by card or by recall of respondent.**

**"Card" = % of all DPT3 vaccinations among children 12-23 months that were verified by card only.**

**"No Card" = % of all DPT3 vaccinations among children 12-23 months that were based on recall of the respondent only.**

**"Measles": does not account for WHO timeline recommendations.**

**"Total" = % of all children 12 - 23 months of age with measles vaccination either verified by card or by recall of respondent.**

**"Card" = % of all Measles vaccinations among children 12-23 months that were verified by card only.**

**"No Card" = % of all Measles vaccinations among children 12-23 months that were based on recall of the respondent only.**



## 5.09 Women's Cancer Screening

	Total Females	Percent Females 18+	PAP Smear	SE	M	Percent Females 40+	Mammography / Breast Exam	SE	M
<b>Residence</b>									
Urban	1,452	83.3	55.4	5.0	1.8	42.4	55.8	5.4	2.3
Rural	111	90.1	52.5	5.4	1.3	48.6	41.4	5.1	0.0
Missing	0								
<b>Income_Quintile</b>									
Q1(poorest)	210	76.8	38.3	2.8	2.4	40.1	34.3	4.2	0.0
Q2	267	80.0	43.0	5.1	2.3	43.3	47.7	7.1	4.2
Q3	317	82.2	50.4	8.2	2.4	37.3	53.1	6.0	0.6
Q4	346	86.2	59.5	3.5	0.8	50.5	53.9	4.1	1.8
Q5(richest)	400	89.9	70.2	1.4	1.3	43.1	71.5	4.1	2.8
Missing	23								
<b>Age</b>									
18-29	385	24.6	48.5	6.8	1.0				
30-39	255	16.3	66.3	3.8	1.8				
40-49	273	17.4	64.3	4.9	2.4	17.4	57.1	6.2	1.8
50-59	207	13.2	58.7	2.7	1.8	13.2	60.5	5.6	2.4
60-69	190	12.2	36.6	7.8	2.4	12.2	44.9	6.1	2.4
70+	254	16.2	0.0	0.0		16.2	0.0	0.0	
Missing	0								
Total	1,563	83.8	55.2	4.7	1.8	42.9	54.7	5.6	2.1

M = % Missing of those with Pap Smear/Mammography SE= Standard Error of % with Pap Smear/Mammography

\*\*\*Note: Values in the age row under the "Females 18 plus" and "Females 40 plus" columns are column percent values.

**Women's Cancer Screening Indicators:**

"Papsmear need" = % of all female respondents aged 18-69

"Papsmear" coverage = % of those female respondents aged 18-69 who self-reported receiving a pap smear test during pelvic examination in the last 3 years.

"Mammography need" = % of all female respondents aged 40-69

"Mammography" coverage = % of those female respondents aged 40-69 who self-reported receiving a mammography test or breast examination in the last 3 years.

\*\*\*\*In several countries women above age 69 were asked the cancer screening questions, and for this reason they are included in the table. For countries that did not ask this questions of women above age 69, these values are expected to be missing.

## 5.10 Oral Health and Injuries

	Oral Health Problems					Road Traffic Accidents				Injury			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
<b>Sex</b>													
Female	1,563	28.0	85.8	1.4	2.2	1.2	45.1	13.0	0.0	3.1	52.3	6.0	7.2
Male	1,414	27.2	80.8	2.4	1.3	1.5	55.2	16.9	0.0	3.3	55.7	6.1	0.0
Missing	0												
<b>Residence</b>													
Urban	2,763	27.8	84.1	1.4	1.9	1.3	48.2	13.3	0.0	3.2	54.5	4.8	4.0
Rural	214	25.5	74.6	5.3	0.0	1.4	80.5	14.3	0.0	3.5	48.3	12.2	0.0
Missing	0												
<b>Income_ Quintile</b>													
Q1(poorest)	417	16.2	66.3	5.3	0.0	2.2	56.0	14.4	0.0	4.1	58.2	9.1	0.0
Q2	528	23.1	78.8	4.0	0.0	1.9	68.3	19.5	0.0	3.6	57.2	12.5	0.0
Q3	601	24.1	79.7	2.8	3.7	1.1	30.2	19.5	0.0	3.5	46.7	6.2	16.5
Q4	651	30.2	86.1	3.4	2.4	1.0	77.2	13.4	0.0	2.2	59.9	4.7	0.0
Q5(richest)	750	37.3	89.9	1.5	1.6	1.1	17.7	11.4	0.0	3.1	49.9	9.7	0.0
Missing	31												
<b>Age</b>													
18-29	793	35.4	82.8	3.2	1.8	1.8	35.3	16.8	0.0	2.8	8.2	6.6	8.2
30-44	811	28.6	84.2	1.6	2.6	1.8	55.5	14.4	0.0	3.2	0.0	8.9	0.0
45-59	636	27.3	83.9	1.8	1.6	0.8	34.4	17.5	0.0	2.9	4.5	10.6	4.5
60-69	337	22.2	79.0	3.9	0.0	1.4	100.0	0.0	0.0	2.7	0.0	7.4	0.0
70-79	268	14.9	82.2	3.7	2.1	0.5	42.5	34.6	0.0	4.3	7.3	4.3	7.3
80+	133	15.5	100.0	0.0	0.0	0.0	0.0	0.0	0.0	5.9	0.0	10.0	0.0
Missing	0												
Total	2,977	27.6	83.5	1.6	1.8	1.3	50.6	12.1		3.2	54.0	4.3	3.7

M = % Missing of % Covered      SE= Standard Error of % Covered

### Indicators:

#### Oral Health:

"Need" = % of all respondents indicating any problems in the last 12 months with their mouth and/or teeth.

"Coverage" = % of respondents reporting problems with their mouth and teeth who indicated receiving dental or medical care in last 12 months.

#### Road Traffic Accident:

"Need" = % of all respondents indicating having suffered bodily injury from a road traffic accident in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury from a road traffic accident in last 12 months who indicating having received treatment for their injuries on-site or at health facility within 1 hour of accident .

#### Injuries from Non-traffic Accidents:

"Need" = % of all respondents indicating having suffered bodily injury due to non-traffic accidents in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury due to non-traffic accidents in last 12 months who indicating having received treatment for injuries on-site or at health facility within 1 hour of accident

## 5.11 Self Reported Non-Communicable Conditions

	Angina					Osteo-Arthritis				Asthma				Diabetes			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
<b>Sex</b>																	
Female	1,563	5.6	40.5	4.6	0.0	14.0	46.5	4.4	0.0	9.7	46.6	3.8	0.0	5.2	74.4	4.8	0.0
Male	1,414	5.7	36.6	5.6	0.0	4.5	33.4	5.7	2.3	7.5	35.0	6.2	0.0	4.9	67.8	5.9	0.0
Missing	0																
<b>Residence</b>																	
Urban	2,763	5.8	39.2	4.5	0.0	9.6	43.6	2.9	0.6	8.8	41.5	2.9	0.0	5.0	69.8	4.8	0.0
Rural	214	3.7	26.8	15.1	0.0	8.8	43.7	7.4	0.0	7.1	46.9	9.5	0.0	5.9	88.8	5.4	0.0
Missing	0																
<b>Income_Quintile</b>																	
Q1(poorest)	417	4.2	34.2	7.8	0.0	13.4	44.8	6.9	0.0	8.0	44.0	7.5	0.0	3.4	72.0	10.2	0.0
Q2	528	7.7	40.3	10.1	0.0	12.9	43.8	5.9	2.2	7.8	40.3	8.5	0.0	5.7	91.5	5.8	0.0
Q3	601	6.0	44.6	9.7	0.0	11.4	47.0	9.7	0.0	8.2	38.4	7.4	0.0	6.4	56.3	9.6	0.0
Q4	651	6.1	37.1	4.1	0.0	7.0	32.1	4.2	0.0	5.7	49.0	6.0	0.0	4.7	65.1	8.5	0.0
Q5(richest)	750	4.4	34.1	7.4	0.0	5.7	46.9	5.5	0.0	12.1	41.3	5.4	0.0	4.8	76.5	6.3	0.0
Missing	31																
<b>Age</b>																	
18-29	793	3.3	12.0	3.8	0.0	1.4	21.7	9.8	0.0	9.2	28.3	4.0	0.0	1.4	24.6	20.9	0.0
30-44	811	4.0	10.4	5.2	0.0	4.0	41.5	9.1	0.0	7.4	45.4	5.2	0.0	2.4	53.5	10.9	0.0
45-59	636	7.0	41.8	12.5	0.0	8.3	47.6	5.4	2.8	8.9	49.4	4.5	0.0	5.1	82.0	6.0	0.0
60-69	337	7.3	58.6	10.9	0.0	21.3	43.7	5.4	0.0	9.9	55.8	10.3	0.0	10.8	88.3	4.5	0.0
70-79	268	9.2	55.0	6.9	0.0	27.1	46.9	5.9	0.0	9.2	31.7	6.1	0.0	13.2	72.5	6.4	0.0
80+	133	11.6	76.7	5.1	0.0	32.0	40.1	14.9	0.0	6.9	55.4	24.3	0.0	12.7	62.2	23.4	0.0
Missing	0																
Total	2,977	5.7	38.6	4.2	0.0	9.5	43.6	2.8	0.5	8.6	41.8	2.8	0.0	5.1	71.4	4.5	0.0

**M = % Missing for Covered**

**SE= Standard Error of %Covered**

**Self Reported Non-Communicable Conditions Indicators:**

"Need" = % of total respondents indicating having received a diagnosis of the condition (angina, arthritis, asthma, diabetes) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (angina, arthritis, asthma, diabetes) indicating they have received treatment or screening for the condition in the last 2 weeks.

## 5.12 Mental Health Conditions

	Total	Depression				Schizophrenia			
		Need	Coverage	SE	M	Need	Coverage	SE	M
Sex									
Female	1,563	14.6	37.1	2.7	0.0	0.8	45.5	9.6	0.0
Male	1,414	6.0	51.3	4.6	0.0	0.7	46.7	20.2	0.0
Missing	0								
Residence									
Urban	2,763	10.6	40.1	2.2	0.0	0.7	47.7	11.5	0.0
Rural	214	9.7	52.4	11.2	0.0	0.6	18.5	19.7	0.0
Missing	0								
Income Quintile									
Q1(poorest)	417	10.9	31.8	5.0	0.0	1.1	10.7	10.5	0.0
Q2	528	12.8	36.9	5.9	0.0	0.5	61.7	31.5	0.0
Q3	601	8.3	55.8	6.3	0.0	0.5	84.5	8.2	0.0
Q4	651	11.3	35.6	4.3	0.0	0.6	45.4	30.5	0.0
Q5(richest)	750	9.5	43.8	3.0	0.0	0.9	46.7	14.0	0.0
Missing	31								
Age									
18-29	793	6.6	42.2	8.9	0.0	0.2	17.1	20.1	0.0
30-44	811	9.1	25.3	4.5	0.0	0.6	76.1	22.8	0.0
45-59	636	13.3	45.4	3.4	0.0	1.4	39.1	15.1	0.0
60-69	337	16.9	44.0	5.8	0.0	1.4	47.9	16.8	0.0
70-79	268	12.9	54.3	9.0	0.0	0.5	0.0	0.0	0.0
80+	133	7.5	47.5	5.5	0.0	0.0	0.0	0.0	0.0
Missing	0								
Total	2,977	10.5	40.9	2.1	0.0	0.7	46.0		0.0

**M = % Missing for Covered**

**SE= Standard Error of %Covered**

**Mental Health Conditions Indicators:**

"Need" = % of total respondents indicating having received a diagnosis of the condition (depression, schizophrenia) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (depression, schizophrenia) indicating they have received treatment or screening for the condition in the last 2 weeks.

### 5.13 Cataracts

		Cataracts			
	Total 60+	Need	Coverage	SE	M
Sex					
Female	444	13.0	44.2	5.9	0.0
Male	293	7.7	45.2	8.3	5.4
Missing	0				
Residence					
Urban	688	11.3	44.5	5.2	1.6
Rural	48	4.5	45.4	16.3	0.0
Missing	0				
Income_Quintile					
Q1(poorest)	142	11.3	48.2	14.1	0.0
Q2	156	8.8	67.4	10.5	0.0
Q3	156	11.5	49.5	15.4	0.0
Q4	158	9.3	22.3	7.2	8.2
Q5(richest)	115	14.3	40.0	6.1	0.0
Missing	10				
Age					
60-69	337	3.6	42.5	18.2	0.0
70-79	268	12.4	47.5	6.3	3.7
80+	133	26.2	42.3	10.0	0.0
Missing	0				
Total	737	10.9	44.5	5.0	1.5

**M = % Missing of Covered    SE= Standard Error of %Covered**

**Cataracts Indicators:**

**Total = Number of respondents 60 years of age or more.**

**"Need" = % in need (respondents 60 years of age or more with self-reported diagnosis of cataracts in the last 5 years).**

**"Coverage" = % respondents 60 years of age or more with self-reported diagnosis of cataracts having indicated receiving eye surgery to remove cataracts in the last 5 years)**

**6.01 In general, rate your health today - q2000**

	Very Good	SE	Good	SE	Moderate	SE	Bad	SE	Very Bad	SE	Missing	SE	N
<b>Gender</b>													
Male	29.6	1.8	55.2	2.0	13.7	0.7	1.3	0.3	0.1	0.0	0.1	0.0	1,414
Female	22.6	2.5	52.6	2.3	21.4	1.2	3.0	0.4	0.4	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	25.9	2.3	53.8	2.0	17.9	1.0	2.1	0.3	0.2	0.1	0.1	0.0	2,763
Rural	26.3	2.9	54.4	2.7	16.0	2.0	2.6	0.8	0.3	0.2	0.3	0.3	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	23.8	3.4	49.7	3.6	21.8	2.2	4.1	0.8	0.6	0.3	0.0	0.0	417
Q2	21.3	3.4	51.9	3.2	22.7	2.8	3.5	1.2	0.6	0.3	0.0	0.0	528
Q3	24.9	3.7	57.7	3.1	15.5	1.2	1.7	0.4	0.2	0.1	0.0	0.0	601
Q4	25.3	3.9	54.8	2.1	18.0	3.2	1.8	0.5	0.0	0.0	0.0	0.0	651
Q5(richest)	32.3	2.4	54.4	2.1	12.6	0.8	0.8	0.2	0.0	0.0	0.0	0.0	750
M	9.5	2.7	36.8	5.2	40.8	5.4	3.5	3.2	0.0	0.0	9.3	2.7	31
<b>Age</b>													
18-29	41.0	2.5	49.3	3.2	7.5	1.4	1.8	0.5	0.1	0.0	0.2	0.1	793
30-44	31.5	3.2	55.0	3.1	12.0	1.2	1.3	0.4	0.1	0.1	0.1	0.1	811
45-59	18.8	2.0	58.0	1.7	19.3	1.8	3.1	0.7	0.7	0.2	0.0	0.0	636
60-69	9.7	1.7	57.8	3.0	28.7	2.1	3.6	1.3	0.1	0.1	0.0	0.0	337
70-79	10.3	1.7	52.8	2.6	34.1	2.7	2.5	0.9	0.3	0.1	0.0	0.0	268
80+	7.5	2.3	46.2	3.9	45.5	4.0	0.8	0.4	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	25.9	2.2	53.9	1.8	17.7	0.9	2.2	0.3	0.2	0.1	0.1	0.0	2,977

## 6.02 Difficulty with 'work and household activities' in last 30 days - q2001

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	84.3	1.0	8.2	0.6	5.9	0.7	1.3	0.4	0.1	0.0	0.1	0.0	1,414
Female	75.5	2.0	11.7	1.0	9.8	0.9	2.7	0.5	0.2	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	80.0	1.1	9.9	0.6	7.8	0.6	2.0	0.4	0.1	0.0	0.1	0.0	2,763
Rural	76.1	2.6	11.8	2.4	9.4	1.2	2.3	0.8	0.2	0.2	0.3	0.3	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	73.5	2.4	10.4	1.7	11.9	1.2	3.9	1.2	0.3	0.1	0.1	0.1	417
Q2	75.3	2.3	11.2	1.3	10.1	1.3	3.0	0.9	0.3	0.2	0.0	0.0	528
Q3	82.3	1.6	8.1	0.9	7.8	1.4	1.8	0.6	0.1	0.1	0.0	0.0	601
Q4	79.9	1.2	11.0	1.2	7.5	1.2	1.4	0.4	0.2	0.2	0.0	0.0	651
Q5(richest)	84.9	1.1	9.7	0.9	4.4	0.7	0.9	0.3	0.0	0.0	0.0	0.0	750
M	58.5	10.4	8.9	5.8	17.8	6.0	7.5	6.9	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	89.7	1.3	6.4	0.8	2.0	0.5	1.6	0.7	0.1	0.0	0.2	0.1	793
30-44	85.0	1.3	9.2	1.0	4.9	0.7	0.8	0.2	0.0	0.0	0.1	0.1	811
45-59	80.0	1.3	8.4	1.1	8.5	0.9	2.7	0.8	0.4	0.1	0.0	0.0	636
60-69	65.2	2.8	16.6	2.6	15.3	1.5	2.5	1.1	0.5	0.5	0.0	0.0	337
70-79	66.2	7.3	14.2	4.8	17.3	2.8	2.1	0.8	0.0	0.0	0.2	0.2	268
80+	50.5	5.5	19.9	3.2	21.9	3.0	7.6	3.8	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	79.7	1.2	10.0	0.6	8.0	0.6	2.0	0.3	0.1	0.0	0.1	0.0	2,977



## 6.03 Difficulty with 'moving around' in last 30 days - q2010

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	91.6	1.1	4.8	0.8	2.8	0.6	0.5	0.2	0.2	0.1	0.1	0.1	1,414
Female	85.3	0.9	7.9	0.8	4.8	0.5	1.9	0.3	0.0	0.0	0.2	0.1	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	88.2	0.7	6.5	0.4	3.9	0.4	1.2	0.2	0.1	0.0	0.1	0.1	2,763
Rural	89.4	1.5	5.8	1.3	2.4	0.7	1.9	0.5	0.0	0.0	0.4	0.3	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	83.5	1.7	6.7	1.2	7.7	1.4	2.0	0.6	0.1	0.1	0.0	0.0	417
Q2	85.4	2.7	7.0	1.1	4.1	1.2	2.8	0.9	0.3	0.3	0.4	0.2	528
Q3	89.2	1.6	6.1	1.2	3.7	0.8	1.0	0.5	0.0	0.0	0.0	0.0	601
Q4	90.1	1.2	5.9	0.9	3.1	0.7	0.8	0.3	0.0	0.0	0.0	0.0	651
Q5(richest)	91.8	0.8	6.0	0.8	1.9	0.4	0.2	0.0	0.2	0.1	0.0	0.0	750
M	60.7	5.9	20.4	4.9	9.0	2.5	2.7	1.2	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	95.6	0.8	3.0	0.7	0.5	0.2	0.7	0.4	0.0	0.0	0.2	0.1	793
30-44	91.5	1.0	5.5	0.5	2.1	0.6	0.6	0.2	0.0	0.0	0.2	0.1	811
45-59	88.6	1.9	5.5	1.0	3.4	0.6	1.8	0.4	0.5	0.2	0.2	0.1	636
60-69	82.8	3.3	9.3	2.2	6.7	1.2	1.0	0.5	0.1	0.1	0.0	0.0	337
70-79	73.9	2.4	14.0	2.1	9.9	1.6	2.2	1.0	0.0	0.0	0.0	0.0	268
80+	65.4	4.3	13.6	2.2	17.1	2.8	4.0	2.2	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	88.3	0.7	6.4	0.4	3.8	0.4	1.2	0.2	0.1	0.0	0.1	0.0	2,977

## 6.04 Difficulty with 'vigorous activities' in last 30 days - q2011

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	53.0	2.8	10.4	1.2	16.0	1.5	11.8	1.6	8.3	0.8	0.5	0.2	1,414
Female	39.5	1.7	11.4	0.6	15.1	1.1	16.4	1.1	17.2	1.1	0.4	0.2	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	46.1	2.2	10.6	0.8	15.3	1.2	14.5	1.0	13.0	0.9	0.5	0.2	2,763
Rural	43.0	4.8	15.2	2.1	18.3	3.4	10.9	2.1	12.2	2.3	0.4	0.4	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	42.6	3.3	8.2	1.8	16.9	1.8	16.2	1.9	16.1	2.2	0.0	0.0	417
Q2	42.6	4.6	10.3	1.4	15.7	1.7	15.9	2.4	15.0	2.5	0.4	0.2	528
Q3	41.0	3.8	9.3	1.2	16.3	3.8	16.4	1.7	16.2	2.2	0.8	0.5	601
Q4	48.3	3.1	10.8	1.0	13.8	1.4	13.9	1.8	12.7	1.7	0.6	0.4	651
Q5(richest)	52.5	1.8	14.1	1.8	15.8	1.2	10.4	0.9	7.1	0.8	0.1	0.1	750
M	30.0	5.2	14.1	5.6	9.5	4.1	15.2	5.9	24.0	7.6	7.3	3.2	31
<b>Age</b>													
18-29	65.0	3.4	15.0	1.7	10.2	2.0	6.6	1.0	2.8	0.7	0.4	0.2	793
30-44	56.5	2.9	10.7	1.0	16.9	1.5	9.8	1.7	5.6	0.8	0.4	0.3	811
45-59	41.8	2.3	11.1	1.1	17.5	1.6	19.3	2.3	9.4	1.3	0.9	0.4	636
60-69	22.3	1.9	9.9	2.6	23.7	3.0	21.8	2.2	22.2	1.8	0.0	0.0	337
70-79	17.1	1.6	4.5	1.0	16.5	2.9	24.9	2.6	36.7	1.9	0.3	0.1	268
80+	4.1	3.6	2.2	0.8	7.0	2.0	21.9	2.6	64.8	4.8	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	45.9	2.1	10.9	0.7	15.5	1.1	14.2	0.9	13.0	0.9	0.5	0.1	2,977

## 6.05 Difficulty with 'self-care' in last 30 days - q2020

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	97.6	0.4	1.1	0.2	0.8	0.3	0.3	0.1	0.0	0.0	0.2	0.1	1,414
Female	96.7	0.5	1.7	0.2	1.2	0.4	0.2	0.1	0.0	0.0	0.2	0.1	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	97.2	0.3	1.4	0.2	1.0	0.2	0.2	0.1	0.0	0.0	0.2	0.1	2,763
Rural	96.6	1.1	1.9	0.9	1.3	0.6	0.2	0.2	0.0	0.0	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	94.3	1.2	3.2	1.0	2.0	0.7	0.5	0.2	0.0	0.0	0.0	0.0	417
Q2	96.7	0.9	1.9	0.6	0.8	0.4	0.3	0.3	0.0	0.0	0.2	0.2	528
Q3	96.8	0.6	1.2	0.3	1.5	0.7	0.2	0.1	0.0	0.0	0.3	0.2	601
Q4	97.8	0.6	0.9	0.3	1.1	0.4	0.1	0.2	0.0	0.0	0.1	0.0	651
Q5(richest)	99.1	0.3	0.5	0.4	0.2	0.1	0.2	0.1	0.0	0.0	0.0	0.0	750
M	86.6	5.8	6.1	2.7	0.0	0.0	0.0	0.0	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	99.0	0.4	0.4	0.2	0.5	0.3	0.0	0.0	0.0	0.0	0.2	0.1	793
30-44	98.7	0.3	0.4	0.2	0.6	0.2	0.0	0.0	0.0	0.0	0.3	0.1	811
45-59	97.1	0.7	0.9	0.2	1.1	0.4	0.7	0.3	0.0	0.0	0.2	0.1	636
60-69	96.2	0.7	2.1	0.7	1.5	0.4	0.2	0.2	0.0	0.0	0.0	0.0	337
70-79	91.6	1.3	5.3	1.0	2.3	1.2	0.4	0.1	0.0	0.0	0.3	0.1	268
80+	90.3	3.2	7.2	2.7	1.8	1.4	0.8	0.4	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	97.1	0.3	1.4	0.2	1.0	0.2	0.2	0.1	0.2	0.1			2,977

**6.06      Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	97.4	0.4	1.2	0.2	0.9	0.3	0.1	0.1	0.2	0.2	0.1	0.0	1,414
Female	96.8	0.6	2.2	0.4	0.8	0.4	0.1	0.0	0.0	0.0	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	97.1	0.3	1.7	0.3	0.8	0.2	0.1	0.1	0.1	0.1	0.1	0.0	2,763
Rural	96.7	1.1	1.8	0.8	1.4	0.7	0.2	0.2	0.0	0.0	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	94.3	1.2	3.5	1.0	1.7	0.6	0.6	0.3	0.0	0.0	0.0	0.0	417
Q2	96.5	0.8	1.9	0.7	0.7	0.3	0.3	0.3	0.6	0.6	0.0	0.0	528
Q3	97.5	0.6	1.3	0.4	1.2	0.8	0.0	0.0	0.0	0.0	0.0	0.0	601
Q4	97.5	0.6	1.5	0.4	0.9	0.4	0.0	0.0	0.0	0.0	0.2	0.1	651
Q5(richest)	98.7	0.4	1.1	0.3	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0	750
M	86.6	5.8	6.1	2.7	0.0	0.0	0.0	0.0	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	98.3	0.6	1.2	0.6	0.4	0.3	0.0	0.0	0.0	0.0	0.2	0.1	793
30-44	99.0	0.3	0.4	0.1	0.4	0.2	0.0	0.0	0.0	0.0	0.2	0.1	811
45-59	96.9	0.7	1.6	0.5	1.1	0.4	0.3	0.2	0.0	0.0	0.1	0.1	636
60-69	95.7	0.6	2.4	0.6	1.7	0.4	0.2	0.2	0.0	0.0	0.0	0.0	337
70-79	93.9	1.9	4.1	1.2	1.8	1.3	0.1	0.1	0.0	0.0	0.0	0.0	268
80+	88.8	4.7	7.0	3.2	1.2	1.1	0.8	0.4	2.3	2.5	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	97.1	0.3	1.7	0.3	0.8	0.2	0.1	0.1	0.1	0.1	0.1	0.0	2,977

**6.07 Amount of 'bodily aches or pains' in last 30 days - q2030**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	68.7	1.6	20.7	1.1	8.1	0.9	2.4	0.4	0.1	0.0	0.1	0.0	1,414
Female	58.8	1.8	20.7	1.2	15.7	0.9	4.4	0.7	0.3	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	63.7	1.5	20.3	0.9	12.3	0.7	3.4	0.5	0.2	0.1	0.1	0.0	2,763
Rural	60.8	2.3	26.2	1.8	9.3	1.4	3.7	0.8	0.0	0.0	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	61.2	3.1	17.8	1.7	14.6	1.8	6.2	1.4	0.2	0.2	0.1	0.1	417
Q2	58.8	3.0	20.9	1.9	15.2	1.9	4.6	1.0	0.4	0.3	0.0	0.0	528
Q3	65.8	2.4	19.6	2.4	11.3	1.0	3.2	0.7	0.0	0.0	0.0	0.0	601
Q4	66.1	1.8	21.2	1.7	10.3	1.8	2.3	0.9	0.1	0.1	0.0	0.0	651
Q5(richest)	64.6	1.4	22.7	0.7	10.3	1.0	2.0	0.4	0.3	0.1	0.0	0.0	750
M	44.2	5.4	20.0	6.8	20.3	6.4	8.2	7.7	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	70.2	2.8	20.8	1.9	6.4	0.8	2.4	0.5	0.0	0.0	0.2	0.1	793
30-44	69.2	2.1	19.6	1.6	8.2	1.0	2.6	0.8	0.3	0.2	0.1	0.1	811
45-59	61.8	2.2	20.3	1.6	13.9	1.5	3.5	0.7	0.5	0.2	0.0	0.0	636
60-69	53.1	3.1	24.1	2.1	17.7	2.3	4.9	2.2	0.0	0.0	0.2	0.1	337
70-79	51.7	6.0	21.8	5.8	22.0	2.0	4.2	1.2	0.3	0.3	0.0	0.0	268
80+	45.6	10.7	18.4	6.7	26.4	3.5	9.5	3.8	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	63.5	1.4	20.7	0.9	12.1	0.6	3.4	0.4	0.2	0.1	0.1	0.0	2,977

**6.08 Amount of 'bodily discomfort' in last 30 days - q2031**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	71.8	1.2	18.6	0.9	7.7	0.9	1.7	0.3	0.1	0.0	0.1	0.0	1,414
Female	62.5	1.5	19.7	1.4	13.7	0.8	3.7	0.4	0.3	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	67.0	1.1	18.8	0.8	11.2	0.6	2.7	0.3	0.2	0.1	0.1	0.0	2,763
Rural	66.2	3.0	24.2	2.7	6.6	1.3	3.0	0.9	0.0	0.0	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	62.9	3.2	18.7	2.3	13.5	2.1	4.5	1.0	0.4	0.2	0.0	0.0	417
Q2	64.3	3.3	18.5	2.6	12.7	2.0	4.2	1.0	0.3	0.3	0.0	0.0	528
Q3	69.6	1.9	18.0	2.7	9.9	1.4	2.5	0.5	0.0	0.0	0.0	0.0	601
Q4	68.5	1.3	18.2	1.3	11.7	1.7	1.5	0.5	0.1	0.1	0.0	0.0	651
Q5(richest)	68.6	0.9	21.2	1.3	7.8	0.9	2.0	0.5	0.3	0.1	0.0	0.0	750
M	42.1	7.2	32.1	6.6	16.4	6.2	2.1	2.4	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	73.9	2.1	18.9	1.7	4.6	0.9	2.5	0.8	0.0	0.0	0.2	0.1	793
30-44	72.2	1.7	18.6	1.4	6.6	1.2	2.1	0.4	0.3	0.2	0.1	0.1	811
45-59	65.6	1.8	18.9	1.1	12.5	1.2	2.7	0.6	0.3	0.2	0.0	0.0	636
60-69	56.5	2.9	22.3	1.8	19.1	2.2	2.2	1.0	0.0	0.0	0.0	0.0	337
70-79	54.8	4.7	21.3	5.4	20.3	2.1	3.3	0.7	0.3	0.3	0.0	0.0	268
80+	50.9	6.2	14.1	2.3	25.7	3.9	8.5	3.4	0.8	0.4	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	66.9	1.0	19.2	0.8	10.8	0.6	2.7	0.3	0.2	0.1	0.1	0.0	2,977

**6.09      Difficulty with 'concentrating or remembering things' in last 30 days - q2050**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	82.5	1.0	11.5	0.6	4.7	0.8	0.7	0.2	0.4	0.2	0.2	0.1	1,414
Female	72.2	1.6	17.2	1.1	7.8	0.8	2.6	0.4	0.1	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	76.6	1.2	14.7	0.8	6.5	0.5	1.7	0.3	0.3	0.1	0.2	0.0	2,763
Rural	82.7	2.5	11.8	2.1	4.4	1.1	1.1	0.6	0.0	0.0	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	74.7	1.7	12.6	1.6	9.2	1.3	2.8	0.8	0.5	0.3	0.2	0.2	417
Q2	76.1	3.4	15.3	2.7	7.6	1.3	0.9	0.5	0.0	0.0	0.0	0.0	528
Q3	78.6	2.3	13.9	1.1	5.6	1.2	1.7	0.6	0.1	0.2	0.0	0.0	601
Q4	78.0	1.6	14.6	1.0	5.0	0.9	2.1	0.5	0.0	0.0	0.3	0.1	651
Q5(richest)	77.6	1.1	15.0	1.1	5.4	0.7	1.3	0.3	0.6	0.5	0.0	0.0	750
M	61.8	6.0	23.6	5.5	7.4	5.5	0.0	0.0	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	86.2	2.4	10.9	1.8	2.2	0.5	0.6	0.3	0.0	0.0	0.2	0.1	793
30-44	85.6	1.3	9.8	1.1	2.6	0.3	1.4	0.4	0.1	0.1	0.5	0.2	811
45-59	73.7	2.7	16.4	1.3	7.0	1.3	2.2	0.7	0.7	0.5	0.0	0.0	636
60-69	70.1	3.6	16.6	2.2	9.3	1.7	3.5	1.7	0.5	0.5	0.0	0.0	337
70-79	59.7	4.9	22.5	2.2	16.6	3.6	1.1	0.3	0.0	0.0	0.0	0.0	268
80+	39.0	4.8	34.3	3.6	22.0	3.2	4.7	1.4	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	77.1	1.3	14.5	0.8	6.3	0.5	1.7	0.3	0.2	0.1	0.2	0.0	2,977

## 6.10 Difficulty with 'learning a new task' in last 30 days - q2051

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	90.0	0.8	6.3	0.5	2.6	0.6	0.9	0.4	0.1	0.1	0.1	0.0	1,414
Female	85.2	0.8	8.3	0.6	5.0	0.5	1.4	0.4	0.0	0.0	0.1	0.1	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	87.2	0.7	7.4	0.5	4.0	0.4	1.2	0.3	0.0	0.1	0.1	0.1	2,763
Rural	91.1	1.5	5.7	1.4	2.4	0.8	0.8	0.3	0.0	0.0	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	81.9	1.9	8.3	1.4	6.0	1.4	3.7	1.3	0.0	0.0	0.0	0.0	417
Q2	84.2	3.3	9.1	2.1	5.4	1.1	1.0	0.4	0.0	0.0	0.2	0.2	528
Q3	88.1	1.3	7.7	0.9	3.0	0.7	1.1	0.4	0.2	0.2	0.0	0.0	601
Q4	88.1	1.0	7.9	0.8	3.5	0.7	0.6	0.5	0.0	0.0	0.0	0.0	651
Q5(richest)	92.2	0.5	4.6	0.4	2.6	0.6	0.6	0.3	0.0	0.0	0.0	0.0	750
M	76.7	5.5	10.6	3.7	5.4	6.0	0.0	0.0	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	95.5	0.8	3.4	0.8	0.6	0.5	0.3	0.2	0.0	0.0	0.2	0.1	793
30-44	94.7	0.4	3.9	0.4	0.8	0.2	0.5	0.2	0.0	0.0	0.1	0.1	811
45-59	87.8	1.9	7.5	0.9	3.6	1.1	0.9	0.3	0.0	0.0	0.2	0.1	636
60-69	80.6	3.2	10.3	2.0	6.1	1.7	2.7	1.4	0.3	0.4	0.0	0.0	337
70-79	71.0	3.8	16.2	1.9	9.3	2.0	3.6	1.4	0.0	0.0	0.0	0.0	268
80+	44.3	3.5	25.3	3.4	27.0	3.0	3.4	1.8	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	87.5	0.7	7.3	0.5	3.9	0.4	1.2	0.3	0.0	0.0	0.1	0.1	2,977



**6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	93.2	0.7	4.6	0.6	1.6	0.4	0.4	0.1	0.0	0.0	0.2	0.1	1,414
Female	93.9	0.6	3.5	0.6	1.8	0.3	0.5	0.1	0.2	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	93.5	0.7	4.0	0.6	1.8	0.3	0.5	0.1	0.1	0.0	0.1	0.1	2,763
Rural	94.8	1.5	4.1	1.3	0.5	0.4	0.4	0.2	0.2	0.2	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	90.7	1.6	5.5	1.2	2.4	1.0	1.2	0.5	0.1	0.1	0.0	0.0	417
Q2	92.6	1.4	5.0	1.1	1.9	0.7	0.5	0.3	0.0	0.0	0.0	0.0	528
Q3	93.5	1.0	3.6	0.7	2.3	0.7	0.5	0.1	0.0	0.0	0.2	0.1	601
Q4	95.0	0.7	3.4	0.8	1.5	0.4	0.1	0.2	0.0	0.0	0.0	0.0	651
Q5(richest)	95.0	1.1	3.4	1.0	1.0	0.6	0.4	0.1	0.3	0.1	0.0	0.0	750
M	86.9	4.7	5.8	1.7	0.0	0.0	0.0	0.0	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	93.8	1.2	3.7	1.0	1.7	0.6	0.3	0.1	0.3	0.2	0.2	0.1	793
30-44	96.0	0.4	2.5	0.6	0.7	0.2	0.5	0.2	0.0	0.0	0.3	0.2	811
45-59	93.1	1.2	4.8	1.3	1.3	0.3	0.8	0.3	0.0	0.0	0.0	0.0	636
60-69	91.9	2.5	4.3	1.3	3.3	1.7	0.5	0.5	0.0	0.0	0.0	0.0	337
70-79	89.5	2.6	7.0	1.8	3.1	1.1	0.4	0.4	0.0	0.0	0.0	0.0	268
80+	92.5	2.4	4.9	2.5	2.4	1.2	0.0	0.0	0.3	0.3	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	93.6	0.6	4.0	0.6	1.7	0.2	0.5	0.1	0.1	0.0	0.1	0.1	2,977

## 6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	86.6	1.3	8.7	0.8	3.3	0.6	1.2	0.3	0.1	0.1	0.1	0.0	1,414
Female	83.0	1.6	10.6	1.3	5.1	0.5	1.2	0.2	0.1	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	84.5	1.4	9.8	1.0	4.3	0.4	1.2	0.1	0.1	0.1	0.1	0.0	2,763
Rural	87.4	2.3	8.4	2.4	3.8	1.2	0.4	0.2	0.0	0.0	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	85.2	1.9	8.5	1.4	4.3	1.0	1.8	0.6	0.2	0.2	0.0	0.0	417
Q2	85.4	1.9	8.4	1.3	5.0	0.8	0.9	0.3	0.2	0.1	0.0	0.0	528
Q3	82.9	2.2	10.3	1.3	5.5	1.3	1.0	0.2	0.2	0.1	0.0	0.0	601
Q4	86.4	1.8	10.1	1.7	2.8	0.5	0.7	0.2	0.0	0.0	0.0	0.0	651
Q5(richest)	84.0	1.2	10.4	1.0	4.1	0.4	1.6	0.6	0.0	0.0	0.0	0.0	750
M	81.1	6.3	10.5	3.8	1.1	1.2	0.0	0.0	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	84.2	2.0	9.0	1.3	5.1	0.7	1.6	0.4	0.0	0.0	0.2	0.1	793
30-44	86.1	2.5	9.5	1.9	3.7	0.8	0.5	0.2	0.1	0.1	0.1	0.1	811
45-59	83.0	2.2	11.0	1.4	4.2	1.0	1.4	0.4	0.4	0.2	0.0	0.0	636
60-69	83.6	1.9	10.7	1.9	4.1	1.7	1.7	0.7	0.0	0.0	0.0	0.0	337
70-79	85.0	1.8	10.6	1.2	3.9	1.3	0.6	0.3	0.0	0.0	0.0	0.0	268
80+	89.8	2.6	4.6	2.8	4.1	1.4	1.5	0.7	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	84.7	1.3	9.7	0.9	4.3	0.4	1.2	0.1	0.1	0.1	0.1	0.0	2,977

## 6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	86.0	1.8	6.4	0.8	5.1	1.0	1.8	0.5	0.4	0.2	0.3	0.2	1,414
Female	83.6	1.5	8.9	1.0	4.3	0.8	2.8	0.3	0.3	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	84.9	1.6	7.6	0.8	4.6	0.8	2.4	0.2	0.3	0.1	0.2	0.1	2,763
Rural	82.8	2.7	8.9	1.4	6.5	2.5	1.7	0.5	0.2	0.2	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	81.2	2.6	9.5	1.4	5.5	1.3	3.0	1.3	0.9	0.6	0.0	0.0	417
Q2	82.0	1.5	6.8	0.9	6.5	1.5	4.2	0.8	0.4	0.3	0.0	0.0	528
Q3	84.8	2.6	8.1	1.4	4.9	2.0	1.9	0.5	0.3	0.0	0.0	0.0	601
Q4	86.5	1.2	7.1	0.9	4.0	0.7	2.2	0.5	0.2	0.1	0.1	0.0	651
Q5(richest)	87.8	1.3	7.2	1.0	3.6	0.4	0.9	0.4	0.1	0.0	0.4	0.4	750
M	66.5	7.7	16.5	5.5	4.3	1.9	5.4	2.4	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	90.8	1.9	4.5	1.3	3.6	0.8	0.3	0.1	0.4	0.3	0.4	0.2	793
30-44	92.0	1.4	4.5	0.9	2.0	0.5	1.2	0.4	0.1	0.0	0.3	0.2	811
45-59	82.3	2.0	9.0	1.1	5.2	1.6	3.1	0.7	0.4	0.2	0.0	0.0	636
60-69	79.5	4.7	9.9	2.2	7.5	2.5	2.8	0.9	0.3	0.1	0.0	0.0	337
70-79	70.8	4.0	16.4	2.2	6.4	2.2	5.8	1.7	0.6	0.2	0.0	0.0	268
80+	57.7	3.0	18.0	2.9	14.8	2.8	8.6	1.9	0.8	0.4	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	84.7	1.5	7.7	0.8	4.7	0.8	2.3	0.2	0.3	0.1	0.2	0.1	2,977

## 6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	85.7	3.0	6.6	1.2	4.6	1.2	2.6	0.9	0.4	0.2	0.2	0.1	1,414
Female	81.1	3.4	9.5	1.3	6.1	1.7	3.1	0.8	0.2	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	83.8	3.0	7.9	1.1	5.2	1.4	2.7	0.8	0.3	0.1	0.2	0.1	2,763
Rural	76.4	3.7	11.3	1.5	7.9	2.7	4.3	1.4	0.2	0.2	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	78.4	4.2	8.7	2.5	6.6	1.3	5.2	1.5	1.1	0.7	0.0	0.0	417
Q2	77.8	3.5	8.1	1.3	7.5	1.6	6.4	1.7	0.2	0.1	0.0	0.0	528
Q3	83.1	3.2	8.0	1.4	6.3	2.2	2.4	0.6	0.1	0.1	0.0	0.0	601
Q4	85.1	2.4	7.9	1.1	5.6	1.6	1.2	0.3	0.2	0.1	0.1	0.0	651
Q5(richest)	89.2	1.6	7.7	1.0	2.2	0.5	0.8	0.4	0.0	0.0	0.2	0.2	750
M	63.5	6.5	15.7	5.8	8.1	5.4	2.7	1.2	2.7	1.2	7.3	3.2	31
<b>Age</b>													
18-29	92.4	2.0	5.0	0.9	1.8	1.2	0.3	0.2	0.3	0.3	0.2	0.1	793
30-44	92.9	1.6	3.8	0.5	2.4	1.1	0.5	0.3	0.0	0.0	0.3	0.2	811
45-59	70.8	4.7	14.4	2.2	10.2	2.2	4.5	1.2	0.2	0.1	0.0	0.0	636
60-69	74.5	8.3	8.9	2.4	9.2	3.6	6.7	2.6	0.6	0.7	0.0	0.0	337
70-79	76.4	4.6	10.6	2.2	7.4	1.6	4.7	2.4	0.9	0.3	0.0	0.0	268
80+	64.9	6.1	16.0	3.1	8.1	2.5	11.0	2.6	0.0	0.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	83.3	3.1	8.1	1.1	5.4	1.4	2.8	0.8	0.3	0.1	0.1	0.1	2,977

## 6.15 Problem with 'sleeping' in last 30 days - q2080

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	75.8	1.5	14.6	1.2	6.5	0.8	2.8	0.5	0.2	0.1	0.1	0.0	1,414
Female	67.8	1.5	17.1	1.0	9.5	1.0	5.0	0.4	0.6	0.2	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	71.6	1.5	15.7	1.0	8.2	0.7	4.1	0.3	0.4	0.2	0.1	0.0	2,763
Rural	71.8	4.0	19.3	4.0	6.5	1.0	2.4	0.8	0.1	0.1	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	73.5	1.7	12.2	1.4	8.1	1.1	5.8	0.9	0.4	0.3	0.0	0.0	417
Q2	70.6	3.5	14.7	2.0	10.6	1.5	3.8	0.9	0.3	0.1	0.0	0.0	528
Q3	71.9	3.4	17.7	3.4	6.4	1.1	3.6	0.8	0.4	0.2	0.0	0.0	601
Q4	72.6	1.8	15.2	1.5	7.5	0.7	4.1	0.8	0.6	0.2	0.0	0.0	651
Q5(richest)	71.4	1.4	17.8	0.8	7.9	1.2	2.6	0.4	0.3	0.1	0.0	0.0	750
M	37.9	10.3	24.8	7.0	13.1	3.4	17.0	5.0	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	75.1	2.8	16.2	1.8	6.5	0.9	1.6	0.4	0.5	0.2	0.2	0.1	793
30-44	75.7	2.2	15.6	2.3	5.8	0.9	2.4	0.4	0.5	0.2	0.1	0.1	811
45-59	69.6	1.5	14.3	1.3	9.9	1.1	5.9	1.1	0.3	0.1	0.0	0.0	636
60-69	66.9	2.0	15.4	1.9	10.1	2.1	7.2	1.2	0.3	0.2	0.0	0.0	337
70-79	66.1	3.4	16.4	1.9	10.6	1.6	6.8	1.3	0.0	0.0	0.0	0.0	268
80+	57.5	5.0	24.6	3.8	12.8	2.4	4.3	2.9	0.8	0.4	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	71.6	1.5	15.9	1.0	8.1	0.7	4.0	0.3	0.4	0.2	0.1	0.0	2,977

**6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	71.0	1.4	20.1	1.2	6.6	0.5	2.1	0.4	0.0	0.0	0.1	0.0	1,414
Female	58.9	2.9	23.3	1.6	13.2	1.4	4.0	0.2	0.5	0.2	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	64.7	2.4	21.6	1.4	10.0	1.0	3.2	0.2	0.3	0.1	0.1	0.0	2,763
Rural	63.5	4.0	23.7	3.3	10.5	1.8	2.3	0.8	0.0	0.0	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	65.5	2.4	18.9	1.8	10.8	1.6	4.4	0.8	0.4	0.2	0.0	0.0	417
Q2	67.4	3.1	18.7	1.9	10.2	1.4	3.0	0.9	0.7	0.4	0.0	0.0	528
Q3	66.1	2.8	21.7	2.9	10.1	1.8	2.1	0.5	0.0	0.0	0.0	0.0	601
Q4	64.2	2.1	21.8	2.0	10.2	0.9	3.8	0.8	0.0	0.0	0.0	0.0	651
Q5(richest)	62.1	3.6	25.7	3.2	9.3	0.9	2.6	0.9	0.3	0.1	0.0	0.0	750
M	46.8	12.3	22.6	8.8	12.9	3.3	9.0	5.2	1.4	1.7	7.3	3.2	31
<b>Age</b>													
18-29	66.7	4.7	24.4	4.1	6.0	0.7	2.9	0.5	0.0	0.0	0.2	0.1	793
30-44	64.5	3.0	23.1	2.3	9.3	1.3	2.5	0.6	0.4	0.2	0.1	0.1	811
45-59	63.4	3.2	21.3	1.6	11.5	1.6	3.5	1.0	0.3	0.1	0.0	0.0	636
60-69	64.3	2.3	18.2	2.1	12.5	1.4	4.6	1.3	0.3	0.2	0.0	0.0	337
70-79	64.2	3.2	19.0	3.1	13.1	1.4	3.5	1.0	0.3	0.3	0.0	0.0	268
80+	61.0	5.7	15.6	4.1	19.9	2.7	2.7	2.2	0.8	0.4	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	64.6	2.2	21.8	1.2	10.1	1.0	3.1	0.2	0.3	0.1	0.1	0.0	2,977

## 6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	66.0	1.5	22.3	1.5	8.9	1.1	2.1	0.4	0.6	0.2	0.1	0.0	1,414
Female	51.4	2.2	27.4	2.3	14.7	1.2	6.0	0.8	0.5	0.1	0.1	0.0	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	58.4	1.9	24.6	1.5	12.1	0.6	4.2	0.5	0.5	0.1	0.1	0.0	2,763
Rural	56.9	3.2	29.3	2.2	9.9	1.7	3.6	0.7	0.4	0.2	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	55.5	2.7	25.4	2.2	12.6	1.6	5.9	0.9	0.6	0.3	0.0	0.0	417
Q2	53.8	3.3	26.6	2.0	13.4	1.5	6.1	1.7	0.2	0.1	0.0	0.0	528
Q3	60.8	3.0	23.8	4.0	11.4	1.5	3.3	0.6	0.8	0.4	0.0	0.0	601
Q4	61.8	1.7	23.8	1.6	10.6	0.8	3.5	0.5	0.3	0.2	0.0	0.0	651
Q5(richest)	59.1	2.8	25.9	1.7	11.6	1.0	2.7	0.6	0.7	0.3	0.0	0.0	750
M	33.2	7.9	18.8	10.4	28.1	9.2	12.6	3.4	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	62.7	3.3	24.1	2.8	8.7	1.0	3.6	0.6	0.8	0.3	0.2	0.1	793
30-44	62.5	1.8	24.5	1.7	10.6	1.0	1.9	0.2	0.4	0.1	0.1	0.1	811
45-59	54.7	3.1	25.7	2.0	13.3	1.2	5.6	1.2	0.7	0.2	0.0	0.0	636
60-69	52.2	2.9	24.9	3.0	15.8	1.3	7.0	1.2	0.1	0.1	0.0	0.0	337
70-79	51.2	4.8	28.2	3.7	15.9	2.5	4.4	0.8	0.3	0.3	0.0	0.0	268
80+	53.5	3.5	23.1	3.8	16.2	2.7	6.4	2.4	0.8	0.4	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	58.3	1.8	25.0	1.4	12.0	0.6	4.2	0.4	0.5	0.1	0.1	0.0	2,977

**6.18 Problem with 'worry or anxiety' in last 30 days - q2091**

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
<b>Gender</b>													
Male	47.9	2.1	24.2	1.3	18.1	1.0	8.7	2.0	1.0	0.3	0.1	0.0	1,414
Female	40.0	2.2	27.1	1.9	22.0	1.8	10.1	1.0	0.5	0.1	0.2	0.1	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	43.6	1.8	25.9	1.5	20.0	1.3	9.5	1.2	0.8	0.2	0.2	0.1	2,763
Rural	45.5	3.6	23.4	2.6	22.2	3.2	8.7	2.5	0.2	0.2	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	46.7	2.7	22.2	1.7	20.2	2.0	9.5	2.0	1.5	0.4	0.0	0.0	417
Q2	46.1	3.5	21.6	2.1	18.6	2.3	13.6	2.0	0.1	0.2	0.0	0.0	528
Q3	42.9	4.6	27.4	1.8	19.4	3.1	9.6	2.6	0.7	0.4	0.0	0.0	601
Q4	46.9	2.3	23.9	1.9	21.3	2.3	7.4	0.8	0.5	0.2	0.0	0.0	651
Q5(richest)	39.0	2.0	31.0	2.0	20.6	0.9	8.1	0.9	1.0	0.8	0.3	0.1	750
M	31.5	9.8	27.0	10.7	23.8	10.9	10.5	3.7	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	46.1	3.5	26.4	1.8	18.4	2.4	7.5	1.0	1.1	0.5	0.5	0.2	793
30-44	44.7	2.2	26.3	3.6	20.2	1.7	8.1	2.3	0.6	0.2	0.1	0.1	811
45-59	35.9	2.6	28.1	2.9	21.0	1.5	14.2	1.8	0.9	0.2	0.0	0.0	636
60-69	42.6	4.4	21.8	3.2	25.0	2.2	10.4	2.5	0.2	0.1	0.0	0.0	337
70-79	49.8	4.2	24.3	3.2	16.1	2.2	9.5	0.8	0.3	0.3	0.0	0.0	268
80+	52.6	3.5	20.1	2.6	21.6	5.5	4.0	2.0	1.6	0.8	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	43.8	1.7	25.8	1.5	20.1	1.2	9.4	1.2	0.7	0.2	0.2	0.1	2,977



## 7.1 Self-reported utilization of health services and perceived unmet need

	<u>Ambulatory Adult</u>		<u>Ambulatory Child</u>		<u>Inpatient Adult</u>		<u>Inpatient Child</u>		<u>Informal Care</u>		<u>Unmet Care</u>		<u>Non use</u>	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>														
Female	46.1	2.2	8.7	1.4	3.0	0.4	0.6	0.2	5.8	1.4	0.2	0.1	35.7	2.6
Male	34.6	3.3	5.0	1.0	3.7	0.5	0.9	0.4	4.2	1.2	0.2	0.2	51.5	3.4
<b>Area of Residence</b>														
Urban	42.1	2.6	6.7	1.1	2.9	0.3	0.7	0.2	5.8	1.3	0.1	0.0	41.6	3.2
Rural	35.9	3.8	7.8	1.6	4.6	1.0	0.8	0.4	2.3	0.9	0.7	0.6	48.0	4.5
<b>Age Group</b>														
18-29y	36.8	3.9	8.1	2.1	2.0	0.7	0.8	0.3	6.8	2.0	0.1	0.1	45.4	3.4
30-44y	36.2	3.4	14.4	1.7	1.6	0.4	1.5	0.4	5.4	1.0	0.5	0.6	40.3	3.8
45-59y	39.4	2.2	3.1	0.6	5.3	1.2	0.4	0.2	5.4	2.2	0.1	0.1	46.3	3.4
60-69y	48.7	4.2	0.0	0.0	5.7	1.3	0.0	0.0	0.3	0.2	0.0	0.0	45.3	5.0
70+y	59.3	2.6	0.4	0.4	3.0	0.6	0.0	0.0	4.3	1.6	0.0	0.0	33.0	3.0
<b>Education Years</b>														
0	67.2	9.1	0.0	0.0	3.5	1.8	0.0	0.0	0.0	0.0	0.0	0.0	29.3	9.2
1-5	43.5	3.9	2.1	1.2	5.5	1.7	0.1	0.2	2.3	1.8	0.0	0.0	46.5	5.3
6-11	39.3	2.8	7.9	1.6	3.6	0.6	0.8	0.2	3.6	1.1	0.0	0.0	44.8	2.9
12+	40.9	3.0	7.0	0.9	2.5	0.5	0.8	0.2	7.4	1.2	0.5	0.4	40.8	3.2
<b>Income Quintiles</b>														
Q1	32.8	3.4	8.4	2.3	4.6	1.2	1.4	0.8	2.1	0.9	0.0	0.0	50.7	4.1
Q2	41.4	3.7	5.6	1.2	3.2	0.8	1.0	0.6	3.7	1.4	0.1	0.1	44.9	3.9
Q3	37.7	2.5	8.9	2.0	2.8	0.6	0.3	0.1	5.2	1.9	0.9	0.6	44.3	3.1
Q4	42.8	3.1	6.5	1.7	3.1	0.7	0.3	0.1	4.9	1.2	0.1	0.0	42.2	3.0
Q5	46.4	2.5	5.2	0.5	3.2	0.4	0.6	0.4	8.4	0.7	0.0	0.0	36.2	2.8
<b>Ownership</b>														
Government	57.1	2.6	13.8	1.9	8.5	1.7	2.3	0.6	3.4	1.0	0.0	0.0	14.9	2.1
Private	63.5	1.7	9.6	1.9	7.1	0.5	1.0	0.2	9.6	1.5	0.1	0.2	9.0	1.1
NGO														
Other	36.6	9.5	13.4	4.4	0.0	0.0	0.0	0.0	27.1	6.3	0.0	0.0	22.9	16.8
TOTAL	40.7	2.6	6.9	1.0	3.3	0.3	0.7	0.1	5.0	1.3	0.2	0.2	43.1	3.1

This table presents the percent of people reporting utilization of health services when they last needed care. It combines information from Q7000, Q7001 and Q7016. Utilization was recorded in the WHS in three separate categories: ambulatory care, hospital inpatient care, and informal care. Time cut-offs for utilization recall used here are 1 year for ambulatory care and informal care, and 3 years for hospital inpatient care. Utilization of ambulatory and inpatient care services

have been further sub-divided into "adult" and "child" categories. In addition to utilization, two other related aspects are being reported here - unmet need and non-use (regardless of expressed need) of health services. Unmet need was estimated on the basis of responses to Q7000 and Q7004. Non-use of health services is simply the percent of respondents who reported no need in Q7000 or did not utilize services in the time cut-offs mentioned above.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, and ownership (of health facility). Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership was constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services.

## 7.2 Patient assessed characteristics of ambulatory health services

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
<b>Sex</b>																
Female	82.1	3.3	98.5	0.8	97.4	1.0	96.3	1.9	22.4	2.1	33.0	3.9	270.8	62.6	6.3	1.5
Male	70.7	6.1	98.9	0.3	97.8	0.6	97.6	0.7	17.7	2.5	46.5	3.2	177.1	11.8	3.8	1.2
<b>Area of Residence</b>																
Urban	79.0	4.3	98.8	0.3	97.9	0.4	97.8	0.4	20.6	1.6	39.8	3.7	250.2	40.9	5.1	1.1
Rural	71.4	5.9	98.0	1.8	96.4	2.5	93.4	4.9	19.8	3.4	34.0	6.5	175.0	31.5	6.0	2.4
<b>Age Group</b>																
18-29y	75.0	7.5	99.6	0.2	97.9	0.7	97.3	1.0	22.8	4.0	41.4	4.2	165.5	18.3	4.5	1.4
30-44y	77.7	3.2	98.4	0.6	97.4	2.0	93.5	4.0	20.1	2.5	40.8	4.9	290.0	61.4	5.9	1.6
45-59y	77.0	4.7	97.7	1.4	97.5	0.9	98.4	0.7	17.6	1.7	39.8	3.1	187.0	20.1	4.7	1.2
60-69y	78.6	5.1	100.0	0.0	98.5	1.4	98.6	0.7	22.1	4.4	25.9	5.1	191.6	24.2	10.0	6.4
70+y	82.0	6.2	98.0	1.5	96.1	1.5	99.6	0.4	21.7	3.2	36.1	4.1	399.5	140.4	1.5	0.9
<b>Education Years</b>																
0	53.7	16.0	100.0	0.0	100.0	0.0	100.0	0.0	17.6	3.8	36.5	15.6	93.4	50.8	0.0	0.0
1-5	77.8	6.9	98.4	1.3	97.6	2.1	99.5	0.5	25.0	4.5	23.9	7.0	183.1	27.8	5.1	2.0
6-11	78.2	5.0	98.6	0.9	98.6	0.4	98.6	0.5	21.4	2.2	33.8	3.1	264.5	83.4	7.1	1.4
12+	76.9	3.6	98.7	0.3	96.4	1.2	94.4	2.8	18.8	1.9	46.0	5.0	212.5	16.2	3.6	1.3
<b>Income Quintiles</b>																
Q1	71.3	6.4	99.5	0.3	98.5	0.8	98.9	0.6	17.9	2.0	31.6	5.6	118.3	20.8	8.7	2.4
Q2	73.8	5.4	98.9	0.7	93.4	2.9	93.2	6.2	23.1	5.3	26.0	6.5	183.3	22.3	5.8	1.9
Q3	71.0	6.0	98.8	0.5	99.0	0.8	97.0	1.3	23.7	2.5	28.8	7.1	195.7	21.9	10.2	2.3
Q4	84.0	3.1	97.6	1.9	97.8	1.0	98.4	0.7	20.1	2.8	44.2	6.8	339.1	84.2	1.7	0.7
Q5	82.8	4.6	98.7	0.4	98.4	0.4	96.4	0.7	17.5	1.9	55.9	2.5	274.3	30.6	2.1	0.8
<b>Condition</b>																
Child-Visit	77.3	13.1	100.0	0.0	100.0	0.0	98.8	1.0	18.8	3.0	50.9	10.3	105.8	18.6	4.1	2.8
Preventive	100.0	0.0	100.0	0.0	91.8	5.6	100.0	0.0	20.9	5.1	47.4	17.8	232.3	69.5	13.8	13.4
Childbirth	100.0	0.0	100.0	0.0	100.0	0.0	100.0	0.0	15.5	0.5	100.0	0.0			0.0	0.0
Acute	77.4	6.6	97.5	0.9	97.5	1.6	94.0	2.6	22.3	3.7	49.0	4.7	142.6	24.5	7.2	2.5
Injury	60.3	11.7	99.5	0.5	98.2	1.4	99.5	0.5	19.5	3.2	54.0	5.7	133.6	35.9	9.9	5.0
Chronic	85.4	6.5	92.6	5.7	95.3	2.0	96.6	1.8	18.3	1.4	52.7	10.4			8.5	4.3

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
Other	79.4	3.4	99.3	0.2	97.8	0.8	96.5	1.7	19.9	1.9	34.2	2.5	140.8	16.0	3.3	0.6
<b>Ownership</b>																
Government	76.0	4.9	99.2	0.4	96.8	1.0	94.5	3.5	22.8	2.9	27.6	3.8	86.9	11.4	6.7	1.8
Private	78.7	3.6	98.3	0.7	97.9	0.9	97.5	0.6	19.6	1.8	45.3	3.4	280.4	46.0	4.8	1.0
NGO																
Other	58.7	18.1	100.0	0.0	100.0	0.0	100.0	0.0	14.3	4.7	18.8	6.7	263.7	115.4	0.0	0.0
<b>Self-Assessed Health</b>																
Very-Good	74.0	5.2	99.6	0.2	99.2	0.4	97.8	0.9	22.3	2.5	43.7	4.7	151.3	23.4	7.2	2.9
Good	78.5	4.2	98.9	0.2	98.6	0.4	97.9	0.5	18.7	1.7	39.8	3.4	280.4	51.8	3.9	0.8
Moderate	78.8	4.2	97.3	2.3	92.8	2.4	92.1	5.2	23.5	4.2	32.1	4.0	176.0	17.1	5.2	1.7
Bad	76.8	9.3	95.3	4.0	96.9	1.7	92.8	4.2	17.1	2.8	25.9	7.4	245.3	51.9	11.4	6.3
Very-Bad	45.5	35.1	100.0	0.0	100.0	0.0	100.0	0.0	16.0	4.0	0.0	0.0	36.0	0.8	45.5	35.1
TOTAL	77.4	4.1	98.7	0.5	97.6	0.6	96.8	1.2	20.4	1.6	38.6	2.9	233.2	39.4	5.3	1.0

This table looks at various characteristics of a visit to an ambulatory care facility, for those who utilized such services in the previous 12 months (Q7204 and Q7205). Characteristics of a visit reported here include:

- (1) The percent of patients visiting a medical doctor (of those visiting any ambulatory provider) (Q7302).
- (2) Patients' rating of *skills*, *equipment*, and *drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7304, Q7305 and Q7306, respectively.
- (3) Barriers or facilitators to care access have been reported here for those who utilized ambulatory care services in the last 12 months. They include:
  - (a) *Mean time taken to reach facility*: patient reported times from Q7307 were used to calculate averages across sample sub-groups.
  - (b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7308.
  - (c) *Mean cost of visit*: patient reported costs in Q7309-Q7313 were summed up for an individual and then averaged across sample sub-groups.
  - (d) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination listed in Q7328-Q7334.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include: sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents could select one of the

following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

### 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>												
Female	15.8	1.7	27.9	2.0	7.2	1.8	3.8	1.4	10.4	2.8	18.4	3.4
Male	12.9	2.9	26.9	3.4	4.9	0.9	5.1	1.2	11.9	2.9	19.3	3.1
<b>Area of Residence</b>												
Urban	15.3	1.8	29.8	2.3	6.3	1.0	4.8	1.4	11.7	1.5	19.4	2.1
Rural	12.2	3.1	18.8	4.7	6.1	2.1	2.8	1.8	8.3	2.5	16.5	4.6
<b>Age Group</b>												
18-29y	17.9	4.9	30.9	4.7	8.0	2.1	4.1	1.1	10.4	2.1	18.9	3.9
30-44y	14.0	2.7	26.7	3.6	6.0	1.3	3.0	1.0	13.0	2.0	20.2	2.6
45-59y	13.5	2.9	26.3	2.9	5.0	2.4	7.2	3.1	10.9	2.4	19.2	2.7
60-69y	15.7	4.9	28.0	5.9	9.9	5.7	3.2	2.9	12.2	6.1	23.3	7.1
70+y	10.2	2.4	23.7	3.6	1.8	1.2	2.4	1.9	4.7	2.0	6.5	3.1
<b>Education Years</b>												
0	8.7	6.6	34.8	19.7	4.1	4.3	4.1	4.3	4.1	4.3	4.1	4.3
1-5	12.4	2.8	30.3	7.7	5.2	1.8	3.2	1.8	7.3	3.3	14.8	4.1
6-11	16.6	2.2	28.7	2.5	6.1	1.9	4.5	1.7	9.7	2.0	18.5	2.5
12+	13.1	2.3	25.7	2.7	6.6	1.1	4.4	0.9	13.2	1.3	20.0	2.2
<b>Income Quintiles</b>												
Q1	15.4	3.6	27.2	4.4	7.6	2.2	3.8	1.6	9.0	2.4	19.3	3.6
Q2	13.7	4.2	34.5	7.3	5.6	1.9	7.7	2.9	9.0	2.7	16.0	3.2
Q3	18.8	3.2	26.0	3.2	9.1	2.6	2.0	0.8	11.5	2.2	19.9	3.8
Q4	12.5	3.5	22.3	3.2	2.5	1.0	3.7	1.9	11.8	2.7	20.1	2.6
Q5	13.5	2.9	29.4	3.2	6.7	2.0	5.2	1.7	12.7	1.4	17.7	2.2
<b>Condition</b>												
Child-Visit	3.1	2.0	15.4	4.1	5.5	2.4	3.7	1.7	4.9	2.4	7.7	2.3
Preventive	8.2	5.6	29.8	12.2	0.0	0.0	0.0	0.0	8.2	5.6	8.2	5.6
Childbirth	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Acute	10.1	1.8	30.7	4.0	4.6	1.5	3.7	1.2	18.4	3.0	17.5	2.7
Injury	17.2	6.2	17.3	6.1	8.5	5.1	5.7	3.6	4.2	2.0	14.4	5.3
Chronic	12.2	4.4	26.3	6.0	6.6	3.9	3.7	2.0	17.8	6.3	26.0	7.7
Other	13.0	1.8	27.4	2.3	6.0	0.9	3.4	1.0	7.6	1.4	15.9	1.8
<b>Ownership</b>												
Government	17.7	2.4	34.7	3.7	9.9	2.1	4.9	1.4	14.5	3.5	25.7	3.3
Private	11.8	1.8	24.4	2.5	4.5	0.8	3.5	0.8	9.2	1.6	15.2	1.7
NGO												
Other	6.3	3.3	13.7	7.2	7.4	8.1	7.4	8.1	12.1	7.3	16.9	7.3
<b>Self-Assessed Health</b>												
Very-Good	12.5	2.5	27.5	3.2	8.9	2.5	4.9	1.8	15.5	3.3	23.4	3.8
Good	13.7	2.1	26.0	2.2	4.2	0.6	3.4	0.9	9.4	1.4	16.6	1.6
Moderate	15.6	2.8	31.8	4.2	8.7	2.4	5.4	2.4	9.7	2.9	20.4	4.3
Bad	11.8	6.0	31.4	14.0	7.2	5.0	0.0	0.0	11.9	5.0	8.1	3.9
Very-Bad	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TOTAL	14.6	1.7	27.5	2.1	6.3	0.9	4.3	1.1	11.0	1.3	18.7	1.8

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

### 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Autonomy				Confidentiality				Choice of		Basic Amenities			
	Treatment Information		Involvement		Talk Privately		Confidentiality of Records		Healthcare Provider		Cleanliness		Space	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female	24.9	2.9	31.2	3.0	13.0	3.9	7.7	2.3	27.2	5.2	17.8	2.4	24.9	2.7
Male	23.6	3.5	27.1	3.2	9.3	1.7	7.8	1.4	27.3	3.4	10.7	2.2	17.5	4.0
Area of Residence														
Urban	25.4	1.8	30.6	2.1	10.8	2.3	7.9	1.9	25.5	2.6	14.9	1.4	21.4	1.6
Rural	20.5	6.2	25.4	6.3	14.1	5.3	7.2	2.9	33.7	9.0	15.0	4.1	23.6	3.6
Age Group														
18-29y	24.7	4.0	28.4	3.6	13.0	2.8	9.2	2.0	25.1	4.6	14.7	3.3	23.5	4.2
30-44y	27.3	3.3	32.1	3.4	12.6	4.9	8.0	1.4	30.6	5.2	12.4	2.5	23.6	4.1
45-59y	23.3	2.7	31.6	3.0	11.8	2.2	5.8	2.4	28.6	3.8	16.8	2.7	19.6	2.4
60-69y	28.8	6.8	31.5	6.6	9.0	5.7	13.1	7.4	26.2	6.8	21.2	5.5	29.2	7.0
70+y	11.5	3.8	14.9	3.7	5.8	2.8	2.7	1.8	18.7	3.4	10.0	2.6	9.4	2.7
Education Years														
0	17.5	13.1	13.3	12.5	19.9	10.9	12.1	8.5	40.5	20.9	34.8	12.4	4.2	4.3
1-5	22.7	4.6	23.4	5.2	8.9	2.8	9.5	5.2	28.6	5.9	18.9	5.4	22.2	6.3
6-11	21.1	2.0	28.0	2.2	10.3	2.8	7.7	2.3	28.6	3.8	17.3	2.0	24.4	2.2
12+	28.2	2.6	32.5	3.0	13.0	3.2	7.5	1.0	25.3	3.9	11.2	1.5	19.7	1.9
Income Quintiles														
Q1	22.5	4.3	26.1	4.6	10.8	2.7	9.0	2.4	34.0	5.0	22.2	5.2	26.9	4.5
Q2	18.2	3.9	25.1	5.3	15.3	5.7	6.3	1.9	37.4	7.7	14.2	3.1	25.5	3.5
Q3	30.4	3.8	32.3	3.9	11.3	3.6	9.7	2.9	28.7	4.0	15.9	3.4	24.9	3.3
Q4	22.0	2.8	30.5	2.2	11.5	3.8	7.3	2.3	21.5	5.3	12.6	2.4	17.0	3.0
Q5	25.5	2.6	30.1	4.1	9.7	2.3	6.8	2.0	20.6	1.9	11.6	2.6	18.2	1.7
Condition														
Child-Visit	12.7	3.6	14.0	5.9	11.7	3.5	7.3	3.2	16.6	6.3	8.2	2.5	13.7	5.2
Preventive	16.5	11.2	16.5	11.2	17.3	16.6	0.0	0.0	8.0	8.2	17.3	16.6	16.5	11.2
Childbirth	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Acute	27.5	4.1	34.8	5.2	11.5	4.8	10.6	3.3	34.5	6.1	12.5	2.5	23.5	3.4
Injury	27.4	6.7	20.4	6.4	8.1	5.0	7.6	5.0	24.8	7.3	6.3	2.7	12.0	5.2
Chronic	33.4	8.4	35.7	7.3	22.2	7.9	6.9	2.9	38.0	8.8	22.2	8.1	17.3	4.5
Other	21.3	2.4	27.8	2.7	8.8	2.1	6.1	1.2	24.4	3.2	13.0	1.7	20.7	1.9
Ownership														
Government	27.9	3.0	32.9	3.3	18.4	4.6	10.2	2.5	39.8	3.7	29.7	3.9	34.9	4.0
Private	22.7	2.4	28.0	2.2	8.2	1.5	6.1	1.2	20.6	2.3	6.5	1.0	14.2	1.6
NGO														
Other	16.9	7.3	16.9	7.3	4.7	2.5	3.1	1.7	34.9	13.7	7.4	8.1	7.4	8.1
Self-Assessed Health														
Very-Good	28.2	3.6	35.0	4.3	15.3	4.2	9.7	2.6	29.2	3.4	14.0	2.7	17.5	2.5
Good	22.2	2.6	26.6	2.6	7.9	1.5	6.8	1.4	25.2	2.5	12.5	1.3	20.9	1.6
Moderate	24.6	4.3	29.9	4.9	16.0	6.0	6.5	2.2	30.8	9.2	18.0	3.3	24.7	2.6
Bad	30.8	11.3	29.0	11.0	11.7	5.3	1.8	1.3	18.6	7.3	16.9	7.4	17.5	11.1
Very-Bad	45.5	35.1	45.5	35.1	0.0	0.0	0.0	0.0	45.5	35.1	0.0	0.0	0.0	0.0
TOTAL	24.3	2.0	29.5	2.0	11.5	2.3	7.7	1.6	27.2	3.3	14.9	1.4	21.9	1.5



The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.4 Patient assessed characteristics of hospital inpatient health services

	Patients staying for more than 6 days on last hospital visit		Patients rating characteristics of health care provider as adequate / more than adequate				Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Patients waiting more than a month to be admitted		Mean number of patients sleeping in the same room		Mean cost for a hospital visit		Percent of population perceiving discrimination			
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	%	SE	No	SE	mean cost	SE	%	SE
Sex																				
Female	27.5	3.1	97.4	1.0	97.1	1.0	98.1	0.8	25.4	3.6	60.9	3.2	1.5	0.6	2.9	0.7	1,553.0	0.5	6.6	1.3
Male	36.4	3.9	98.3	0.5	96.8	1.0	96.0	1.1	38.5	9.5	45.6	3.2	1.8	1.0	2.7	0.7	973.6	0.5	7.8	3.0
Area of Residence																				
Urban	30.2	2.9	97.3	0.7	96.7	0.8	97.0	0.5	24.8	3.4	53.4	2.9	2.2	0.7	2.7	0.6	1,374.2	0.5	8.0	2.1
Rural	34.7	7.7	99.2	0.5	97.9	2.0	97.8	1.4	49.2	9.3	56.8	5.0	0.0	0.0	2.9	0.8	1,106.4	0.5	4.6	2.2
Age Group																				
18-29y	15.2	3.5	95.0	2.6	92.4	2.7	96.2	1.1	24.7	4.4	54.8	4.1	1.6	1.0	3.4	0.9	1,543.0	0.5	11.9	3.2
30-44y	22.4	4.0	98.0	0.6	97.4	0.9	96.9	1.0	31.7	4.0	64.9	4.1	0.5	0.5	2.7	0.8	1,730.2	0.5	5.5	1.7
45-59y	39.4	6.0	97.9	0.9	98.0	0.9	96.7	1.8	40.3	14.9	48.6	5.3	1.2	0.8	2.7	0.8	1,335.7	0.5	5.9	2.3
60-69y	43.2	6.6	99.2	0.8	99.2	0.8	97.4	2.8	33.6	9.2	45.9	9.1	0.0	0.0	2.7	1.0	571.0	0.5	8.9	3.3
70+y	53.5	5.3	100.0	0.0	99.3	0.3	100.0	0.0	21.8	4.8	46.3	7.3	6.7	2.6	2.1	0.8	535.3	0.5	3.4	2.2
Education Years																				
0	30.7	26.4	100.0	0.0	100.0	0.0	100.0	0.0	33.5	9.2	24.3	14.3	0.0	0.0	2.5	0.8	34.1	0.5	12.0	13.0
1-5	56.5	6.9	100.0	0.0	100.0	0.0	100.0	0.0	24.9	6.1	43.9	6.3	4.7	3.2	2.7	0.8	469.0	0.5	3.7	2.6
6-11	31.6	4.8	97.4	1.0	96.7	1.5	96.6	0.8	34.9	7.4	49.8	2.6	2.1	1.0	3.0	0.7	998.7	0.5	9.8	1.9
12+	23.3	2.8	97.6	0.6	96.4	1.2	97.2	0.8	27.2	4.9	65.2	2.8	0.0	0.0	2.5	0.8	1,997.9	0.5	4.0	1.5
Income Quintiles																				
Q1	38.3	6.5	99.5	0.5	97.4	1.5	98.7	1.0	29.3	3.3	45.1	7.3	0.0	0.0	3.8	1.0	325.8	0.5	5.4	2.3
Q2	30.2	5.0	98.6	0.7	98.1	1.5	97.9	0.9	50.5	14.1	41.1	4.8	0.5	0.5	3.2	0.9	1,093.0	0.5	9.8	3.3
Q3	34.5	5.2	98.3	0.8	95.3	2.2	98.6	0.9	29.8	6.0	52.5	6.4	2.3	1.0	2.6	0.8	979.4	0.5	12.0	5.3
Q4	29.2	4.1	95.1	1.4	93.6	3.0	94.9	2.1	27.2	5.9	56.8	5.6	4.3	2.1	2.3	0.8	1,230.0	0.5	4.2	1.7
Q5	25.1	4.3	97.1	0.5	99.6	0.5	95.7	0.9	18.5	1.4	76.5	3.8	1.6	0.4	1.8	0.6	3,037.6	0.5	4.9	0.9
Condition																				
Child-Visit	20.6	6.7	97.7	1.4	96.4	1.7	97.8	1.3	27.6	4.2	58.2	9.8	0.0	0.0	2.8	0.7	217.1	131.8	12.4	4.6
Fever	17.0	8.7	98.3	1.5	96.8	2.1	98.3	1.5	31.7	8.0	66.4	15.2	0.0	0.0	3.3	1.0	172.3	98.5	11.0	6.3
Childbirth	7.7	2.7	98.2	1.1	99.1	0.6	98.0	1.2	21.3	2.3	68.4	4.3	0.0	0.0	2.9	1.2	2,448.1	799.4	8.8	3.5
Arthritis	9.2	11.2	100.0	0.0	100.0	0.0	100.0	0.0	11.0	6.1	72.3	25.8	0.0	0.0	2.1	1.1	102.9	16.9	18.5	20.8
Asthma	26.4	12.3	87.3	6.7	87.3	6.7	84.2	11.5	27.1	8.3	52.4	13.4	0.0	0.0	2.8	0.7	49.6	12.0	10.8	10.5
Heart-Disease	68.0	9.1	100.0	0.0	100.0	0.0	100.0	0.0	46.0	18.8	21.8	7.2	0.0	0.0	3.3	1.8	272.2	71.2	3.0	1.9
Injury	33.4	8.9	97.3	1.5	92.6	3.2	97.3	2.1	64.2	29.0	22.6	7.1	0.0	0.0	2.8	1.0	1,737.3	1,066.6	9.4	5.7
Minor-Surgery	28.8	4.5	96.0	1.5	100.0	0.0	96.3	1.5	25.7	6.5	60.2	5.2	7.5	3.1	2.8	1.0	654.8	187.2	0.9	1.0
Other	39.5	4.1	98.4	0.5	95.3	1.6	97.1	1.5	28.7	4.9	55.0	4.1	0.4	0.4	2.6	0.7	586.2	202.5	8.5	2.6
Ownership																				
Government	31.8	4.2	98.4	0.9	97.2	1.0	99.6	0.3	33.1	3.9	43.2	4.8	3.0	2.0	3.9	0.9	150.0	34.0	11.4	3.4
Private	30.5	2.7	97.1	0.6	96.6	0.9	95.4	0.8	29.3	8.0	62.3	2.7	0.6	0.2	1.8	0.6	1,451.5	310.1	4.1	1.0
NGO																				
Other	50.0	0.0	100.0	0.0	100.0	0.0	100.0	0.0	83.0	0.5	0.0	0.0	0.0	0.0	20.5	0.5			0.0	0.0
Self-Assessed Health																				
Very-Good	17.0	3.3	100.0	0.0	97.0	3.1	99.1	1.0	26.4	4.3	67.8	4.5	3.3	1.9	1.9	0.6	449.2	136.8	7.1	3.2
Good	26.7	3.3	96.5	1.1	96.8	0.9	96.6	0.8	28.8	4.5	57.0	2.6	0.9	0.4	3.0	0.7	758.1	291.3	8.2	1.5
Moderate	47.3	6.9	98.9	0.5	97.4	1.0	96.8	0.9	39.3	13.8	39.8	6.3	2.0	0.9	2.8	0.8	1,465.8	447.0	3.4	1.7
Bad	44.4	12.1	92.9	3.4	92.9	3.4	95.1	3.1	26.1	5.3	53.0	11.3	0.0	0.0	2.7	1.0	1,030.6	131.8	17.1	9.6
Very-Bad	100.0	0.0	100.0	0.0	100.0	0.0	100.0	0.0	35.5	0.5	0.0	0.0	0.0	0.0	2.5	0.5	3,000.5	0.5	0.0	0.0
TOTAL	31.4	2.7	97.8	0.7	97.0	0.8	97.2	0.6	31.1	5.7	54.2	2.3	1.6	0.8	2.8	0.6	1,295.5	0.5	7.1	1.5

This table looks at various characteristics of a visit to a hospital inpatient care facility, for those who utilized such services in the past 3 years (Q7200, Q7201 and Q7402). Characteristics of a visit reported here include:

(1) The percent of patients who stayed 6 days or more at an inpatient facility during their last visit (Q7404).

(2) Patients' rating of *skills, equipment, and drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7405, Q7406 and Q7407, respectively.

(3) Barriers or facilitators to care access have been reported here for those who utilized inpatient care services in the last 3 years. They include:

(a) *Mean time taken to reach facility*: patient reported times from Q7408 were used to calculate averages across sample sub-groups.

(b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7409.

(c) *Patients waiting more than a month to be admitted*: percent of patients who reported waiting more than a month to be admitted in Q7410.

(d) *Mean number of patients sleeping in same room*: averages across sample sub-groups have been constructed for numbers reported in Q7417.

(e) *Mean cost of visit*: patient reported costs in Q7411-Q7415 were summed up for an individual and then averaged across sample sub-groups.

(f) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination asked in Q7433-Q7439.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication				Autonomy			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions		Treatment Information		Involvement	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>																
Female	11.4	2.8	11.5	2.0	6.2	1.3	9.9	2.1	13.1	2.9	17.7	2.6	20.0	2.6	21.5	2.4
Male	13.4	3.4	19.0	2.5	4.6	2.1	4.7	1.8	8.5	1.7	15.1	2.1	18.9	2.1	19.9	2.4
<b>Area of Residence</b>																
Urban	9.5	1.8	14.8	1.4	5.7	1.4	7.4	1.6	11.5	1.9	17.5	1.7	21.0	1.9	21.2	1.8
Rural	20.3	5.0	14.5	4.9	4.9	2.1	8.6	3.7	10.0	4.1	13.8	5.1	15.4	4.9	19.5	5.6
<b>Age Group</b>																
18-29y	16.1	2.8	19.9	3.6	8.6	2.5	13.9	3.2	15.3	3.3	27.3	6.7	27.7	6.5	22.5	4.7
30-44y	7.5	2.9	12.5	2.4	6.4	1.7	5.8	1.3	10.3	2.5	13.1	2.8	15.1	2.8	18.0	3.4
45-59y	17.5	5.7	13.7	3.4	4.2	2.0	6.1	2.6	9.1	2.8	11.2	2.8	19.8	4.2	23.1	4.1
60-69y	15.3	4.7	14.7	4.5	3.4	3.0	5.5	4.0	10.7	4.3	19.2	7.7	18.3	4.8	23.8	4.0
70+y	5.1	3.1	13.3	4.0	2.5	1.6	6.8	4.5	10.0	4.6	13.3	6.2	17.6	7.8	17.5	8.3
<b>Education Years</b>																
0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	18.4	14.2	18.4	14.2	23.2	14.2	23.2	14.2
1-5	9.5	3.7	11.2	4.9	3.9	1.5	6.4	5.3	10.2	5.0	18.1	6.0	22.8	5.8	22.2	5.6
6-11	14.0	3.2	15.4	3.0	5.6	1.6	7.9	1.8	10.5	2.4	15.0	2.6	18.2	2.7	19.5	2.7
12+	11.0	2.5	15.4	4.1	6.1	1.8	8.0	1.7	12.0	2.1	18.4	2.7	20.5	3.3	22.1	3.0
<b>Income Quintiles</b>																
Q1	14.3	3.4	17.2	4.1	5.8	1.9	9.6	3.0	8.8	3.2	16.6	3.6	18.5	3.7	23.2	5.0
Q2	21.1	5.4	14.5	3.5	3.8	1.6	12.4	3.6	13.1	3.5	17.8	4.2	22.4	4.3	17.6	4.3
Q3	13.9	3.8	14.3	4.4	11.6	5.5	6.1	3.5	12.8	5.1	11.7	4.8	15.7	3.8	20.6	4.0
Q4	10.2	2.9	19.2	5.5	3.7	1.1	3.0	0.9	12.3	3.9	19.1	3.6	21.0	3.5	28.3	3.5
Q5	2.5	0.5	9.8	1.9	3.6	0.6	6.7	1.1	9.5	1.8	18.0	1.9	20.6	2.4	16.0	2.1
<b>Condition</b>																
Child-Visit	16.2	5.5	19.8	5.6	8.4	3.3	7.6	2.9	12.9	4.6	17.4	5.2	20.4	6.0	19.5	5.8
Fever	11.7	6.1	13.6	7.4	9.0	5.8	7.4	4.9	15.1	11.8	15.3	13.3	23.9	15.1	21.8	15.0
Childbirth	7.5	2.3	6.2	2.4	7.1	3.0	11.1	3.7	13.1	3.9	16.1	3.6	14.1	3.9	16.4	4.3
Arthritis	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	18.5	20.8	18.5	20.8	18.5	20.8	0.0	0.0
Asthma	24.4	9.4	35.5	12.2	4.3	4.4	7.8	4.7	11.3	6.3	24.1	11.1	30.8	13.1	31.9	13.0
Heart-Disease	9.8	7.6	0.0	0.0	0.0	0.0	3.1	3.2	4.6	5.0	7.7	6.2	19.2	5.7	19.2	5.7
Injury	14.8	10.2	21.6	7.2	0.6	0.6	1.5	0.8	4.5	1.9	13.6	5.5	16.3	5.0	23.2	6.9
Minor-Surgery	13.5	5.9	18.7	4.6	2.5	1.0	3.9	1.7	9.7	2.3	13.8	2.5	18.7	2.6	18.7	2.8
Other	12.6	4.2	16.2	2.8	8.2	1.9	10.1	4.2	13.3	3.4	20.4	3.1	22.2	3.4	23.1	3.5
<b>Ownership</b>																
Government	16.8	3.2	17.8	2.9	5.6	1.9	8.6	2.6	11.6	2.9	17.2	3.4	19.6	3.1	23.4	3.4
Private	8.8	2.8	12.1	2.1	5.7	1.0	7.1	1.2	11.2	1.6	16.7	1.9	19.5	2.2	19.1	2.4
NGO																
Other	0.0	0.0	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	50.0	0.0	0.0	0.0
<b>Self-Assessed Health</b>																
Very-Good	8.1	2.4	8.4	2.8	2.5	1.7	4.9	1.6	8.6	4.0	12.7	3.4	15.5	3.0	14.8	3.3
Good	9.5	2.0	17.5	2.4	7.3	1.4	8.0	2.0	12.6	2.2	17.5	2.6	20.4	2.6	21.0	2.5
Moderate	17.4	6.4	13.3	2.5	4.0	1.4	7.6	2.5	9.4	1.9	15.5	2.3	19.3	3.2	21.4	2.8
Bad	30.0	9.9	17.1	9.6	10.0	10.0	16.8	9.5	19.4	12.2	37.1	14.8	30.6	10.1	43.7	12.2
Very-Bad	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TOTAL	12.3	2.6	14.7	1.6	5.5	1.1	7.7	1.5	11.1	1.7	16.6	1.8	19.6	2.0	20.8	2.0

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Confidentiality				Choice of health care provider		Basic Amenities				Social Support			
	Talk Privately		Confidentiality of records			SE	Cleanliness		Space		Family Visit		External Contact	
	%	SE	%	SE			%	SE	%	SE	%	SE	%	SE
Sex														
Female	24.6	4.7	13.1	3.4	40.7	3.6	20.8	3.7	31.9	3.6	27.4	2.8	24.9	2.5
Male	15.5	2.4	9.7	2.4	38.1	4.1	15.2	2.4	22.5	2.6	20.4	2.6	22.6	3.3
Area of Residence														
Urban	19.2	2.4	11.6	3.1	38.0	2.2	18.8	1.8	27.7	1.8	24.2	1.8	23.6	2.0
Rural	24.8	6.8	11.4	4.3	44.2	6.5	16.9	7.1	28.0	7.7	24.8	5.8	24.6	3.8
Age Group														
18-29y	20.2	4.0	12.4	3.2	53.6	3.6	28.1	3.2	32.1	4.0	35.3	4.1	30.7	3.9
30-44y	26.8	7.6	8.4	1.8	34.5	5.0	20.2	6.4	35.7	6.0	22.2	3.4	21.4	3.3
45-59y	14.7	4.1	13.8	4.8	38.5	5.5	11.9	3.0	23.3	4.1	24.4	4.6	26.8	6.6
60-69y	20.0	2.9	12.8	3.7	40.7	4.9	15.5	3.5	22.3	3.7	15.2	3.8	14.8	2.6
70+y	17.5	4.7	11.8	6.7	29.0	6.8	11.6	3.4	15.4	3.6	20.4	4.1	22.7	4.3
Education Years														
0	18.4	14.2	30.4	15.7	30.4	15.7	0.0	0.0	3.8	4.0	0.0	0.0	0.0	0.0
1-5	24.4	6.9	15.1	5.6	31.0	7.1	9.7	4.0	18.7	5.3	12.5	4.2	15.3	4.7
6-11	19.4	4.6	10.4	2.6	45.1	4.0	22.2	4.0	30.0	4.4	26.6	3.0	28.8	2.4
12+	21.4	3.4	11.3	2.7	34.4	3.7	16.1	2.2	28.3	3.5	25.6	2.5	20.1	3.1
Income Quintiles														
Q1	25.7	8.3	13.4	3.8	44.9	7.7	32.1	8.4	40.5	8.8	25.9	5.0	23.8	5.7
Q2	23.8	5.2	14.8	4.0	45.0	6.2	18.8	4.5	29.1	5.8	28.0	4.9	29.9	5.6
Q3	20.0	4.6	11.5	5.4	45.3	4.6	20.1	4.0	28.9	4.4	25.0	4.2	29.3	4.4
Q4	23.2	4.5	10.7	4.1	33.6	5.2	8.7	2.5	21.5	3.9	20.5	4.6	21.2	5.3
Q5	12.0	2.6	7.9	1.8	31.3	3.9	12.3	1.4	20.0	2.7	23.0	2.2	16.8	2.6
Condition														
Child-Visit	32.0	13.8	8.0	3.0	35.8	9.0	35.0	13.2	40.2	12.4	28.1	6.6	25.3	6.1
Fever	55.2	21.2	10.5	6.9	29.7	17.2	63.4	17.5	67.2	16.5	27.0	13.6	27.0	13.6
Childbirth	20.1	6.4	7.9	2.3	33.8	5.0	18.6	4.5	26.6	7.6	23.6	3.7	24.7	4.0
Arthritis	18.5	20.8	18.5	20.8	18.5	20.8	18.5	20.8	18.5	20.8	0.0	0.0	18.5	20.8
Asthma	28.6	12.7	13.7	6.4	69.3	13.6	20.7	10.6	27.5	12.9	36.8	15.2	34.9	15.4
Heart-Disease	14.9	5.2	12.3	5.3	30.0	7.6	0.0	0.0	5.2	4.2	18.7	11.1	25.9	10.1
Injury	16.2	6.4	7.2	3.6	51.3	9.0	8.6	4.3	21.8	5.7	21.5	6.4	26.1	10.3
Minor-Surgery	17.6	3.2	6.2	2.6	37.5	3.9	19.1	3.2	23.2	4.2	28.3	4.8	23.3	4.7
Other	19.1	3.7	15.6	4.9	40.8	4.9	18.0	2.7	30.6	3.7	23.6	2.7	21.9	2.8
Ownership														
Government	25.6	4.7	11.6	3.4	52.0	4.5	32.8	4.8	39.9	5.0	31.5	3.8	31.5	4.0
Private	17.1	2.6	10.9	1.8	30.4	3.1	8.8	1.6	19.4	1.9	19.6	1.9	18.6	2.6
NGO														
Other	0.0	0.0	0.0	0.0	50.0	0.0	50.0	0.0	50.0	0.0	0.0	0.0	0.0	0.0
Self-Assessed Health														
Very-Good	15.1	5.4	1.1	0.7	35.8	7.4	13.7	2.7	23.9	3.7	20.3	4.2	16.6	5.4
Good	22.5	4.8	12.3	2.7	38.5	3.4	22.6	4.0	30.2	3.4	24.4	2.8	24.2	2.1
Moderate	18.4	3.5	13.7	3.8	39.1	4.5	13.5	2.4	23.3	3.7	22.2	3.6	23.9	4.0
Bad	31.8	11.2	25.3	11.1	68.6	10.5	27.4	9.9	48.1	12.4	61.6	11.6	52.2	11.9
Very-Bad	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TOTAL	20.6	2.8	11.6	2.5	39.6	2.3	18.4	2.2	27.8	2.4	24.3	2.0	23.9	1.8

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

## 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"

	Prompt Attention		Dignity		Communication		Autonomy		Confidentiality		Choice		Basic Amenities		Social Support	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>Sex</b>																
Female	21.8	1.6	31.4	2.0	43.7	2.1	20.0	1.8	38.2	1.8	23.1	2.0	31.3	2.3	20.2	2.0
Male	20.6	2.2	25.5	3.2	35.2	3.5	16.3	2.0	31.3	2.9	18.4	2.2	25.9	2.4	19.3	2.3
<b>Area of Residence</b>																
Urban	22.0	1.9	29.2	2.5	40.2	3.0	19.8	2.1	36.7	1.9	22.4	2.2	29.5	2.3	21.9	2.2
Rural	18.8	3.4	26.7	4.0	37.9	4.0	13.2	2.3	29.2	3.2	15.9	2.9	26.4	4.4	12.5	2.7
<b>Age Group</b>																
18-29y	23.5	3.1	26.9	3.7	42.0	4.0	20.9	2.8	41.1	3.3	22.1	2.5	34.9	3.5	24.7	2.8
30-44y	22.6	2.5	29.8	2.7	41.1	3.2	19.5	2.8	35.0	2.5	22.7	2.7	30.1	2.8	19.5	2.6
45-59y	21.8	2.4	30.5	2.3	41.3	3.0	18.4	2.5	35.4	2.1	21.7	2.8	27.6	2.8	20.1	2.4
60-69y	15.8	1.8	26.5	3.0	35.1	2.8	13.2	1.8	28.7	3.0	17.7	2.3	22.7	3.5	14.8	2.3
70+y	16.6	2.2	26.2	3.6	30.6	3.2	13.6	2.2	26.1	4.1	14.0	2.7	19.7	2.4	13.2	2.4
<b>Education Years</b>																
0	19.6	9.5	10.9	6.6	23.1	7.5	2.4	1.9	11.7	6.1	6.6	5.2	21.9	7.9	4.7	3.3
1-5	8.2	1.9	17.7	3.2	22.3	3.2	9.2	2.4	18.5	4.1	9.4	2.0	14.0	3.5	11.4	2.5
6-11	18.7	2.2	25.1	2.4	35.1	2.6	15.7	1.9	30.3	2.0	17.7	2.2	24.5	2.6	18.7	2.3
12+	27.0	1.9	35.5	2.2	49.2	2.2	23.6	2.2	44.7	2.8	27.5	2.6	37.0	2.5	23.1	2.5
<b>Income Quintiles</b>																
Q1	17.1	3.1	20.5	3.6	29.1	2.9	12.2	3.0	19.1	2.5	14.0	3.0	23.9	3.4	14.5	2.8
Q2	19.1	3.1	28.4	4.2	37.9	3.5	18.2	3.0	32.2	4.2	18.2	3.6	27.5	3.7	17.1	2.8
Q3	18.0	2.6	24.5	3.1	33.1	4.1	14.0	1.8	31.8	2.5	14.2	2.4	21.3	2.9	18.1	4.3
Q4	19.3	2.2	30.5	3.3	42.2	2.7	19.1	2.1	39.4	2.7	23.0	3.5	30.2	2.9	20.7	2.6
Q5	31.5	3.8	37.0	1.9	53.1	2.8	25.9	3.2	48.6	3.4	32.5	3.7	38.9	3.3	26.8	1.7
<b>Ownership</b>																
Government	14.7	2.5	20.0	2.5	31.4	3.2	12.3	2.0	23.0	1.9	11.2	2.1	21.3	3.0	17.4	3.4
Private	23.1	1.3	31.2	3.4	44.5	2.2	20.1	1.4	38.3	2.2	24.2	2.2	31.7	2.4	20.8	2.1
NGO																
Other	13.4	4.4	32.4	11.2	42.0	9.2	10.9	5.4	42.9	9.6	13.4	4.4	38.7	11.5	14.8	5.0
<b>Self-Assessed Health</b>																
Very-Good	25.5	3.0	29.3	3.5	43.7	4.8	24.0	3.9	40.2	3.2	27.4	3.8	35.0	3.6	25.6	3.6
Good	19.6	1.4	27.5	2.4	37.4	2.3	15.4	1.8	33.0	2.4	17.8	1.9	25.3	2.2	17.3	1.7
Moderate	21.7	2.7	31.9	3.7	40.2	3.1	18.8	2.7	34.1	3.5	22.4	2.4	30.7	3.6	18.3	2.6
Bad	13.9	4.4	29.5	6.6	43.0	7.6	15.3	6.4	35.5	6.4	17.6	5.2	21.9	5.8	17.5	4.6
Very-Bad	0.0	0.0	0.0	0.0	55.5	17.6	18.9	18.0	0.0	0.0	0.0	0.0	47.7	13.8	43.4	13.8
<b>Type of Care</b>																
Ambulatory-Adult	19.6	1.4	28.2	3.0	40.3	2.3	18.0	1.3	34.2	2.2	18.4	1.7	29.0	2.7	19.5	2.7
Ambulatory-Child	19.5	3.7	21.2	3.3	37.7	5.0	17.3	3.4	33.9	4.1	22.6	3.8	28.6	4.5	21.1	3.8
Inpatient-Adult	18.0	3.1	33.2	6.9	38.2	6.0	16.2	4.3	36.1	6.9	22.6	4.2	25.2	4.1	21.1	4.2
Inpatient-Child	10.8	5.0	9.2	5.3	24.9	8.3	6.2	3.1	18.0	5.6	11.6	5.4	29.8	6.6	5.3	2.2
Non-Users	23.6	2.3	30.1	2.5	39.9	3.4	19.1	2.7	36.1	2.6	23.0	2.9	29.0	2.7	19.9	2.1
TOTAL	21.2	1.6	28.6	2.3	39.7	2.6	18.3	1.8	35.0	2.0	20.9	1.9	28.8	2.1	19.8	1.9

This table shows the percent of people who rated a responsiveness domain as "Extremely Important". All survey respondents were queried on their perception of the importance of each of the eight responsiveness domains in section 7100 of the WHS where each domain was described in detail (Q7100-Q7107). Respondents were asked to say whether the domain was "Extremely Important", "Very Important", "Moderately Important", "Slightly Important" or "Not



Important".

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, ownership (of health facility), self-assessed health, and type of care. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, self-assessed health and type of care sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad". Type of care is a constructed sample sub-group combining information from Q7000, Q7001 and Q7016. There are five categories of type of care: "Ambulatory adult", "Ambulatory child", "Inpatient adult", "Inpatient child" and "Non users".

---

## 8.0 Expenditure Background Indicators (2002)

---

GDP per capita (\$)	3,620
Total health expenditure per capita (\$)	361
Total population (000)	3,391
Life expectancy at birth (Both sexes)	75.2
Life expectancy at birth (male)	71.0
Life expectancy at birth (female)	79.3
Healthy life expectancy at birth (Both sexes)	66.2
Healthy life expectancy at birth (Male)	63.0
Healthy life expectancy at birth (Female)	69.4
Government funded health expenditure as a percentage of total health expenditure	29.0
Social security funded health expenditure as a percentage of total health expenditure	15.6
Private health insurance premiums as a percentage of total health expenditure	53.2
Out-of-pocket payment as a percentage of total health expenditure	17.8

## 8.1 Distribution of health payments across different population groups

	Mean Household expenditure (EXP) in local currency	% Poor	% Impoverished	% Catastrophic	OOP % EXP	OOP % non-subsistence spending	Mean OOP in local currency
<b>Catastrophic</b>							
No	11,281	15.7	0.7		2.3	3.8	232
Yes	8,873	13.1	14.1		37.4	55.5	3730
<b>Poor</b>							
No	12,751		1.2	2.4	3.4	5.3	366
Yes	2,978		0.0	2.0	2.0	3.4	44
<b>Insurance</b>							
No	5,404	24.8	0.9	2.4	2.8	4.6	178
Yes	11,462	15.2	1.1	2.4	3.2	5.0	321
<b>Residence</b>							
rural	48,134	28.2	1.2	2.9	3.4	5.8	244
urban	8,078	14.6	1.0	2.3	3.2	4.9	321
<b>Income_Quintile</b>							
1st(poorest)	3,280	76.0	4.0	2.3	2.2	3.9	63
2nd	5,061	0.0	1.1	2.0	2.1	4.1	107
3rd	6,793	0.0	0.0	2.2	2.7	4.6	177
4th	8,861	0.0	0.0	2.6	3.9	6.1	353
5th(richest)	32,209	0.0	0.0	2.8	5.0	6.3	879
total	11,224	15.6	1.0	2.4	3.2	5.0	315

OOP: Out of pocket

*Household consumption expenditure (EXP)* comprises both monetary and in-kind payment on all goods and services and the money value of the consumption of home-made products.

*Poor* - a household is considered poor when its total household expenditure is smaller than its subsistence spending  
*Impoverishment* - a non-poor household is impoverished by health payment when it becomes poor after paying for health services.

*Catastrophic health expenditure* occurs when a household's total out-of-pocket health payments equals or exceeds 40% of the households capacity to pay or non-subsistence spending.

*Out of pocket health payments (OOP)* refer to payments made by households at the point they receive health services. It includes doctor's consultation fees, purchases of medication and hospital bills. Spending on alternative and/or traditional medicine is included. Expenditure on health related transportation and special nutrition is excluded. Insurance reimbursements are also excluded.

*Subsistence spending* - household subsistence spending is the minimum requirement to maintain a basic life in a society. A poverty line is used in the analysis as subsistence spending.

*Food expenditure* is the amount spent on all foodstuffs by the household plus the value of the family's own food production consumed within the household. It excludes expenditure on alcoholic beverages, tobacco and food consumption outside the home (e.g hotel and restaurants).

*Household capacity to pay* - is defined as a household's non-subsistence spending.

---

## 8.2 Percent distribution of structure of out-of-pocket health payments

---

	Inpatient	Outpatient	Traditional	Drugs	Other	Total
<b>Catastrophic</b>						
No	1.3	2.1	1.7	47.4	47.5	100
Yes	16.0	7.0	0.7	27.0	49.3	100
<b>Poor</b>						
No	5.4	3.5	1.4	41.6	48.1	100
Yes	7.4	1.7	1.0	46.8	43.1	100
<b>Insurance</b>						
No	6.6	2.7	2.2	47.6	40.8	100
Yes	5.4	3.5	1.4	41.6	48.1	100
<b>Residence</b>						
rural	0.4	0.6	0.2	45.3	53.5	100
urban	5.8	3.7	1.5	41.5	47.6	100
<b>Income_Quintile</b>						
1st(poorest)	3.9	1.9	2.1	50.5	41.6	100
2nd	0.9	2.2	3.5	51.6	41.8	100
3rd	2.7	1.8	3.0	64.4	28.1	100
4th	2.1	1.7	1.5	46.6	48.1	100
5th(richest)	8.0	4.8	0.7	33.3	53.1	100
total	5.4	3.5	1.4	41.7	48.0	100

---

### 8.3 Financial sources used by households for paying for health services

	Savings	Sold Items	Borrow from relatives &	Borrow from others	Health insurance	Current Income	Other
<b>OOP%non_subsistence_spending</b>							
less than 10%	1.7	0.6	5.1	0.6	3.0	90.4	1.0
10-20%	5.5	0.0	10.2	0.9	2.1	96.5	0.8
20-40%	4.8	1.3	7.1	1.6	3.7	93.7	1.8
above 40%	6.2	3.0	14.3	4.6	4.3	94.1	1.6
<b>Hospitalization</b>							
No	2.2	0.6	5.6	0.8	3.1	91.4	0.9
Yes	4.1	0.8	11.2	0.7	2.7	88.5	4.2
<b>Insurance</b>							
No	0.0	0.0	9.8	2.0	0.0	77.6	3.8
Yes	2.4	0.7	5.8	0.7	3.2	91.8	1.0
<b>Residence</b>							
rural	0.8	0.6	3.8	0.3	5.4	93.7	1.0
urban	2.5	0.7	6.1	0.8	2.8	91.0	1.1
<b>Income_Quintile</b>							
1st(poorest)	0.4	0.7	4.6	0.7	2.8	84.5	1.0
2nd	1.5	0.0	4.8	0.8	2.6	89.6	2.0
3rd	1.8	1.3	8.2	0.9	2.9	93.2	1.1
4th	2.4	0.8	6.7	0.7	3.6	93.3	1.0
5th(richest)	5.7	0.4	5.3	0.8	3.2	95.7	0.4
total	2.4	0.6	5.9	0.8	3.0	91.2	1.1

## 8.4 Insurance Coverage

	Insured	SE	Mandatory Insurance	SE	Voluntary Insurance	SE	Insured as dependent	SE	Mandatory and Voluntary	SE
Q1(poorest)	14.3	3.4	9.9	3.1	15.7	3.4	8.9	3.1	9.9	3.3
Q2	16.8	3.4	16.0	4.7	16.5	3.0	18.3	7.0	13.1	3.4
Q3	19.7	2.0	21.1	2.5	19.5	2.1	21.7	4.0	21.4	5.4
Q4	22.9	2.1	24.8	2.8	22.2	1.7	23.2	4.2	22.3	3.1
Q5(richest)	26.3	6.3	28.2	6.9	26.2	6.4	27.9	8.5	33.3	10.5
Total	100		100		100		100		100	
Number	7,609.0		2,278.0		5,862.0		564.0		581.0	

## 10.1 Inability to control the important things in life - q8000

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
<b>Gender</b>													
Male	46.1	3.1	27.5	1.4	22.8	2.6	2.3	0.3	0.7	0.2	0.6	0.4	1,414
Female	42.7	2.3	25.5	1.3	26.4	2.2	3.9	0.4	0.7	0.2	0.8	0.2	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	44.6	2.5	26.1	1.1	24.6	2.2	3.1	0.2	0.8	0.2	0.8	0.2	2,763
Rural	40.1	4.2	31.0	3.2	25.4	3.1	3.3	0.8	0.2	0.1	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	38.1	3.6	27.4	2.0	28.8	2.3	3.7	0.8	1.5	0.7	0.4	0.4	417
Q2	40.9	3.2	31.2	3.4	23.6	2.3	3.3	0.9	0.7	0.3	0.3	0.2	528
Q3	45.2	4.2	21.7	1.7	29.7	4.4	2.4	0.7	0.5	0.4	0.6	0.3	601
Q4	42.3	2.8	29.4	1.9	23.0	2.0	3.9	0.6	0.4	0.2	1.0	0.2	651
Q5(richest)	51.3	2.7	24.3	1.1	20.2	1.6	2.8	0.8	0.8	0.3	0.7	0.2	750
M	42.0	6.0	18.1	3.9	32.6	5.4	0.0	0.0	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	46.3	3.6	24.1	1.6	25.0	2.5	3.4	1.0	0.5	0.4	0.7	0.2	793
30-44	41.5	3.1	28.7	1.1	26.0	3.1	2.1	0.6	0.5	0.2	1.2	0.3	811
45-59	44.2	2.6	26.3	1.9	24.0	2.7	3.5	0.7	1.1	0.3	1.0	0.4	636
60-69	43.0	2.0	24.7	1.8	25.1	1.8	5.9	1.9	1.3	0.7	0.0	0.0	337
70-79	45.0	2.6	31.5	2.4	22.3	3.1	0.8	0.5	0.3	0.1	0.0	0.0	268
80+	51.6	4.6	22.0	3.0	21.8	3.8	3.8	1.5	0.8	0.4	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	44.3	2.3	26.5	1.0	24.7	2.1	3.1	0.2	0.7	0.2	0.7	0.2	2,977

## 10.2 Inability to cope - q8001

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
<b>Gender</b>													
Male	42.0	2.5	27.6	1.6	25.5	1.9	3.6	0.3	0.6	0.2	0.6	0.4	1,414
Female	34.1	2.3	27.4	1.6	30.8	1.9	5.7	0.7	1.3	0.3	0.8	0.2	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	38.0	2.4	27.2	1.4	28.4	1.6	4.7	0.5	1.0	0.2	0.8	0.2	2,763
Rural	35.5	3.6	31.6	2.5	26.7	2.1	5.0	1.1	1.0	0.8	0.1	0.2	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	34.9	3.6	25.2	2.0	32.9	2.6	5.2	1.1	1.4	0.7	0.4	0.4	417
Q2	34.6	2.9	29.2	3.9	29.6	2.5	5.7	1.6	0.6	0.3	0.3	0.2	528
Q3	37.7	4.0	25.6	2.8	31.5	3.0	3.3	0.6	1.4	0.8	0.6	0.3	601
Q4	36.7	2.9	28.7	1.8	26.9	1.7	6.0	0.6	0.7	0.3	1.0	0.2	651
Q5(richest)	42.6	3.0	28.3	1.2	24.0	2.7	3.4	0.3	0.9	0.3	0.7	0.2	750
M	41.0	4.8	20.5	7.8	15.6	3.6	15.7	4.9	0.0	0.0	7.3	3.2	31
<b>Age</b>													
18-29	40.7	4.2	26.6	2.6	27.7	2.1	3.5	0.4	0.9	0.3	0.7	0.2	793
30-44	35.8	2.7	29.6	1.6	29.7	2.0	2.7	0.5	1.1	0.4	1.2	0.3	811
45-59	36.0	2.6	27.7	2.1	28.6	2.5	5.9	1.5	0.8	0.3	1.0	0.4	636
60-69	33.1	2.2	27.0	2.5	28.8	2.4	9.5	1.3	1.6	0.8	0.0	0.0	337
70-79	42.1	2.8	28.5	3.2	23.7	2.4	5.7	1.6	0.0	0.0	0.0	0.0	268
80+	44.5	4.4	18.4	3.6	30.5	4.4	4.9	2.0	1.7	0.9	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	37.8	2.2	27.5	1.3	28.3	1.5	4.7	0.4	1.0	0.2	0.7	0.2	2,977



### 10.3 Satisfaction with your health - q8002

	Very dissatisfied	SE	Dissatisfied	SE	Neither satisfied nor dissatisfied	SE	Satisfied	SE	Very Satisfied	SE	Missing	SE	N
<b>Gender</b>													
Male	2.4	0.5	2.8	0.4	11.6	1.6	59.0	2.0	23.1	3.2	1.1	0.3	1,414
Female	3.5	0.6	4.1	0.6	16.8	1.7	57.0	1.7	17.8	2.3	0.9	0.2	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	3.0	0.5	3.4	0.5	14.4	1.7	58.0	1.7	20.1	2.9	1.0	0.2	2,763
Rural	2.5	0.9	4.8	0.9	12.4	2.0	56.5	3.4	23.3	2.7	0.4	0.4	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	2.2	0.6	4.3	1.1	16.0	2.1	53.6	2.5	23.1	2.7	0.9	0.5	417
Q2	3.5	1.0	5.4	1.0	15.2	3.1	56.3	2.8	19.2	3.8	0.3	0.2	528
Q3	3.0	0.9	4.2	0.8	12.1	2.2	61.1	1.7	18.7	3.2	0.9	0.4	601
Q4	1.8	0.6	2.1	0.4	14.9	3.0	60.4	2.0	19.1	3.1	1.6	0.5	651
Q5(richest)	4.1	0.6	2.4	0.2	12.8	0.8	57.6	3.3	22.5	3.8	0.7	0.2	750
M	0.0	0.0	3.5	3.2	41.5	8.0	38.7	6.8	9.0	3.1	7.3	3.2	31
<b>Age</b>													
18-29	4.1	0.9	2.5	1.1	10.2	0.8	52.2	3.0	30.0	2.6	1.1	0.4	793
30-44	2.6	0.5	2.1	0.6	10.8	1.4	61.5	2.7	21.7	3.3	1.4	0.4	811
45-59	2.5	0.5	5.5	1.0	17.3	2.6	55.4	1.7	17.9	3.3	1.4	0.4	636
60-69	2.9	2.1	6.3	0.9	17.3	2.6	60.0	2.2	13.3	2.3	0.1	0.1	337
70-79	1.3	0.4	2.7	0.6	20.9	2.3	67.5	3.2	7.7	2.9	0.0	0.0	268
80+	4.0	2.0	3.2	2.4	25.3	6.8	57.7	4.1	9.9	4.0	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	3.0	0.4	3.5	0.4	14.3	1.7	57.9	1.7	20.3	2.8	1.0	0.2	2,977

## 10.4 Feeling safe when alone at home - q8011

	Completely safe	SE	Very safe	SE	Moderately safe	SE	Slightly safe	SE	Not safe at all	SE	Missing	SE	N
<b>Gender</b>													
Male	6.4	0.7	23.7	1.7	40.4	2.3	23.1	2.1	5.8	0.8	0.6	0.2	1,414
Female	5.1	0.8	19.5	2.4	39.4	1.2	24.9	1.7	10.4	0.8	0.7	0.2	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	5.5	0.6	21.5	1.8	40.0	1.4	24.3	1.8	8.0	0.5	0.7	0.2	2,763
Rural	8.2	2.0	20.7	3.3	38.1	3.0	21.3	2.9	11.5	2.8	0.2	0.2	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	6.1	1.5	24.8	2.6	35.5	2.5	22.1	2.6	11.1	2.1	0.4	0.4	417
Q2	8.0	1.3	15.6	2.4	40.3	2.1	24.4	1.6	10.9	1.4	0.9	0.3	528
Q3	3.5	0.7	18.9	2.5	39.8	3.0	30.2	4.1	7.5	1.0	0.2	0.1	601
Q4	5.4	0.7	21.3	3.7	40.1	1.8	24.1	3.0	8.3	1.0	0.8	0.3	651
Q5(richest)	5.8	0.7	26.4	1.5	42.0	3.2	20.1	1.5	5.2	0.8	0.5	0.2	750
M	10.7	4.7	12.5	4.4	36.8	7.5	20.6	5.9	12.1	7.3	7.3	3.2	31
<b>Age</b>													
18-29	6.9	1.0	24.0	2.5	39.2	3.0	23.1	1.2	6.2	0.7	0.7	0.2	793
30-44	6.0	0.8	20.1	1.7	40.7	2.3	25.8	2.5	6.9	1.1	0.5	0.3	811
45-59	5.8	1.2	21.8	3.5	41.7	1.7	20.6	2.0	9.2	1.0	0.8	0.3	636
60-69	4.2	1.8	21.9	2.2	34.1	2.5	28.2	4.3	10.7	1.9	0.9	0.3	337
70-79	3.1	0.6	18.2	4.0	44.7	1.6	20.1	3.7	13.4	1.2	0.6	0.6	268
80+	5.4	1.8	18.9	7.7	34.8	3.7	33.6	6.2	7.3	2.8	0.0	0.0	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	5.7	0.6	21.5	1.7	39.9	1.3	24.1	1.8	8.2	0.5	0.6	0.2	2,977

## 10.5 Feeling safe when walking down the street alone after dark - q8012

	Completely safe	SE	Very safe	SE	Moderately safe	SE	Slightly safe	SE	Not safe at all	SE	Missing	SE	N
<b>Gender</b>													
Male	4.4	1.3	15.2	2.7	32.6	1.4	30.5	3.9	16.9	1.4	0.4	0.2	1,414
Female	1.8	0.9	8.2	1.8	24.4	4.3	38.0	2.9	27.0	3.8	0.6	0.2	1,563
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Residence</b>													
Urban	2.8	1.1	11.2	2.4	28.1	3.1	34.9	3.5	22.5	2.8	0.5	0.2	2,763
Rural	6.7	2.3	15.6	3.1	31.2	3.0	28.6	3.7	17.9	3.5	0.0	0.0	214
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
<b>Income</b>													
Q1(poorest)	4.8	2.0	17.0	2.6	27.3	4.1	27.3	3.1	23.2	4.8	0.4	0.4	417
Q2	6.0	1.9	11.4	2.0	26.2	2.0	31.3	4.4	24.8	3.4	0.3	0.1	528
Q3	1.9	0.7	8.1	2.8	25.9	4.2	41.0	2.9	22.5	4.8	0.6	0.3	601
Q4	3.2	1.2	13.0	3.3	26.1	4.7	36.3	4.2	20.8	4.0	0.5	0.2	651
Q5(richest)	0.9	0.4	10.0	1.6	34.5	4.2	33.8	3.2	20.4	2.5	0.4	0.1	750
M	0.0	0.0	10.9	3.6	20.9	6.0	31.4	6.9	29.5	8.9	7.3	3.2	31
<b>Age</b>													
18-29	3.1	1.2	13.0	3.1	32.6	1.7	33.3	4.2	17.5	1.6	0.5	0.2	793
30-44	3.3	1.2	10.9	1.4	30.7	4.6	34.6	3.6	20.1	3.5	0.4	0.2	811
45-59	4.1	1.6	11.3	2.9	29.6	2.6	32.8	2.9	21.2	3.6	1.0	0.4	636
60-69	2.7	1.5	12.7	1.6	20.1	4.5	34.3	4.3	30.2	3.2	0.0	0.0	337
70-79	1.0	0.4	10.8	4.5	23.6	4.3	34.3	3.2	30.2	6.1	0.0	0.0	268
80+	1.5	1.6	6.0	4.5	12.1	4.8	48.4	9.5	31.3	3.7	0.8	0.4	133
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	3.0	1.1	11.5	2.3	28.3	3.0	34.4	3.4	22.2	2.7	0.5	0.2	2,977