

REPORT OF NAMIBIA

TABLE OF CONTENTS

1	Introduction
2	Methodology
2.1	Sampling
2.2.1	Geographic Component of the Survey
2.2.2	The map
2.3	Survey Metrics
2.3.1	Sample deviation index
2.3.2	HH Sample deviation index
2.3.3	Response rate
2.3.4	Missing data
2.3.5	Reliability
3	Socio-Demographic Profile
3.1	Household composition by residence and age groups according to income quintile
3.2	Selected age distributions by income quintile
3.3	Characteristics of head of household by income quintile
3.4	Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex
3.5	Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education
4	Risk Factors
4.1	Prevalence of smoking and average daily tobacco consumption
4.2	Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers
4.3	Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

- 4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water
- 4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility
- 4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used
- 5 Coverage
 - 5.01 Tuberculosis
 - 5.02 HIV Prevention and Screening
 - 5.03 Antenatal Care and Mother to Child Transmission of HIV
 - 5.04 Birth Delivery Care
 - 5.05 Child's Health - Common Illness
 - 5.06 Child's Health - Danger Signs
 - 5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card
 - 5.08 Vaccination Status of Children 12-23 Months
 - 5.09 Women's Cancer Screening
 - 5.10 Oral Health and Injuries
 - 5.11 Self Reported Non-Communicable Conditions
 - 5.12 Mental Health Conditions
 - 5.13 Cataracts
- 6 Health State Descriptions
 - 6.01 In general, rate your health today - q2000
 - 6.02 Difficulty with 'work and household activities' in last 30 days - q2001
 - 6.03 Difficulty with 'moving around' in last 30 days - q2010
 - 6.04 Difficulty with 'vigorous activities' in last 30 days - q2011
 - 6.05 Difficulty with 'self-care' in last 30 days - q2020
 - 6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021

- 6.07 Amount of 'bodily aches or pains' in last 30 days - q2030
- 6.08 Amount of 'bodily discomfort' in last 30 days - q2031
- 6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050
- 6.10 Difficulty with 'learning a new task' in last 30 days - q2051
- 6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060
- 6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061
- 6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071
- 6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072
- 6.15 Problem with 'sleeping' in last 30 days - q2080
- 6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081
- 6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090
- 6.18 Problem with 'worry or anxiety' in last 30 days - q2091
- 6.19 Adjusted for vignette responses
- 7 Health System Responsiveness
 - 7.1 Self-reported utilization of health services and perceived unmet need
 - 7.2 Patient assessed characteristics of ambulatory health services
 - 7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1
 - 7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2
 - 7.4 Patient assessed characteristics of hospital inpatient health services
 - 7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1
 - 7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2
 - 7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"
- 8 Health Expenditure, Insurance and Human Resources for Health

- 8.0 Expenditure Background Indicators (2002)
- 8.1 Distribution of health payments across different population groups
- 8.2 Percent distribution of structure of out-of-pocket health payments
- 8.3 Financial sources used by households for paying for health services
- 8.4 Insurance Coverage
- 8.5 Human resources for health
- 9.1 Adult Mortality Chart 1
- 9.2 Adult Mortality Chart 2
- 9.3 Adult Mortality Chart 3
- 9.4 Adult Mortality Chart 4
- 10 Stress and Social Capital
- 10.1 Inability to control the important things in life - q8000
- 10.2 Inability to cope - q8001
- 10.3 Satisfaction with your health - q8002
- 10.4 Feeling safe when alone at home - q8011
- 10.5 Feeling safe when walking down the street alone after dark - q8012

1. Introduction

The World Health Organization initiated the World Health Survey (WHS) as a *data collection platform* to obtain information on the health of populations and on health systems. The WHS responds to the need of countries for a detailed and sustainable health information system and gathers data through surveys to measure essential population health parameters. The WHS uses standard survey procedures and instruments for general population surveys to gather comparable data across WHO Member States. These methods and instruments have been refined through scientific review of literature, extensive consultations with international experts and large-scale pilot tests. WHS is designed to evolve through its implementation by continuous input from collaborators, including policy makers, survey institutions, scientists and other interested parties. The data is jointly owned by the countries and the WHO, and there is a commitment for a long-term data collection, building local capacity and using this information for health policy.

The objectives of the World Health Survey were to:

- develop a **means** of providing:
 - **valid, relevant, reliable** and **comparable** information on **outcomes, functions** and **inputs** to health systems
 - to **supplement** the information provided by routine **HIS**
- build the **evidence base** necessary for policy-makers:
 - to **routinely monitor** if health systems are achieving the **desired goals**
 - and in some countries, to assess if the **increase in health resources** is having the desired effect

The introductory chapter of each country report will provide details on the current sources of data from routine health information systems and ad hoc surveys. It will describe the health situation in the country and identify gaps. It will also present a brief overview on the health system organization within the country.

2. Sampling

The WHO sampling guidelines emphasized scientific principles of survey sampling as explicit standards for quality, included examples of good sampling plans, and identified quality assurance standards for countries to adhere to. WHO and technical advisers provided technical support to countries when needed. The following principles were used for sampling in the WHS:

1. The sample should target the *de facto* population (i.e. all people living in that country including guest workers, immigrants and refugees) and not *de jure* population (i.e. the citizens of that country alone) with complete geographic coverage of the country, if possible, or a representative geographic segment of the country.
2. A *sample frame* with 90% coverage that included all key subgroups of interest. Enumeration or listing of households to update the frame was recommended.
3. The WHS sample identified all adult members of the general population aged 18 years or older as its sample. Households are selected using a random, stratified sampling procedure. One individual per household is then selected through a random selection procedure using the Kish table method.
4. The WHS used a sampling strategy which encompasses a known non-zero selection probability for any individual included in the study. This makes it possible to extrapolate the data to the whole population. Households and individuals are selected with a known probability in order to provide selection probabilities for each household and individual in the data set.
5. The WHS sampling strategy was without replacement and categories of non-respondents were clearly defined.
6. The WHS used GPS devices in certain countries and digitized maps to geocode the data within specified guidelines.

Each national report will present a detailed description of the sampling strategy and the allocation tables

2.2.1 Geographic Component of the Survey

Geographic Information refers to any information associated with a location on the Earth's surface. This location can be associated to a coordinate (latitude/longitude), a place's name, a postal address or any other geographic object.

Household Surveys represent an important source of information. If the geographic location of the surveyed clusters is collected at the same time that the interview is conducted users are then able to integrate the spatial dimension in their analysis

The geographic component of the WHO WHS is stored in two sections of the questionnaire:

- 0100 Sampling Information allowing the link between the location of each cluster and the sampling frame used for the survey. This section has also been used in order to collect the type of setting in which each household was located at the time of the survey. This section of the questionnaire has been used for all the countries part of the WHO WHS
- 0200 Geocoding Information for the collection of the latitude and longitude of each household. This section has only been used in 27 countries where GPS devices were available or have been provided. The choice of these countries has been done on the basis of the availability of digital maps delimiting the surveyed clusters, the field data collection planning and their distribution within the different WHO regions. Specific training and field material have been prepared and used by the interviewers or field supervisors in these countries in order to collect the latitude and longitude of each of the surveyed household.

As the sampling design may not always include the administrative structure of the country and as the codes entered in the data set for the sampling level needed to be labelled, a key correspondence table containing this information has been collected after the survey itself.

Once all the survey data and the key correspondence table available a specific protocol has been applied to the two sections mentioned earlier in order to clean eventual data collection or data entry mistakes as well as to fill the gaps in close collaboration with the survey institution.

The questionnaire as well as the training material and data cleaning protocols used can be downloaded from the WHO WHS web site (<http://www.who.int/healthinfo/survey/en/index.html>) in the "Survey Instrument and Related Documents" section.

Namibia is part of the 27 countries where Global Positioning System (GPS) devices have been used during the survey in order to collect the location of each surveyed household.

In order to take advantage of the level of precision given by the use of these devices, but also take into account country respective data confidentiality policies regarding the distribution of such data to third parties, several additional geographical variables have been calculated:

- the weighted centre of gravity of each cluster part of the survey.
- different parameters and indexes giving an indication of the dispersion of the households interviewed in the cluster around its centre of gravity. These parameters and indexes can be used to explain possible heterogeneity in the answers obtained for a same cluster to questions presenting a geography dimension (accessibility for example).

The final geographic subset for Namibia contains:

- the cleaned information stored in the section 0100 and 0200
- the labels attached to the codes used in the data set for identifying each level of the sampling when available
- the 1st and 2nd administrative level names and codes coming from the Second Administrative Level Boundaries data set project (SALB, http://www3.who.int/whosis/gis/salb/salb_home.htm) allowing the link with the administrative structure of the country (if not part of the sampling frame).
- the geographic variables described above

The use of the GPS devices offers a large range of application which goes from the production of thematic maps, through the performing of spatial analysis to modelling using the spatial dimension of some of the surveyed data.

The generation of the maps and/or application of the methods that are now briefly described may require specific Geographic Information System (GIS) software and technical expertise. In addition to that, in order to create thematic maps, extract figures from other layers or perform spatial analysis it is first necessary to make sure that all the digital layers used are homogeneous in terms of: projection, representativity and sometimes scale of representation. The first one insures the proper overlay of the different layers in the GIS; the second is necessary in order to make a correct analysis of the results; the last one becomes important when we try to attribute cluster to other geographical entities like administrative units for example. In any case this underlines the importance of giving preference to digital layers being well documented (meta information available).

The capacity that exist within a GIS to create thematic maps by overlaying different layers or by linking each feature to a specific data set is largely used nowadays. One example of such map is the sampling distribution map reported at the beginning of this report which has been generated overlaying the location of each surveyed cluster on top of the extension of the a population distribution grid with the indication of the delimitation of the administrative units.

As it can be seen it is not necessary to go down to the location of each household for generating a thematic map which already provides a lot of useful information. Depending on the variables it is in fact possible to generate thematic maps at any higher level part of the sampling frame or administrative structure of the country.

The other potential application is the possibility to extract new parameters from spatially distributed variables stored in other digital layers. The parameters generally used are either linked to:



- the location of the cluster itself (e.g altitude)
- the geographic relation (distance or travelling time) which exist between the location of the cluster and the location of other geographic objects (roads, health facility, populated place, water point,.).

These parameters are then used to aggregate the clusters according to new units of analysis integrating therefore the geographic dimension as one of the parameters.

The final example of application that can be mentioned regards the possibility to link the data collected in the context of the WHO WHS to other data collected at the same level (cluster) or any above aggregated level (districts for example) in order to look for possible relationships that would allow the creation of models. If applicable these models may generate figures for units (administrative, statistical) that would not have been integrated in the original sampling frame.

This section only represent a short introduction to the possibilities that the geographic dimension of the surveyed information can offer. A more detailed description of the methods used in order to generate maps, perform specific analysis or apply models are presented in a publication that can be downloaded from the WHO WHS web site (Ebener et al., forthcoming).

NAMIBIA WHS 2003 Sampling Distribution



Data Source:

- Cluster Location: collected with GPS devices during WHS 2003
- Administrative Boundaries: SALB Data set 2000
- Population figures: extracted from the Landscan database 2001


Map Projection: UN-projected (Geographic)
Reference System: WGS 84 datum

Map created 21 July 2004

The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.
© WHO 2004. All rights reserved




2.3.1 Sample deviation index



SDI

2.3.2 HH Sample deviation index



HDI

2.3.3 Response rate

Household

Selected	4,656
----------	-------

Interviewed	4,340
-------------	-------

Percentages	93
-------------	----


Individual

Selected	4,296
----------	-------

Interviewed	4,238
-------------	-------

Percentages	99
-------------	----

2.3.4 Missing data



2.3.5 Reliability

Insufficient / No data available

3.1 Household composition by residence and age groups according to income quintile

	Percent Rural	SE	Mean Number of									
			Household members	SE	Children < 2 years	SE	Children < 5years	SE	Children 5 -17 years	SE	Persons 18+ years	SE
Q1(poorest)	89.6	1.70	4.6	0.14	0.2	0.02	0.7	0.04	1.7	0.09	2.2	0.07
Q2	79.4	2.62	5.2	0.17	0.3	0.02	0.7	0.04	1.9	0.09	2.6	0.08
Q3	77.0	2.31	5.6	0.19	0.2	0.02	0.6	0.04	2.0	0.11	3.0	0.09
Q4	48.4	4.15	5.0	0.15	0.2	0.02	0.5	0.03	1.7	0.07	2.8	0.10
Q5(richest)	23.8	3.73	4.7	0.18	0.1	0.01	0.4	0.03	1.4	0.09	2.8	0.10
All	67.3	0.79	5.0	0.09	0.2	0.01	0.6	0.02	1.7	0.05	2.6	0.04

Note: table based on de jure members of household i.e. usual residents

3.2 Selected age distributions by income quintile

	Females Age 18–49	Persons Age 50+	Persons Age 60+
Q1(poorest)	890	639	413
Q2	955	595	411
Q3	1,131	575	382
Q4	859	335	191
Q5(richest)	856	334	155
All	4,691	2,478	1,553

Note: table based on de jure members of household i.e. usual residents

3.3 Characteristics of head of household by income quintile

	Mean Age of Head	SE	Median Education of Head	Percent Female Head	SE
Q1 (poorest)	50	0.88	1	52.0	2.51
Q2	49	0.99	2	47.4	2.21
Q3	49	1.28	2	44.6	1.99
Q4	43	0.85	3	44.9	2.46
Q5 (richest)	44	0.82	4	35.3	2.13
All	47	0.62	2	45.2	1.08

Education

1=no formal education

2=less than primary school

3=primary school completed

4=secondary school completed

5=high school (or equivalent) completed

6=college/pre-university/university completed

7=post graduate degree completed

Note: table based on de jure members of household i.e. usual residents

3.4 Household population: percent distribution of the de jure population by age, residence, marital status and education according to sex

	Male	SE	Female	SE	Missing	Total	SE
Age							
0-4	12.7	0.40	10.7	0.40		11.60	0.30
5-14	29.0	0.70	26.7	0.70		27.80	0.60
15-29	29.1	0.60	29.9	0.70		29.50	0.50
30-44	15.1	0.60	16.0	0.40		15.60	0.40
45-59	7.2	0.30	7.9	0.30		7.60	0.20
60-69	3.0	0.20	4.0	0.30		3.50	0.20
70-79	2.2	0.20	2.5	0.20		2.30	0.20
80+	1.3	0.10	2.0	0.20		1.70	0.10
Missing	0.4	0.10	0.3	0.10		0.30	0.10
Residence							
Urban	28.5	1.10	26.7	1.20		27.50	1.10
Rural	71.5	1.10	73.3	1.20		72.50	1.10
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Marital_Status							
Never	74.9	0.60	70.9	0.60		72.80	0.50
Currently	17.6	0.50	17.3	0.50		17.40	0.40
Separated	0.3	0.10	0.5	0.10		0.40	0.10
Divorced	0.4	0.10	1.1	0.10		0.80	0.10
Widowed	1.2	0.10	5.5	0.30		3.50	0.20
CoHabitng	2.4	0.20	2.3	0.20		2.30	0.20
Missing	3.2	0.50	2.5	0.40		2.80	0.40
Education							
NoFormal	29.3	0.90	29.1	1.00		29.20	0.90
LessthanPrimary	35.8	0.80	33.1	0.60		34.40	0.60
Primary	21.5	0.70	23.9	0.80		22.80	0.70
Secondary	9.2	0.50	10.2	0.50		9.70	0.50
HighSchool	0.1	0.00	0.1	0.00		0.10	0.00
College	1.6	0.20	1.7	0.20		1.70	0.20
PostGraduate	1.2	0.20	0.7	0.10		0.90	0.10
Missing	1.4	0.20	1.1	0.20		1.20	0.20
Total	100.0		100.0			100.00	
Number	9,346.0		10,866.0			20,212.00	
Number of Households						4,340.00	

3.5 Background characteristics of respondents: percent distribution of men and women by age, residence, marital status and education

	Male	SE	Female	SE	Missing	Total	SE
Age							
18-29	43.3	1.70	39.4	1.40		41.20	1.20
30-44	30.6	1.50	30.4	1.30		30.50	1.00
45-59	15.5	1.20	17.3	1.10		16.40	0.90
60-69	6.2	0.90	7.5	0.80		6.90	0.60
70-79	3.4	0.60	4.1	0.50		3.80	0.40
80+	1.0	0.30	1.3	0.20		1.10	0.20
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Residence							
Urban	35.7	1.60	30.5	1.40		33.00	1.10
Rural	64.3	1.60	69.5	1.40		67.00	1.10
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Marital_Status							
Never	57.6	1.70	52.1	1.60		54.70	1.30
Currently	34.4	1.50	30.2	1.40		32.20	1.10
Separated	0.8	0.20	1.4	0.30		1.10	0.20
Divorced	1.0	0.50	2.4	0.50		1.80	0.30
Widowed	1.5	0.30	9.5	0.90		5.70	0.50
CoHabitng	4.6	0.60	4.3	0.50		4.40	0.40
Missing	0.1	0.00	0.2	0.10		0.10	0.10
Education							
NoFormal	22.2	1.60	22.3	1.50		22.20	1.20
LessthanPrimary	25.7	1.60	25.1	1.40		25.40	1.10
Primary	28.5	1.70	31.2	1.30		29.90	1.10
Secondary	18.4	1.40	18.2	1.20		18.30	1.00
HighSchool	0.0	0.00	0.0	0.00		0.00	0.00
College	3.3	0.90	2.6	0.40		2.90	0.50
PostGraduate	1.8	0.50	0.7	0.20		1.20	0.30
Missing	0.0	0.00	0.0	0.00		0.00	0.00
Total	100.0		100.0			100.00	
Number	1,996.0		2,249.0			4,245.00	

4.1 Prevalence of smoking and average daily tobacco consumption

	Current daily smoker	SE	Smoker, not daily	SE	Non-smoker	SE	Missing	Average daily tobacco consumption*	SE	Number of respondents
Sex										
Male	20.4	1.50	5.1	0.70	64.4	1.80	10.0	7.0	0.50	1,996
Female	8.7	0.80	2.7	0.40	80.4	1.30	8.2	5.4	0.50	2,249
Missing										0
Residence										
Urban	14.5	1.30	3.0	0.60	74.2	1.60	8.3	8.0	0.70	1,399
Rural	14.1	1.20	4.2	0.50	72.2	1.70	9.5	5.8	0.50	2,846
Missing										0
Income Quintile										
Q1	21.9	2.20	6.1	1.20	66.4	2.70	5.6	5.6	1.00	821
Q2	13.8	1.70	3.4	0.70	76.4	2.20	6.4	5.9	0.80	843
Q3	11.8	1.50	3.8	0.90	73.5	2.10	10.9	5.2	1.00	938
Q4	9.4	1.40	3.5	1.00	73.7	2.70	13.4	7.6	0.80	699
Q5	14.2	1.70	3.0	1.10	74.5	2.30	8.3	10.1	1.00	726
Missing										219
Age										
18-29	7.6	0.80	3.4	0.60	77.7	1.50	11.3	6.2	0.70	1,751
30-44	15.6	1.60	3.9	0.80	71.3	1.90	9.2	7.1	0.50	1,294
45-59	21.6	2.20	2.6	0.90	68.8	2.60	7.0	7.4	1.10	697
60-69	22.7	3.60	5.5	1.50	66.5	3.90	5.3	5.0	0.70	293
70-79	22.2	4.40	7.7	2.90	68.9	5.50	1.3	5.5	1.50	160
80+	30.1	5.90	13.6	4.20	51.3	7.50	5.0	2.4	0.60	49
Missing										0
All	14.2	0.90	3.8	0.40	72.9	1.20	9.1	6.5	0.40	4,245

* Denominator used is current daily smoker

4.2 Prevalence of life time abstainers, non-heavy drinkers, infrequent heavy drinkers and frequent heavy drinkers

	Life time abstainers	SE	Non-heavy drinkers	SE	Infrequent Heavy Drinkers*	SE	Frequent Heavy drinkers**	SE	Missing	Number of respondents
Sex										
Male	44.7	1.80	35.5	1.70	7.7	0.90	1.6	0.40	10.5	1,996
Female	63.6	1.40	24.4	1.20	3.3	0.60	0.4	0.10	8.3	2,249
Missing										0
Residence										
Urban	57.9	1.90	25.5	1.60	7.0	0.90	0.8	0.20	8.8	1,399
Rural	53.2	1.70	31.7	1.50	4.5	0.70	1.0	0.30	9.6	2,846
Missing										0
Income Quintile										
Q1	51.4	2.20	34.7	2.10	7.1	1.20	1.0	0.40	5.8	821
Q2	56.5	2.60	31.1	2.60	4.9	1.20	0.6	0.40	7.0	843
Q3	54.7	2.90	29.3	2.40	3.5	0.90	1.1	0.60	11.4	938
Q4	54.5	2.70	25.6	2.50	6.2	1.40	0.4	0.20	13.3	699
Q5	56.9	2.60	27.6	2.30	5.8	1.10	1.1	0.50	8.6	726
Missing										219
Age										
18-29	56.3	1.90	27.3	1.60	4.5	0.80	0.8	0.30	11.1	1,751
30-44	50.3	1.90	30.7	1.80	8.0	1.40	1.4	0.50	9.6	1,294
45-59	54.9	2.60	32.0	2.60	5.5	1.30	0.4	0.20	7.1	697
60-69	56.8	4.20	32.7	3.80	1.6	0.80	1.7	0.90	7.2	293
70-79	67.2	5.00	28.9	4.90	1.4	0.90	0.0	0.00	2.5	160
80+	62.6	6.80	32.3	6.70	0.0	0.00	0.0	0.00	5.0	49
Missing										0
All	54.7	1.30	29.6	1.10	5.3	0.50	0.9	0.20	9.3	4,245

*Infrequent heavy drinkers: 1-2 days with 5 or more standard drinks per week(last 7 days)

**Frequent heavy drinkers: 3 or more days with 5 or more standard drinks per week(last 7 days)

4.3 Prevalence of insufficient intake of fruit and vegetables and insufficient physical activity

	Fruit and vegetable intake					Physical Activity					Number of respondents
	*Insufficient	SE	Sufficient	SE	Missing	**Insufficient	SE	Sufficient	SE	Missing	
Sex											
Male	89.0	1.00	11.0	1.00	0.0	19.2	1.10	70.8	1.70	10.0	1,996
Female	89.6	0.90	10.4	0.90	0.0	31.6	1.40	60.5	1.50	7.9	2,249
Missing											0
Residence											
Urban	81.8	1.50	18.2	1.50	0.0	27.8	1.50	63.9	1.50	8.3	1,399
Rural	92.9	0.90	7.1	0.90	0.0	24.8	1.30	66.1	1.70	9.1	2,846
Missing											0
Income Quintile											
Q1	95.7	1.20	4.3	1.20	0.0	26.4	2.10	68.6	2.30	5.0	821
Q2	92.5	1.30	7.5	1.30	0.0	28.8	2.80	65.1	2.90	6.1	843
Q3	91.2	1.40	8.8	1.40	0.0	23.1	2.00	65.9	2.30	11.0	938
Q4	88.1	1.60	11.9	1.60	0.0	24.4	2.10	62.4	2.90	13.2	699
Q5	76.2	2.80	23.8	2.80	0.0	28.7	2.40	63.2	2.40	8.1	726
Missing											219
Age											
18-29	89.9	1.10	10.1	1.10	0.0	22.3	1.60	66.8	2.00	10.9	1,751
30-44	86.9	1.30	13.1	1.30	0.0	22.8	1.50	68.2	1.80	9.1	1,294
45-59	88.3	1.80	11.7	1.80	0.0	30.1	2.50	63.3	2.70	6.6	697
60-69	94.1	2.20	5.9	2.20	0.0	33.3	3.80	61.2	3.90	5.5	293
70-79	94.6	1.60	5.4	1.60	0.0	47.1	5.20	51.2	5.30	1.7	160
80+	95.0	2.80	5.0	2.80	0.0	51.8	6.40	43.1	6.00	5.0	49
Missing											0
All	89.3	0.80	10.7	0.80	0.0	25.8	1.00	65.4	1.30	8.8	4,245

*Insufficient intake of fruit/vegetables: less than 5 servings of fruit/vegetables per one typical day

**Insufficient physical activity: less than 150 minutes per week(last 7 days) spent on walking/moderate activity/vigorous activity

4.4 Access to improved drinking-water: percent distribution of households by type of access to drinking water

	Piped into households	SE	other improved sources	SE	Total improved sources	SE	No improved sources	SE	Missing	Number of households
Residence										
Urban	62.2	3.80	25.4	3.50	87.5	1.10	2.9	0.70	9.5	1,419
Rural	21.7	2.40	46.8	2.10	68.5	2.00	21.8	1.90	9.6	2,918
Missing										0
Income_Quintile										
Q1(poorest)	5.1	0.90	56.7	2.40	61.7	2.60	31.8	2.50	6.4	995
Q2	9.2	1.40	63.3	2.50	72.5	2.40	20.4	2.30	7.1	887
Q3	30.1	2.40	46.1	2.70	76.2	2.30	12.2	1.90	11.5	865
Q4	68.3	3.10	14.9	2.00	83.2	2.00	4.1	1.00	12.7	684
Q5(richest)	87.3	1.40	1.2	0.40	88.6	1.40	0.2	0.20	11.2	671
Missing										235
Total	35.0	2.00	39.8	1.80	74.7	1.40	15.7	1.20	9.6	4,338

other improved sources: public standpipe/protected tube well/borehole/protected dug well/protected spring/rainwater collection

no improved sources: unprotected well/unprotected spring/water taken directly from a pond or stream/tanker truck water/vendor provided water

4.5 Access to improved sanitation: percent distribution of households by type of sanitation facility

	Flush toilet to sewage system	SE	Other improved toilet facilities	SE	Total improved sanitation	SE	No improved sanitation	SE	Missing	Number of households
Residence										
Urban	50.0	3.60	9.4	1.30	59.3	3.70	32.0	3.70	8.7	1,419
Rural	10.9	2.10	6.7	1.00	17.6	2.40	73.0	2.40	9.4	2,918
Missing										0
Income_Quintile										
Q1(poorest)	0.3	0.20	0.8	0.30	1.1	0.40	92.4	1.20	6.6	995
Q2	2.8	0.90	3.5	0.80	6.3	1.20	86.9	1.70	6.8	887
Q3	10.2	1.70	11.8	1.60	22.0	2.10	66.6	2.40	11.4	865
Q4	49.5	3.40	16.2	2.20	65.7	3.00	22.4	2.20	11.9	684
Q5(richest)	77.5	1.90	9.8	1.30	87.2	1.50	3.0	0.80	9.8	671
Missing										235
Total	23.7	1.90	7.6	0.80	31.3	2.00	59.6	2.00	9.1	4,338

other improved sanitation facilities: connection to septic system/pour-flush latrine/covered dry latrine(with privacy) [provided facilities not shared]
no improved sanitation: uncovered dry latrine(without privacy)/bucket latrine/no facilities(open defecation)

4.6 Solid Fuel use for cooking: percent distribution of households by type of cooking fuel used

	Electricity, Gas	SE	Kerosene	SE	Solid fuels on improved stove	SE	Solid fuels on traditional stove	SE	Total solid fuel use	SE	Missing	Number of households
Residence												
Urban	62.9	3.30	5.2	1.50	0.1	0.10	21.9	2.80	22.0	2.80	9.9	1,419
Rural	13.3	2.50	1.1	0.40	0.5	0.10	75.8	2.60	76.3	2.70	9.3	2,918
Missing												0
Income_ Quintile												
Q1(poorest)	0.8	0.30	1.7	0.60	0.0	0.00	91.6	1.30	91.6	1.30	5.9	995
Q2	7.3	1.20	4.0	1.20	0.1	0.10	81.6	2.20	81.7	2.20	7.0	887
Q3	18.1	2.60	3.9	0.80	0.6	0.30	65.9	3.20	66.5	3.20	11.5	865
Q4	63.1	3.00	1.9	0.60	0.6	0.30	22.1	2.30	22.7	2.30	12.3	684
Q5(richest)	84.9	1.60	0.1	0.10	0.4	0.30	2.4	0.70	2.8	0.80	12.2	671
Missing												235
Total	29.6	2.00	2.4	0.60	0.3	0.10	58.2	2.00	58.5	2.00	9.5	4,338

5.01 Tuberculosis

		Tuberculosis			
	Total	Need	Coverage	SE	M
Sex					
Female	2,249	18.4	18.1	2.5	0.2
Male	1,996	16.2	25.8	3.5	0.0
Missing	0				
Residence					
Urban	1,399	12.9	30.3	3.3	0.0
Rural	2,846	19.5	18.6	2.9	0.1
Missing	0				
Income_Quintile					
Q1(poorest)	821	24.8	24.3	4.8	0.2
Q2	843	19.3	14.9	3.2	0.0
Q3	938	15.6	24.4	5.6	0.0
Q4	699	16.3	24.2	5.8	0.0
Q5(richest)	726	8.6	15.8	5.1	0.0
Missing	219				
Age					
18-29	1,751	17.8	18.8	3.2	0.0
30-44	1,294	14.4	21.3	3.6	0.0
45-59	697	16.4	23.0	4.8	0.3
60-69	293	22.4	27.1	7.6	0.0
70-79	160	29.7	32.3	11.0	0.8
80+	49	21.8	6.0	5.1	0.0
Missing	0				
Total	4,239	17.4	21.5	2.3	0.1

M = % Missing of Covered SE= Standard Error of % Covered

Tuberculosis Indicators:

"Need" = % of all respondents with self-reported continuous cough lasting 3 weeks, OR had blood in phlegm, OR had coughed blood in the last 12 months

"Coverage" = % of respondents with self reported symptoms who indicated receiving a TB screening test in the last 12 months.

5.02 HIV Prevention and Screening

	Total (persons of reproductive age 18-49)	Voluntary Counselling and Testing for HIV/AIDS				Condom Usage			
		Need	Coverage	SE	M	Need	Coverage	SE	M
Sex									
Females	1,727	64.3	15.5	1.4	25.5	31.2	48.6	3.1	2.3
Males	1,606	66.6	3.7	0.6	66.6	37.4	64.8	3.0	0.3
Missing									
Residence									
Urban	1,208	71.8	14.2	1.4	46.3	33.9	62.8	3.3	1.1
Rural	2,125	61.8	6.8	0.9	45.3	34.4	53.9	2.9	1.3
Missing									
Income_Quintile									
Q1(poorest)	610	72.0	5.9	1.5	45.0	33.2	40.6	6.1	0.5
Q2	641	65.4	8.8	1.4	44.3	37.1	54.1	5.1	1.8
Q3	735	60.0	7.1	1.9	46.5	38.2	56.9	5.0	1.8
Q4	562	61.6	9.9	1.5	47.6	33.8	62.2	4.8	1.8
Q5(richest)	615	70.3	16.8	2.5	45.3	29.3	77.6	4.4	0.1
Missing	169								
Total	3,333	65.4	9.7	0.8	45.7	34.2	57.1	2.2	1.2

M = % Missing of Covered SE= Standard Error of % Covered

HIV Prevention and Screening Indicators:

"Total" = number of respondents of reproductive age.

"Voluntary Counselling and Testing for HIV/AIDS Need" = % of respondents of reproductive age who are sexually active, i.e. have had sexual intercourse in the last 12 months.

"Voluntary Counselling and Testing for HIV/AIDS Coverage" = % of respondents of reproductive age who are sexually active who indicated having taken an HIV test and receiving the results of test.

"Condom Usage Need" = % of respondents of reproductive age whose last sexual encounter during the last 12 months is classified as risky sexual behavior, i.e. having had sexual intercourse with a non-cohabitating partner .

"Condom Usage Coverage" = % of respondents of reproductive age who used a condom last time had risky sex during the last 12 months (with non-cohabitating partner).

5.03 Antenatal Care and Mother to Child Transmission of HIV

	Antenatal Care											Maternal to Child HIV Trans.			
	Women 18-49	Birth in last 5 years	Cov	SE	Visit 1x	Visit 2x	Visit 3x	Blood pressure checked	Blood sample taken	Told Signs of Pregnancy Complications	M	Birth in last 2 years	Cov	SE	M
Residence															
Urban	598	36.2	44.8	3.3	73.8	70.6	67.3	75.6	74.9	57.0	19.6	16.5	30.5	4.2	9.9
Rural	1,129	40.4	37.6	2.9	72.7	70.1	67.2	73.1	71.5	57.2	17.1	18.3	22.9	4.0	9.1
Missing															
Income_Quintile															
Q1(poorest)	339	50.0	29.0	3.6	69.4	66.9	62.2	68.2	61.8	49.9	17.4	21.0	13.3	5.8	9.4
Q2	345	41.8	40.6	4.9	70.7	67.2	64.3	73.8	73.3	56.2	20.3	21.7	24.3	6.8	12.2
Q3	361	33.5	47.2	6.2	80.0	77.4	76.6	73.3	80.6	61.6	14.2	13.6	19.3	7.6	8.7
Q4	295	37.6	47.2	5.3	76.7	73.3	68.4	77.1	73.6	59.3	18.1	20.7	39.9	7.2	8.4
Q5(richest)	297	30.4	47.4	5.2	73.6	71.8	71.8	80.7	79.9	64.2	19.8	11.2	39.0	9.1	9.8
Missing	89														
Total	1,727	39.0	39.9	2.3	73.0	70.3	67.2	73.9	72.6	57.1	17.9	17.6	25.4	3.0	9.4

M = % Missing of Covered SE= Standard Error of %Covered

Antenatal Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least 3 times or more by health care professional, received blood pressure, blood sample taken, and told signs about complications for their last pregnancy (aggregate antenatal care indicator).

"Visit 1x" "Visit 2x" "Visit 3x" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having been seen at least once, twice or three times or more respectively by a health care professional during their last pregnancy.

"Blood pressure checked" = % of female respondents of reproductive age with live birth in the last 5 years who had blood pressure taken when they saw health care professional during their last pregnancy.

"Blood sample taken" = % of female respondents of reproductive age with live birth in the last 5 years who had a blood sample taken when they saw health care professional during their last pregnancy.

"Told Signs of Pregnancy Complications" = % of female respondents of reproductive age with live birth in the last 5 years who had been told signs of pregnancy complications when they saw health care professional during their last pregnancy.

Maternal to Child HIV Transmission Indicators:

"Birth in last 2 years" = % of female respondents of reproductive age indicating having had a live birth in the last 2 years.

"Cov" = % of female respondents of reproductive age with live birth in the last 2 years who indicated having been counseled and tested and given results for HIV during their last pregnancy.

5.04 Birth Delivery Care

	Delivery Care					Place of Delivery						
	Women 18-49	Need	Coverage	SE	M	Public Health Facility	SE	Private Health Facility	SE	Home	SE	M
Residence												
Urban	598	36.2	81.6	2.4	12.1	68.0	3.1	9.9	2.3	5.0	1.4	17.2
Rural	1,129	40.4	72.9	3.1	7.6	64.6	3.1	4.8	1.3	20.5	2.7	10.1
Missing												
Income_Quintile												
Q1(poorest)	339	50.0	59.2	5.4	11.0	58.9	5.3	0.3	0.3	28.0	4.8	12.7
Q2	345	41.8	78.1	4.2	7.5	67.7	5.7	5.1	2.6	16.3	3.9	10.9
Q3	361	33.5	82.8	4.4	4.4	76.1	5.1	1.3	0.8	13.7	4.5	8.9
Q4	295	37.6	84.5	3.5	8.9	73.7	4.3	5.1	2.3	7.3	3.1	13.8
Q5(richest)	297	30.4	79.8	4.5	15.7	51.3	5.8	27.1	5.1	4.7	2.8	16.9
Missing	89											
Total	1,727	39.0	75.7	2.3	9.0	65.7	2.3	6.4	1.2	15.5	1.9	12.3

M = % Missing of Covered SE= Standard Error of % Covered Birth Delivery Care Indicators:

Total = Number of female respondents of reproductive age (18-49 years old)

"Birth in Last 5 Years" = % of female respondents of reproductive age indicating having had a live birth in the last 5 years.

"Coverage" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having received assistance with birth delivery from a health care professional (doctor, nurse or midwife) during their last birth.

"Public Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a public health facility.

"Private Health Facility" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered in a private health facility.

"Home" = % of female respondents of reproductive age with live birth in the last 5 years who indicated having their last birth delivered home or in the field (etc.)

5.05 Child's Health - Common Illness

	Total Children under Age of 5	Fever	More liquid	SE	Same or more food	SE	M	Cough	More liquid	SE	Same or more food	SE	M	Diarrhea	More liquid	SE	Same or more food	SE	M
Sex																			
Females	921	32.8	23.1	3.7	52.8	3.5	3.6	27.6	25.2	4.5	53.9	32.8	6.6	11.2	22.4	5.3	52.1	6.5	3.5
Males	760	39.0	22.3	3.2	55.7	3.4	4.4	34.5	21.4	3.5	55.8	39.0	3.9	16.7	29.2	4.6	51.9	5.3	7.3
Missing	76																		
Residence																			
Urban	466	38.6	23.7	3.8	61.9	4.4	1.5	31.5	23.8	5.1	62.5	38.6	2.6	14.2	24.6	4.2	49.4	5.9	2.2
Rural	1,291	34.8	22.6	3.2	51.7	3.0	5.1	30.5	23.7	3.5	52.4	34.8	6.0	13.4	28.3	4.7	52.3	5.4	6.6
Missing																			
Income_Quintile																			
Q1(poorest)	383	39.3	20.0	5.2	53.8	4.3	3.2	30.1	18.4	5.4	51.1	39.3	2.8	16.9	32.5	7.8	46.3	8.0	5.6
Q2	439	28.7	15.0	3.9	52.9	5.6	5.2	25.5	18.1	5.2	51.4	28.7	7.9	10.2	28.9	7.5	50.5	8.3	0.0
Q3	384	36.0	29.0	5.5	50.9	6.7	5.3	34.3	32.8	6.4	55.6	36.0	6.9	17.1	25.8	7.3	55.5	8.8	5.0
Q4	258	37.7	23.8	5.5	46.4	7.0	6.6	30.2	22.3	6.1	42.8	37.7	7.8	12.2	21.5	8.7	41.0	11.9	13.2
Q5(richest)	207	39.9	27.5	6.9	67.0	7.1	0.0	35.7	23.3	7.0	68.7	39.9	0.0	9.8	28.8	10.5	64.2	10.5	0.0
Missing	86																		
Total	1,758	35.8	22.9	2.5	54.6	2.5	4.1	30.7	23.7	2.9	55.2	35.8	5.0	13.6	27.3	3.6	51.5	4.3	5.4

M = % Missing of Covered SE= Standard Error of % Covered

Child's Common Illness Indicators:

Total = Number of children under the age of 5 years at the time of the interview.

"Fever"/"Cough"/"Diarrhea" = % of children under the age of 5 with reported episode of (cough, fever, or diarrhea) in the last 3 months.

"More Liquid" = % of children under 5 with either fever, cough, or diarrhea in the last 3 months, that were given more liquids to drink during their illness.

"Same or More Food" = % of children under 5 with either fever, cough, or diarrhea in the last 3 months, that were given the same amount or more food to eat during their illness.

5.06 Child's Health - Danger Signs

	Children under 5	Fast Breath	Seeking Treatment						Vomit	Seeking Treatment						No Drink or Eat	Seeking Treatment					
			Yes	SE	Public	Private	Trad	M		Yes	SE	Public	Private	Trad	M		Yes	SE	Public	Private	Trad	M
Sex																						
Females	921	0.1	90.2	3.8	91.4	6.6	0.7	0.7	7.2	93.4	2.9	91.3	4.1	2.1	1.0	10.4	93.2	3.4	83.7	13.5	1.5	0.7
Males	760	0.1	80.3	5.8	88.0	5.3	3.2	2.2	10.2	90.7	4.0	87.5	8.0	0.8	0.7	14.0	86.3	4.7	86.4	6.2	1.2	2.6
Missing	76																					
Residence																						
Urban	466	0.1	87.4	5.2	81.0	19.0	0.0	4.4	9.6	97.8	1.6	77.7	14.5	2.8	0.0	12.4	95.8	2.7	74.4	21.0	1.4	0.0
Rural	1,291	0.1	83.6	5.0	92.2	2.6	2.3	1.5	8.5	90.3	3.3	95.4	1.9	0.6	1.1	11.9	86.8	3.8	90.3	4.6	1.3	2.2
Missing																						
Income_Quintile																						
Q1(poorest)	383	0.1	85.1	6.3	98.4	1.6	0.0	0.0	13.6	88.4	4.7	98.8	0.0	1.2	0.0	13.9	87.3	5.4	93.6	0.5	2.4	0.0
Q2	439	0.1	82.2	6.6	94.3	2.9	0.0	0.0	5.7	95.4	4.7	100.0	0.0	0.0	0.0	7.7	88.4	7.0	98.5	0.0	0.0	0.0
Q3	384	0.1	85.1	6.6	90.5	2.1	4.9	4.1	9.5	90.6	5.1	92.5	1.8	0.0	1.8	12.2	91.1	4.5	86.3	9.3	0.0	2.6
Q4	258	0.1	77.3	12.0	83.2	15.9	0.0	6.1	6.5	94.4	4.0	85.8	12.8	0.0	3.4	15.0	86.3	9.3	93.3	6.0	0.0	1.5
Q5(richest)	207	0.0	100.0	0.0	63.6	28.0	8.4	0.0	7.1	100.0	0.0	67.1	24.2	8.7	0.0	11.6	100.0	0.0	44.3	45.7	5.8	0.0
Missing	86																					
Total	1,758	0.1	84.5	4.0	89.5	6.5	1.8	2.2	8.8	92.5	2.4	90.0	5.8	1.3	0.8	12.0	89.3	3.0	85.7	9.4	1.3	1.6

M = % Missing of those Seeking Treatment SE= Standard Error of % Seeking Treatment

Child Danger Signs Indicators:

Total = Number of children under the age of 5 years at the time of the interview.

"Fast Breathing"/"Vomiting"/"No Drink or Eat" = % of children under the age of 5 with reported episodes of (fast breathing, vomiting, or no eating or drinking) in the last 3 months.

"Seeking Treatment" = % of children under 5 with episodes of fast breathing, vomiting, or no eating or drinking in the last 3 months that were taken for care/treatment at either a private health facility, public health facility, or traditional healer (other).

Note: large number of missings are usually due to not having information type of facility (either a no answer or a don't know answer).

5.07 Vaccinations Meeting WHO Timeline Recommendations for Children 12 to 23 months of Age with Vaccination Card

	Children 12-23 months with card	DPT 3 Immunization			Measles		
		%Cov	SE	M	%Cov	SE	M
Sex							
Females	162	57.9	5.4	0.0	50.3	5.2	0.0
Males	138	46.2	5.7	0.0	47.2	6.1	0.0
Missing	17						
Residence							
Urban	75	55.6	5.8	0.0	48.6	5.4	0.0
Rural	241	50.9	4.9	0.0	49.2	4.8	0.0
Missing	0						
Income_Quintile							
Q1(poorest)	89	54.6	6.4	0.0	51.6	6.6	0.0
Q2	77	53.1	8.0	0.0	48.1	7.8	0.0
Q3	68	53.4	8.7	0.0	39.8	8.0	0.0
Q4	41	53.1	10.0	0.0	59.0	10.0	0.0
Q5(richest)	20	22.5	8.5	0.0	39.0	11.5	0.0
Missing	21						
Total	316	52.0	4.0	0.0	49.1	3.9	0.0

M = % Missing of Covered SE= Standard Error of % Covered

Vaccinations Meeting WHO Timeline for Children 12 - 23 months Indicators:

Total = Children 12 - 23 months with a vaccination card seen by the interviewer (does not include children less than 12 months to account for censored observations).

"DPT 3" = % of children 12 - 23 months with a vaccination card seen by the interviewer who received 3 inoculations of DTP not sooner than 6, 10 and 14 weeks respectively and before 12 months of age, and that was verified by the interviewer as being recorded on the card.

"Measles" = % of children 12 - 23 months with a vaccination card seen by the interviewer who received who received 1 inoculation of measles not sooner than 8 months of age and before 12 months of age.

5.08 Vaccination Status of Children 12-23 Months

	Children 12-23 months	Card seen			DPT 3 Immunization							Measles						
		Seen	SE	M	Total		Card		No Card		M	Total		Card		No Card		M
					%Cov	SE	%Cov	SE	%Cov	SE		%Cov	SE	%Cov	SE	%Cov	SE	
Sex																		
Females	239	67.7	3.9	0.0	70.6	3.8	88.2	3.4	11.8	3.4	20.4	77.2	3.7	77.5	3.6	22.5	3.6	14.0
Males	198	69.5	4.6	0.0	71.0	4.3	89.5	3.7	10.5	3.7	15.1	77.8	4.4	78.1	4.8	21.9	4.8	13.0
Missing	26																	
Residence																		
Urban	124	60.4	4.7	0.0	60.5	5.5	85.5	3.6	14.5	3.6	20.9	71.6	4.9	73.5	4.8	26.5	4.8	15.8
Rural	339	71.3	3.4	0.0	74.2	3.4	90.2	2.9	9.8	2.9	15.9	78.9	3.1	79.7	3.4	20.3	3.4	12.0
Missing	0																	
Income_Quintile																		
Q1(poorest)	124	72.1	5.7	0.0	69.3	6.6	96.7	1.9	3.3	1.9	23.2	72.3	5.9	83.1	5.3	16.9	5.3	20.0
Q2	105	73.8	5.6	0.0	79.6	5.3	87.6	5.4	12.4	5.4	12.4	88.8	3.8	77.9	6.0	22.1	6.0	6.7
Q3	98	69.3	7.5	0.0	76.8	6.9	84.0	6.9	16.0	6.9	12.6	78.5	5.8	81.4	6.9	18.6	6.9	9.1
Q4	55	74.0	6.4	0.0	76.3	6.6	92.7	4.2	7.3	4.2	7.8	88.4	4.0	78.8	6.8	21.2	6.8	5.7
Q5(richest)	49	41.0	8.5	0.0	39.8	9.1	73.4	9.2	26.6	9.2	31.5	50.7	9.1	54.4	10.7	45.6	10.7	26.9
Missing	32																	
Total	463	68.4	2.8	0.0	70.5	2.9	89.1	2.4	10.9	2.4	17.3	77.0	2.7	78.1	2.9	21.9	2.9	13.0

M = % Missing (% without adequate information to estimate Total DPT3 or Measles Coverage)

SE= Standard Error of % Total, DPT3 or Measles Coverage

Explanation of Indicators for Vaccination Status of Children 12 - 23 Months:

"Card Seen" = % of children 12 - 23 months of age who had a vaccination card seen by the interviewer.

"DPT 3 Immunization": does not account for WHO timeline recommendations.

"Total" = % of all children 12 - 23 months of age with DPT3 vaccination either verified by card or by recall of respondent.

"Card" = % of all DPT3 vaccinations among children 12-23 months that were verified by card only.

"No Card" = % of all DPT3 vaccinations among children 12-23 months that were based on recall of the respondent only.

"Measles": does not account for WHO timeline recommendations.

"Total" = % of all children 12 - 23 months of age with measles vaccination either verified by card or by recall of respondent.

"Card" = % of all Measles vaccinations among children 12-23 months that were verified by card only.

"No Card" = % of all Measles vaccinations among children 12-23 months that were based on recall of the respondent only.

5.09 Women's Cancer Screening

	Total Females	Percent Females 18+	PAP Smear	SE	M	Percent Females 40+	Mammography / Breast Exam	SE	M
Residence									
Urban	687	97.9	17.8	1.4	13.8	26.4	15.2	2.9	18.3
Rural	1,563	93.2	5.8	0.9	18.2	35.7	6.7	1.2	14.6
Missing	0								
Income_Quintile									
Q1(poorest)	482	90.7	6.6	1.6	10.2	37.9	8.2	2.3	10.8
Q2	449	95.0	4.2	1.0	13.9	32.3	1.5	1.1	17.8
Q3	482	95.7	6.3	1.3	24.9	35.2	7.5	2.9	21.4
Q4	355	94.7	10.8	1.9	17.6	24.3	12.4	4.0	14.1
Q5(richest)	359	98.6	23.0	2.4	12.3	32.5	19.3	4.3	11.1
Missing	124								
Age									
18-29	887	39.4	8.3	1.3	18.8				
30-39	504	22.4	13.8	1.8	14.2				
40-49	336	14.9	13.3	2.2	13.3	14.9	12.9	2.1	13.0
50-59	232	10.3	5.2	1.6	15.3	10.3	6.9	2.2	15.1
60-69	169	7.5	3.0	1.2	23.4	7.5	3.3	1.5	21.0
70+	121	5.4	0.0	0.0		5.4	0.0	0.0	
Missing	0								
Total	2,249	94.6	9.6	0.8	16.8	32.8	8.8	1.2	15.5

M = % Missing of those with Pap Smear/Mammography SE= Standard Error of % with Pap Smear/Mammography

***Note: Values in the age row under the "Females 18 plus" and "Females 40 plus" columns are column percent values.

Women's Cancer Screening Indicators:

"Papsmear need" = % of all female respondents aged 18-69

"Papsmear" coverage = % of those female respondents aged 18-69 who self-reported receiving a pap smear test during pelvic examination in the last 3 years.

"Mammography need" = % of all female respondents aged 40-69

"Mammography" coverage = % of those female respondents aged 40-69 who self-reported receiving a mammography test or breast examination in the last 3 years.

****In several countries women above age 69 were asked the cancer screening questions, and for this reason they are included in the table. For countries that did not ask this questions of women above age 69, these values are expected to be missing.

5.10 Oral Health and Injuries

	Oral Health Problems					Road Traffic Accidents				Injury			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
Sex													
Female	2,249	23.9	53.4	2.8	8.3	1.4	45.0	10.0	12.4	3.2	37.8	6.4	6.0
Male	1,996	13.6	52.2	4.2	3.2	3.0	46.4	9.8	2.9	4.7	31.3	7.2	2.5
Missing	0												
Residence													
Urban	1,399	17.9	61.4	3.1	3.1	2.3	48.7	9.5	9.3	5.3	35.6	5.2	5.1
Rural	2,846	19.7	49.2	3.4	8.1	2.1	44.5	9.3	4.5	3.2	33.0	8.2	3.2
Missing	0												
Income_ Quintile													
Q1(poorest)	821	21.7	44.8	4.5	7.1	1.9	20.7	13.9	16.4	4.0	32.7	13.8	0.0
Q2	843	19.9	56.6	6.1	8.3	1.7	53.4	19.2	0.0	2.8	35.7	12.6	0.0
Q3	938	17.6	44.1	6.5	10.9	2.8	53.5	16.4	1.9	4.6	38.6	11.5	6.4
Q4	699	18.2	51.3	5.3	4.4	1.6	28.5	13.7	12.0	3.4	24.7	7.8	1.8
Q5(richest)	726	17.8	70.1	5.0	1.6	2.8	49.3	13.0	6.7	5.6	36.0	7.4	8.8
Missing	219												
Age													
18-29	1,751	13.6	60.0	4.0	2.9	2.1	3.5	10.3	3.5	4.6	3.7	7.0	3.7
30-44	1,294	19.5	59.6	3.9	6.6	2.1	4.7	10.9	4.7	3.3	5.2	6.8	5.2
45-59	697	22.7	37.8	5.6	9.4	2.2	3.0	17.5	3.0	3.1	6.9	13.4	6.9
60-69	293	30.5	46.8	7.2	10.6	2.5	0.0	18.9	0.0	4.3	0.0	9.9	0.0
70-79	160	35.9	48.2	8.1	7.4	2.7	58.2	31.6	58.2	4.5	0.0	18.9	0.0
80+	49	28.4	46.6	13.6	6.0	0.3	100.0	0.0	100.0	0.0	0.0	0.0	0.0
Missing	0												
Total	4,239	19.1	53.0	2.6	6.5	2.1	46.0	6.9	6.2	3.9	34.1	5.1	4.1

M = % Missing of % Covered SE= Standard Error of % Covered

Indicators:

Oral Health:

"Need" = % of all respondents indicating any problems in the last 12 months with their mouth and/or teeth.

"Coverage" = % of respondents reporting problems with their mouth and teeth who indicated receiving dental or medical care in last 12 months.

Road Traffic Accident:

"Need" = % of all respondents indicating having suffered bodily injury from a road traffic accident in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury from a road traffic accident in last 12 months who indicating having received treatment for their injuries on-site or at health facility within 1 hour of accident .

Injuries from Non-traffic Accidents:

"Need" = % of all respondents indicating having suffered bodily injury due to non-traffic accidents in last 12 months.

"Coverage" = % of respondents reporting having suffered bodily injury due to non-traffic accidents in last 12 months who indicating having received treatment for injuries on-site or at health facility within 1 hour of accident

5.11 Self Reported Non-Communicable Conditions

	Angina					Osteo-Arthritis				Asthma				Diabetes			
	Total	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M	Need	Coverage	SE	M
Sex																	
Female	2,249	8.4	41.1	4.0	0.2	9.6	46.3	4.4	0.2	4.2	64.3	5.9	0.0	2.6	66.5	8.6	1.6
Male	1,996	5.4	40.7	7.1	0.0	8.1	55.7	5.8	0.0	2.1	43.8	10.3	0.5	1.0	67.6	9.7	0.0
Missing	0																
Residence																	
Urban	1,399	6.3	51.3	6.1	0.5	8.2	53.7	4.9	0.0	4.2	54.9	5.7	0.0	2.0	66.4	7.9	1.0
Rural	2,846	7.3	36.6	4.8	0.0	9.3	48.8	5.0	0.1	2.8	60.2	8.6	0.3	1.8	67.0	9.6	1.3
Missing	0																
Income_Quintile																	
Q1(poorest)	821	11.9	27.0	5.2	0.0	9.3	36.3	6.4	0.5	3.4	55.5	12.7	0.0	1.9	71.6	14.9	0.0
Q2	843	7.3	38.6	9.3	0.0	8.1	39.4	8.2	0.0	2.6	83.9	9.5	0.0	1.9	45.2	15.5	4.2
Q3	938	5.6	53.8	9.0	0.0	11.4	58.0	7.9	0.0	2.6	50.0	14.2	0.0	1.8	78.9	10.3	1.6
Q4	699	4.9	52.6	10.0	0.0	8.9	64.1	7.3	0.0	4.5	50.0	12.2	0.7	1.4	73.3	12.2	0.0
Q5(richest)	726	4.9	50.0	11.0	1.3	6.6	50.7	8.2	0.0	3.6	56.5	9.2	0.0	2.5	65.3	9.8	0.0
Missing	219																
Age																	
18-29	1,751	4.6	41.6	7.3	0.0	3.7	53.1	7.6	0.0	1.6	50.7	10.3	0.0	0.6	65.5	14.9	0.0
30-44	1,294	6.0	39.0	7.1	0.0	7.4	46.7	5.8	0.0	4.2	53.1	9.0	0.4	1.6	53.6	11.5	1.3
45-59	697	8.3	43.1	9.1	0.8	15.4	45.8	7.8	0.0	4.6	64.0	14.0	0.0	2.5	84.7	8.7	0.0
60-69	293	18.7	46.4	8.5	0.0	19.2	56.3	8.0	0.0	6.5	67.6	12.6	0.0	7.7	70.4	10.4	3.0
70-79	160	13.5	23.9	9.9	0.0	24.3	62.1	9.6	1.0	1.6	100.0	0.0	0.0	4.6	51.5	20.5	0.0
80+	49	7.4	55.0	16.6	0.0	30.3	40.1	7.9	0.0	2.5	43.4	34.7	0.0	1.0	73.3	26.4	0.0
Missing	0																
Total	4,239	7.0	41.0	3.8	0.2	8.9	50.3	3.8	0.1	3.2	58.0	5.5	0.2	1.9	66.8	6.8	1.2

M = % Missing for Covered

SE= Standard Error of %Covered

Self Reported Non-Communicable Conditions Indicators:

"Need" = % of total respondents indicating having received a diagnosis of the condition (angina, arthritis, asthma, diabetes) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (angina, arthritis, asthma, diabetes) indicating they have received treatment or screening for the condition in the last 2 weeks.

5.12 Mental Health Conditions

	Total	Depression				Schizophrenia			
		Need	Coverage	SE	M	Need	Coverage	SE	M
Sex									
Female	2,249	8.8	30.9	4.5	0.2	2.8	66.8	7.2	0.4
Male	1,996	4.9	30.1	6.7	0.4	2.5	55.1	11.1	7.6
Missing	0								
Residence									
Urban	1,399	5.4	27.8	4.7	0.6	2.4	64.4	11.9	11.5
Rural	2,846	7.8	31.6	5.3	0.2	2.8	60.5	7.2	0.3
Missing	0								
Income Quintile									
Q1(poorest)	821	10.6	34.9	7.2	0.4	5.5	56.9	9.0	0.0
Q2	843	8.6	31.4	8.8	0.0	2.5	76.7	11.9	0.0
Q3	938	5.4	19.9	9.4	0.9	2.5	68.5	14.7	0.0
Q4	699	3.6	26.0	11.9	0.0	1.6	70.5	19.0	0.0
Q5(richest)	726	6.1	35.6	8.0	0.0	1.5	35.6	22.6	37.2
Missing	219								
Age									
18-29	1,751	5.8	27.9	6.4	0.5	2.4	60.1	11.3	9.5
30-44	1,294	5.3	30.7	6.7	0.0	2.7	73.6	9.1	0.0
45-59	697	10.0	33.6	9.3	0.0	2.9	64.7	17.9	0.0
60-69	293	8.2	38.3	16.5	0.0	2.8	58.2	18.7	0.0
70-79	160	17.5	29.1	17.3	1.3	4.1	8.2	9.0	0.0
80+	49	11.3	13.4	10.1	0.0	2.6	41.4	29.5	0.0
Missing	0								
Total	4,239	7.0	30.6	4.1	0.3	2.7	61.6		3.6

M = % Missing for Covered

SE= Standard Error of %Covered

Mental Health Conditions Indicators:

"Need" = % of total respondents indicating having received a diagnosis of the condition (depression, schizophrenia) in last 12 months.

"Coverage" = % of respondents with a self-reported condition (depression, schizophrenia) indicating they have received treatment or screening for the condition in the last 2 weeks.

5.13 Cataracts

		Cataracts			
	Total 60+	Need	Coverage	SE	M
Sex					
Female	290	9.2	42.3	10.5	2.1
Male	213	11.0	68.5	14.8	0.0
Missing	0				
Residence					
Urban	82	17.3	75.8	9.5	0.0
Rural	420	8.6	46.0	13.0	1.6
Missing	0				
Income_Quintile					
Q1(poorest)	110	4.9	53.2	19.7	0.0
Q2	126	12.6	34.1	17.9	0.0
Q3	123	6.6	59.9	17.7	0.0
Q4	72	15.7	78.5	14.4	5.0
Q5(richest)	41	15.4	67.5	14.1	0.0
Missing	31				
Age					
60-69	293	8.2	46.0	12.5	2.3
70-79	160	13.1	59.5	15.3	0.0
80+	49	10.3	73.9	13.3	0.0
Missing	0				
Total	503	10.0	54.5	9.7	1.1

M = % Missing of Covered SE= Standard Error of %Covered

Cataracts Indicators:

Total = Number of respondents 60 years of age or more.

"Need" = % in need (respondents 60 years of age or more with self-reported diagnosis of cataracts in the last 5 years).

"Coverage" = % respondents 60 years of age or more with self-reported diagnosis of cataracts having indicated receiving eye surgery to remove cataracts in the last 5 years)

6.01 In general, rate your health today - q2000

	Very Good	SE	Good	SE	Moderate	SE	Bad	SE	Very Bad	SE	Missing	SE	N
Gender													
Male	40.6	1.6	29.1	1.8	16.2	1.3	3.4	0.6	0.7	0.2	10.0	1.3	1,993
Female	33.0	1.4	29.2	1.3	23.4	1.1	5.5	0.6	1.3	0.4	7.6	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	42.1	1.6	28.2	1.6	17.1	1.4	4.1	0.7	0.7	0.2	7.8	1.0	1,397
Rural	33.8	1.5	29.6	1.6	21.5	1.1	4.8	0.6	1.2	0.3	9.2	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	28.4	2.5	32.6	2.8	25.9	2.0	7.2	1.2	0.7	0.2	5.2	1.1	820
Q2	37.6	2.6	25.4	2.8	24.1	2.1	4.7	1.0	1.8	0.7	6.4	1.4	842
Q3	36.8	2.2	28.1	2.5	17.9	2.2	4.7	1.0	1.6	0.5	10.9	1.8	937
Q4	39.0	2.6	26.2	2.0	18.9	1.8	3.0	0.6	0.5	0.2	12.4	2.3	698
Q5(richest)	42.9	2.5	33.5	2.3	12.3	1.7	2.8	0.8	0.5	0.4	8.0	1.5	725
M	33.2	4.8	29.8	4.8	20.5	3.4	3.4	1.2	0.3	0.2	12.7	3.0	219
Age													
18-29	45.1	1.7	28.3	1.8	12.8	1.4	2.8	0.5	0.2	0.1	10.8	1.3	1,748
30-44	38.4	1.8	31.1	1.8	17.1	1.3	3.6	0.7	0.9	0.3	8.9	1.4	1,292
45-59	27.1	2.3	29.7	2.3	28.8	2.3	6.1	1.2	2.1	0.9	6.1	1.2	696
60-69	19.1	3.0	23.9	3.5	38.3	4.1	10.8	2.1	3.0	1.2	4.9	1.8	293
70-79	9.7	2.8	29.9	4.6	44.2	5.2	11.7	3.3	1.0	1.0	3.4	2.1	160
80+	6.9	2.8	30.5	5.1	42.7	6.2	9.3	2.9	4.3	2.1	6.3	2.6	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	36.6	1.1	29.1	1.2	20.0	0.9	4.5	0.4	1.0	0.2	8.7	0.8	4,239

6.02 Difficulty with 'work and household activities' in last 30 days - q2001

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	54.9	1.8	14.8	1.1	10.7	1.0	6.9	0.8	1.6	0.5	11.1	1.4	1,993
Female	49.4	1.7	17.8	1.2	15.0	1.0	8.0	0.8	1.2	0.3	8.6	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	62.7	1.4	13.5	1.0	10.4	1.0	4.0	0.6	1.1	0.4	8.2	1.0	1,397
Rural	46.7	1.7	17.9	1.1	14.3	0.9	9.1	0.9	1.5	0.4	10.6	1.3	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	42.4	2.5	23.6	1.7	15.2	1.6	11.9	1.7	0.9	0.3	6.0	1.2	820
Q2	50.3	2.7	16.8	1.6	15.5	1.8	9.2	1.5	1.6	0.6	6.6	1.4	842
Q3	50.7	2.4	12.9	1.8	13.5	1.6	6.9	1.4	2.5	0.9	13.5	2.1	937
Q4	54.5	2.5	13.2	1.9	13.0	1.9	4.5	1.0	1.5	0.8	13.4	2.3	698
Q5(richest)	66.0	2.6	14.4	1.6	7.4	1.2	3.2	0.8	0.4	0.2	8.6	1.5	725
M	44.8	4.4	20.1	3.9	11.6	2.8	10.0	2.6	0.4	0.3	13.0	3.0	219
Age													
18-29	58.3	2.0	14.6	1.3	9.2	1.0	5.3	1.0	0.6	0.3	12.0	1.4	1,748
30-44	55.1	2.1	16.4	1.4	11.9	1.1	6.4	1.0	0.9	0.3	9.3	1.4	1,292
45-59	47.0	2.7	17.5	1.9	16.2	1.9	9.0	1.3	2.3	1.1	8.0	1.7	696
60-69	36.1	3.7	18.9	3.1	23.0	3.4	13.6	2.7	3.3	1.3	5.1	1.8	293
70-79	21.0	4.8	25.2	4.5	26.7	3.9	17.7	4.3	3.8	2.2	5.6	2.4	160
80+	9.2	3.7	21.6	5.1	29.6	5.8	21.5	4.2	7.9	4.0	10.2	4.4	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	52.0	1.3	16.4	0.8	13.0	0.7	7.5	0.6	1.4	0.3	9.8	0.9	4,239

6.03 Difficulty with 'moving around' in last 30 days - q2010

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	63.3	1.8	14.3	1.2	7.2	0.8	4.2	0.6	1.1	0.4	9.9	1.3	1,993
Female	59.4	1.5	15.5	1.1	11.6	1.0	5.4	0.6	0.7	0.2	7.4	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	70.8	1.5	11.3	1.2	6.7	0.7	2.6	0.4	0.8	0.3	7.9	1.0	1,397
Rural	56.6	1.8	16.7	1.1	11.0	1.0	6.0	0.6	0.9	0.3	8.9	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	51.7	2.5	22.1	2.5	13.7	1.4	6.7	1.1	1.1	0.6	4.8	1.1	820
Q2	61.1	2.8	16.2	1.8	10.0	1.6	5.5	1.0	1.3	0.6	6.0	1.4	842
Q3	58.7	2.7	15.0	1.6	10.0	1.5	4.8	1.1	0.8	0.4	10.7	1.8	937
Q4	62.9	2.6	12.3	1.9	8.4	1.4	3.1	0.8	0.9	0.5	12.4	2.3	698
Q5(richest)	74.6	2.3	8.8	1.4	5.3	1.1	2.7	0.8	0.4	0.2	8.2	1.6	725
M	58.5	4.5	12.0	3.0	8.2	3.2	8.6	2.8	0.0	0.0	12.7	3.0	219
Age													
18-29	69.3	2.0	10.8	1.1	5.9	1.0	2.7	0.6	0.6	0.4	10.6	1.3	1,748
30-44	66.1	2.0	13.9	1.3	8.1	0.8	2.2	0.4	0.7	0.3	9.0	1.4	1,292
45-59	53.7	2.5	17.5	1.8	12.6	1.6	8.9	1.6	1.1	0.7	6.1	1.2	696
60-69	37.5	3.4	27.7	3.6	19.9	3.2	9.2	2.3	0.7	0.4	4.9	1.8	293
70-79	24.3	4.6	28.2	4.3	25.5	3.9	18.6	3.9	2.1	1.8	1.3	0.9	160
80+	14.9	3.6	30.5	6.3	19.4	5.2	22.6	4.2	7.5	4.0	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	61.2	1.3	14.9	0.8	9.5	0.7	4.9	0.4	0.9	0.2	8.6	0.8	4,239

6.04 Difficulty with 'vigorous activities' in last 30 days - q2011

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	59.8	1.8	12.3	1.3	9.3	1.0	5.2	0.8	3.2	0.6	10.2	1.4	1,993
Female	54.3	1.6	13.3	1.0	10.1	0.8	9.9	0.9	4.7	0.7	7.7	0.8	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	68.2	1.6	8.3	0.9	8.4	0.9	5.1	0.7	2.0	0.4	7.9	1.0	1,397
Rural	51.3	1.8	15.1	1.1	10.4	0.8	8.9	0.8	5.0	0.7	9.3	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	49.4	2.7	16.1	1.9	13.5	1.7	9.3	1.2	6.6	1.3	5.2	1.1	820
Q2	54.6	2.8	15.3	1.8	9.8	1.3	7.9	1.3	6.2	1.3	6.2	1.4	842
Q3	56.0	2.6	10.8	1.7	9.5	1.5	9.5	1.7	3.0	0.8	11.2	2.1	937
Q4	57.9	2.7	10.8	1.6	8.9	1.5	7.1	1.2	2.5	0.8	12.8	2.4	698
Q5(richest)	69.9	2.6	10.1	1.6	7.0	1.4	3.6	0.8	1.1	0.3	8.2	1.6	725
M	50.8	4.6	16.0	4.0	8.4	3.8	7.7	1.9	4.1	1.6	12.9	3.0	219
Age													
18-29	64.4	1.9	10.6	1.1	8.2	0.9	3.9	0.7	1.7	0.5	11.1	1.3	1,748
30-44	59.6	1.9	13.4	1.4	9.7	1.1	6.0	0.8	2.3	0.5	9.0	1.4	1,292
45-59	52.7	2.7	13.8	1.7	11.4	1.5	9.8	1.5	5.9	1.3	6.5	1.3	696
60-69	35.7	3.6	17.3	3.4	13.5	2.7	20.1	3.3	8.3	1.8	5.1	1.8	293
70-79	21.8	3.9	18.6	4.0	13.9	3.3	28.1	5.3	16.4	4.2	1.3	0.9	160
80+	16.6	4.5	19.6	5.2	4.9	2.3	16.1	4.6	37.8	6.7	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	56.9	1.3	12.9	0.8	9.7	0.6	7.7	0.6	4.0	0.5	8.9	0.9	4,239

6.05 Difficulty with 'self-care' in last 30 days - q2020

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	70.0	1.9	10.4	1.1	4.4	0.7	4.3	0.8	0.9	0.4	9.8	1.3	1,993
Female	68.8	1.4	10.0	0.8	7.5	0.8	5.3	0.6	0.9	0.3	7.6	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	78.2	1.5	6.8	0.8	4.1	0.7	2.3	0.4	0.8	0.3	7.8	1.0	1,397
Rural	65.0	1.5	11.9	0.9	7.0	0.7	6.1	0.8	0.9	0.3	9.1	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	56.8	2.5	16.0	1.7	11.6	1.5	9.4	1.5	1.2	0.5	5.0	1.1	820
Q2	71.3	2.1	12.8	1.8	5.3	1.3	3.5	0.9	1.2	0.6	6.0	1.4	842
Q3	69.1	2.6	8.9	1.6	5.4	1.3	4.9	1.3	1.0	0.5	10.7	1.8	937
Q4	72.2	2.9	7.0	1.1	3.5	0.7	3.6	1.1	1.0	0.5	12.7	2.4	698
Q5(richest)	82.3	2.6	4.4	1.1	3.4	1.0	1.8	1.1	0.2	0.1	8.0	1.5	725
M	59.0	5.5	13.3	4.4	8.1	2.6	6.2	2.7	0.7	0.7	12.9	3.0	219
Age													
18-29	72.7	1.8	7.5	1.0	3.8	0.7	4.2	0.9	1.0	0.4	10.9	1.3	1,748
30-44	74.1	1.8	8.9	1.1	5.2	0.9	2.4	0.5	0.6	0.2	8.9	1.4	1,292
45-59	63.9	2.8	12.1	1.6	8.2	1.7	8.7	1.5	0.9	0.5	6.1	1.2	696
60-69	58.1	3.7	19.0	3.0	11.6	2.2	5.6	1.8	0.5	0.4	5.1	1.8	293
70-79	45.9	4.9	24.6	4.5	13.3	3.2	12.0	3.6	2.9	2.1	1.3	0.9	160
80+	50.5	5.8	15.3	4.0	21.2	6.0	5.3	2.2	2.6	1.2	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	69.4	1.1	10.2	0.6	6.1	0.5	4.8	0.6	0.9	0.2	8.6	0.9	4,239

6.06 Difficulty with 'taking care of and maintaining general appearance' in last 30 days - q2021

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	69.0	2.0	10.9	1.3	6.1	1.0	3.3	0.6	0.5	0.3	10.1	1.3	1,993
Female	67.4	1.5	10.7	0.8	7.5	0.8	6.1	0.7	0.7	0.2	7.6	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	76.7	1.6	7.5	0.9	3.8	0.6	3.4	0.5	0.5	0.2	8.0	1.0	1,397
Rural	64.0	1.8	12.4	1.0	8.3	1.0	5.5	0.6	0.7	0.2	9.2	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	56.1	2.4	16.3	1.9	12.7	1.8	8.0	1.2	1.3	0.6	5.5	1.1	820
Q2	68.8	2.2	12.9	1.6	7.4	1.2	4.1	1.0	0.5	0.3	6.1	1.4	842
Q3	67.5	3.0	9.6	1.5	6.8	1.3	4.5	1.0	0.6	0.5	10.9	1.8	937
Q4	69.3	2.9	9.4	1.5	3.8	0.8	4.4	1.1	0.5	0.4	12.5	2.3	698
Q5(richest)	82.6	2.6	4.6	1.1	2.6	0.8	2.2	0.7	0.0	0.0	8.0	1.5	725
M	62.5	5.1	11.3	3.8	6.4	1.8	6.5	2.6	0.7	0.7	12.7	3.0	219
Age													
18-29	72.2	2.0	7.7	0.9	4.9	1.0	3.7	0.7	0.8	0.4	10.9	1.3	1,748
30-44	73.3	2.0	8.6	1.1	6.2	1.3	2.7	0.5	0.1	0.1	9.1	1.4	1,292
45-59	61.2	3.1	15.1	2.0	8.0	1.8	8.4	1.5	0.7	0.4	6.6	1.3	696
60-69	56.2	3.7	21.4	2.9	10.8	2.2	5.6	1.6	1.0	0.6	4.9	1.8	293
70-79	46.9	5.3	21.1	3.6	15.2	3.6	13.7	3.7	1.9	1.8	1.3	0.9	160
80+	32.1	5.7	20.0	4.0	26.2	5.4	15.6	4.9	1.1	0.6	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	68.2	1.3	10.8	0.8	6.9	0.7	4.8	0.5	0.6	0.2	8.8	0.8	4,239

6.07 Amount of 'bodily aches or pains' in last 30 days - q2030

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	53.3	1.9	17.7	1.2	12.9	1.1	5.2	0.9	1.0	0.4	9.9	1.3	1,993
Female	47.3	1.4	18.0	1.2	16.4	1.2	8.8	0.9	1.8	0.4	7.6	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	58.8	1.7	16.8	1.3	9.5	0.8	6.0	0.8	1.1	0.4	7.8	1.0	1,397
Rural	45.9	1.6	18.4	1.2	17.3	1.2	7.7	1.0	1.6	0.3	9.1	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	39.7	2.7	22.1	2.0	19.7	2.2	11.2	1.8	2.3	0.9	4.8	1.1	820
Q2	49.8	3.0	19.1	2.0	16.6	1.7	6.8	1.5	1.0	0.4	6.6	1.4	842
Q3	50.6	2.5	14.3	2.0	17.3	1.9	5.0	1.0	2.0	0.6	10.8	1.8	937
Q4	50.7	2.8	16.7	1.7	11.4	1.6	7.2	1.3	1.6	0.7	12.4	2.3	698
Q5(richest)	62.6	2.6	16.1	1.9	7.7	1.1	5.3	1.1	0.3	0.2	8.0	1.5	725
M	45.8	5.0	21.6	5.1	11.6	3.7	7.9	2.2	0.4	0.4	12.7	3.0	219
Age													
18-29	57.9	1.9	15.6	1.2	10.7	1.2	4.4	0.8	0.8	0.3	10.6	1.3	1,748
30-44	55.7	1.9	17.0	1.4	10.5	1.1	6.8	1.5	1.0	0.3	9.0	1.4	1,292
45-59	40.2	2.7	20.8	2.2	20.0	2.4	10.7	1.8	1.5	0.8	6.8	1.4	696
60-69	28.7	3.3	23.1	3.8	31.9	3.5	8.4	2.0	3.0	1.3	4.9	1.8	293
70-79	13.5	3.4	23.3	5.1	36.2	5.5	18.1	4.0	7.7	3.6	1.3	0.9	160
80+	19.1	4.5	28.9	4.9	21.3	4.3	22.4	4.8	2.7	1.2	5.5	2.6	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	50.2	1.2	17.9	0.9	14.7	0.9	7.1	0.7	1.4	0.3	8.7	0.9	4,239

6.08 Amount of 'bodily discomfort' in last 30 days - q2031

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	54.2	2.0	17.0	1.4	12.9	1.3	4.4	0.7	1.1	0.4	10.3	1.3	1,993
Female	48.6	1.4	19.5	1.2	15.9	0.9	7.6	0.8	1.1	0.2	7.4	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	60.3	1.8	16.2	1.2	10.1	1.0	4.4	0.7	0.9	0.4	8.2	1.0	1,397
Rural	46.8	1.8	19.4	1.3	16.6	1.1	6.9	0.8	1.2	0.3	9.1	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	40.0	2.6	26.6	2.2	18.2	1.8	9.0	1.6	1.4	0.7	4.9	1.1	820
Q2	49.3	2.8	19.6	1.9	16.5	1.9	7.5	1.5	1.0	0.4	6.1	1.4	842
Q3	52.0	2.9	15.3	2.0	16.7	2.0	4.1	0.9	1.2	0.5	10.7	1.8	937
Q4	54.2	2.8	15.3	1.7	11.3	1.6	5.5	1.1	1.3	0.7	12.4	2.3	698
Q5(richest)	64.1	2.7	14.6	1.7	7.8	1.1	3.9	0.9	0.9	0.7	8.7	1.5	725
M	45.0	4.9	18.0	4.4	15.3	4.2	7.3	2.2	0.4	0.4	14.0	3.2	219
Age													
18-29	59.4	2.0	14.8	1.3	10.6	1.3	3.5	0.6	0.8	0.4	10.9	1.3	1,748
30-44	57.1	1.9	17.0	1.4	10.7	1.2	5.5	1.0	0.6	0.3	9.1	1.4	1,292
45-59	39.0	2.8	24.3	2.5	20.5	2.3	8.3	1.6	1.4	0.7	6.4	1.3	696
60-69	32.2	3.4	23.6	3.0	27.1	3.1	9.7	2.4	2.5	1.1	4.9	1.8	293
70-79	10.7	2.7	30.6	5.9	35.7	5.5	18.3	4.5	3.4	2.1	1.3	0.9	160
80+	23.7	6.0	24.0	5.6	21.9	5.4	20.5	4.3	4.4	2.4	5.5	2.6	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	51.2	1.3	18.3	1.0	14.5	0.8	6.1	0.6	1.1	0.3	8.8	0.8	4,239

6.09 Difficulty with 'concentrating or remembering things' in last 30 days - q2050

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	63.4	1.8	14.6	1.1	6.1	0.8	5.0	0.9	1.1	0.4	9.8	1.3	1,993
Female	57.5	1.4	18.8	1.0	9.0	0.8	6.0	0.7	1.2	0.3	7.5	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	69.9	1.4	11.8	1.0	5.3	0.7	4.4	0.6	0.7	0.3	7.9	1.0	1,397
Rural	55.5	1.7	19.3	1.1	8.8	0.9	6.1	0.9	1.4	0.4	8.9	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	50.6	2.6	24.5	1.6	13.2	1.6	5.9	1.2	0.8	0.4	4.9	1.1	820
Q2	59.2	2.4	18.0	1.7	7.6	1.2	7.2	1.5	2.0	0.8	6.0	1.4	842
Q3	59.6	2.4	15.2	1.6	7.6	1.2	5.8	1.2	1.1	0.6	10.7	1.8	937
Q4	62.3	2.9	14.4	1.7	4.4	1.0	5.6	1.2	0.7	0.4	12.5	2.3	698
Q5(richest)	71.5	2.2	11.8	1.6	4.4	0.8	3.3	0.9	1.0	0.7	8.1	1.5	725
M	60.0	4.4	14.9	3.9	7.4	2.2	3.9	1.3	1.1	0.8	12.7	3.0	219
Age													
18-29	64.3	1.9	14.6	1.2	5.5	1.1	3.7	0.8	1.2	0.4	10.7	1.3	1,748
30-44	65.9	1.8	14.0	1.2	6.5	0.9	3.9	0.7	0.7	0.3	9.0	1.4	1,292
45-59	54.6	2.7	22.3	2.2	8.4	1.5	7.7	1.7	0.9	0.5	6.1	1.2	696
60-69	45.2	3.8	24.8	3.4	15.4	2.6	7.2	2.0	2.5	1.5	4.9	1.8	293
70-79	30.6	6.0	24.5	3.8	18.8	3.7	22.3	5.3	2.5	1.9	1.3	0.9	160
80+	33.3	6.2	20.5	6.1	19.7	4.5	19.4	4.6	1.6	1.0	5.5	2.6	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	60.3	1.2	16.8	0.8	7.6	0.6	5.5	0.6	1.2	0.3	8.6	0.8	4,239

6.10 Difficulty with 'learning a new task' in last 30 days - q2051

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	65.4	1.6	15.3	1.2	4.6	0.6	3.7	0.5	1.0	0.4	10.0	1.3	1,993
Female	60.0	1.5	16.6	1.2	8.7	0.9	5.6	0.6	1.7	0.4	7.5	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	73.4	1.6	9.2	1.0	5.2	0.7	3.5	0.5	0.9	0.3	7.9	1.0	1,397
Rural	57.2	1.6	19.4	1.2	7.5	0.8	5.3	0.6	1.6	0.4	9.1	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	53.1	2.3	23.9	2.0	10.1	1.4	6.3	1.1	1.7	0.6	5.0	1.2	820
Q2	61.7	2.3	16.8	2.0	6.3	1.2	5.8	1.1	3.1	1.0	6.2	1.3	842
Q3	60.0	2.4	16.8	2.0	7.3	1.5	4.8	0.8	0.4	0.2	10.8	1.8	937
Q4	65.9	2.9	11.1	1.8	5.8	1.1	4.2	0.8	0.6	0.4	12.4	2.3	698
Q5(richest)	76.5	2.2	8.8	1.4	3.7	0.8	2.0	0.5	0.8	0.7	8.2	1.5	725
M	54.7	3.9	19.9	3.5	6.4	2.1	4.1	1.8	1.6	0.8	13.3	3.0	219
Age													
18-29	67.3	1.7	14.5	1.4	4.4	0.6	2.6	0.6	0.7	0.3	10.6	1.3	1,748
30-44	67.6	1.9	13.0	1.3	5.9	0.9	3.7	0.6	0.6	0.2	9.2	1.3	1,292
45-59	60.6	2.8	18.8	2.2	8.7	1.4	5.2	1.3	0.6	0.3	6.1	1.2	696
60-69	39.4	3.8	26.5	3.3	14.3	2.6	9.6	2.0	5.0	1.8	5.3	1.8	293
70-79	32.5	5.6	25.9	4.7	14.8	3.8	16.6	4.2	7.9	3.9	2.3	1.3	160
80+	22.2	5.4	16.6	4.8	13.9	4.0	29.8	6.0	12.0	4.6	5.5	2.6	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	62.5	1.2	16.0	0.9	6.7	0.6	4.7	0.4	1.3	0.3	8.7	0.8	4,239

6.11 Difficulty with 'personal relationships or participation in the community' in last 30 days - q2060

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	52.5	2.1	11.5	1.3	7.5	1.0	2.8	0.6	14.3	1.5	11.4	1.4	1,993
Female	48.0	1.8	16.7	1.3	9.0	0.8	4.4	0.5	13.0	1.3	9.0	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	48.9	2.4	8.5	0.7	5.2	0.8	2.5	0.4	24.6	2.6	10.2	1.0	1,397
Rural	50.7	2.0	17.1	1.1	9.7	0.8	4.2	0.6	8.2	1.2	10.1	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	50.6	2.7	20.9	2.0	11.2	1.5	5.3	1.2	5.8	1.7	6.3	1.2	820
Q2	54.8	2.7	17.4	1.8	7.4	1.2	4.6	0.9	8.7	1.5	7.0	1.4	842
Q3	46.7	2.8	15.9	2.0	10.5	1.6	4.8	1.1	10.1	1.5	12.0	1.8	937
Q4	49.1	3.4	7.8	1.3	8.4	1.6	2.0	0.7	18.5	2.8	14.3	2.6	698
Q5(richest)	49.9	3.6	7.4	1.4	3.3	1.1	1.3	0.4	27.5	3.5	10.6	1.6	725
M	49.1	5.1	13.5	4.0	6.9	2.3	2.5	1.3	14.6	3.8	13.4	3.0	219
Age													
18-29	54.2	2.2	12.8	1.4	7.8	1.0	2.6	0.6	10.8	1.2	11.8	1.3	1,748
30-44	48.4	2.2	14.4	1.2	6.1	0.8	2.5	0.6	18.0	2.0	10.6	1.4	1,292
45-59	49.9	2.8	15.2	2.1	8.3	1.5	3.2	1.0	15.6	2.1	7.9	1.4	696
60-69	48.5	3.9	18.4	3.4	10.4	2.1	6.1	1.8	8.8	2.2	8.0	2.0	293
70-79	32.0	5.2	14.4	2.8	24.4	4.6	18.4	4.7	7.1	2.1	3.7	1.6	160
80+	21.5	5.1	22.4	4.6	14.0	3.7	16.7	4.7	18.8	4.9	6.6	2.7	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	50.1	1.6	14.3	0.8	8.3	0.6	3.7	0.4	13.6	1.2	10.1	0.9	4,239

6.12 Difficulty with 'dealing with conflicts and tensions' in last 30 days - q2061

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	54.8	2.1	10.8	1.1	5.0	0.8	2.3	0.7	2.3	0.5	24.8	1.9	1,993
Female	51.6	1.8	14.2	1.0	7.1	0.8	3.6	0.6	2.9	0.5	20.6	1.6	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	47.8	2.4	10.1	0.9	5.4	0.8	2.0	0.5	2.6	0.6	32.1	2.6	1,397
Rural	55.7	2.0	13.8	1.0	6.4	0.8	3.5	0.6	2.6	0.5	17.9	1.8	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	57.3	2.7	17.5	1.7	6.7	1.2	4.0	1.0	3.3	1.0	11.2	2.1	820
Q2	57.6	3.0	14.7	1.7	5.6	1.0	3.1	0.8	3.7	1.1	15.4	2.1	842
Q3	52.1	2.7	12.4	1.4	9.3	1.5	3.3	1.4	1.5	0.6	21.4	2.5	937
Q4	48.9	3.4	9.3	1.8	4.7	1.0	3.2	0.9	2.2	0.8	31.7	3.8	698
Q5(richest)	49.3	3.7	8.2	1.4	3.6	1.1	1.4	0.7	2.5	0.7	35.1	3.7	725
M	50.7	4.9	12.0	3.8	5.0	1.6	2.6	1.3	2.6	1.4	27.2	4.6	219
Age													
18-29	56.5	2.3	10.4	1.2	6.0	0.9	2.4	0.8	1.9	0.5	22.8	1.9	1,748
30-44	51.7	2.1	11.3	1.1	5.1	0.8	1.8	0.5	2.6	0.7	27.4	2.1	1,292
45-59	51.7	2.6	15.3	1.8	5.3	1.3	3.8	1.0	3.2	0.9	20.7	2.2	696
60-69	50.4	3.8	20.2	3.2	9.3	2.1	3.9	1.2	2.4	1.0	13.8	2.7	293
70-79	42.8	5.1	21.9	4.0	10.4	2.9	9.6	3.6	6.2	3.7	9.0	2.6	160
80+	39.9	5.0	13.0	5.0	14.3	4.2	17.2	4.5	7.6	3.4	8.0	2.9	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	53.1	1.6	12.6	0.7	6.1	0.6	3.0	0.4	2.6	0.4	22.6	1.5	4,239

6.13 Difficulty with 'seeing and recognizing a person you know across the road (20m)' in last 30 days - q2071

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	59.3	1.8	10.9	1.2	6.4	0.7	2.4	0.5	1.0	0.4	19.9	1.6	1,993
Female	55.9	1.4	12.5	0.8	8.8	0.6	5.7	0.7	1.0	0.3	16.1	1.2	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	62.9	2.0	9.2	0.8	6.9	0.7	2.7	0.4	0.9	0.3	17.4	1.8	1,397
Rural	54.9	1.6	13.0	0.9	8.1	0.6	4.8	0.7	1.1	0.3	18.1	1.4	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	47.5	2.8	15.7	1.9	13.1	1.6	5.6	1.2	0.6	0.3	17.5	2.2	820
Q2	61.3	2.9	10.5	1.4	5.1	1.0	3.9	1.0	0.9	0.5	18.3	2.3	842
Q3	58.9	2.5	10.4	1.6	5.3	0.9	5.9	1.5	0.6	0.4	19.0	2.1	937
Q4	56.4	2.8	12.6	1.8	6.6	1.0	2.8	0.8	1.4	0.7	20.2	2.6	698
Q5(richest)	65.7	2.6	8.6	1.1	9.1	1.2	1.5	0.5	1.8	0.9	13.3	2.0	725
M	51.8	4.9	15.1	4.1	6.5	1.8	4.7	1.7	1.6	1.1	20.3	3.6	219
Age													
18-29	64.0	1.8	8.2	1.0	4.5	0.6	2.2	0.5	0.8	0.4	20.3	1.6	1,748
30-44	64.2	1.9	9.7	1.0	5.4	0.9	1.9	0.5	0.4	0.2	18.4	1.6	1,292
45-59	49.7	2.9	20.0	2.1	10.8	1.5	3.5	0.8	1.3	0.6	14.7	1.7	696
60-69	35.4	3.4	20.2	2.7	16.9	2.6	14.7	2.7	0.2	0.2	12.6	3.0	293
70-79	18.4	3.4	15.4	3.1	27.2	4.9	20.8	4.6	7.9	3.6	10.3	3.0	160
80+	22.3	4.3	14.2	4.0	18.0	4.5	22.0	6.4	4.7	2.8	18.8	5.3	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	57.5	1.2	11.7	0.7	7.7	0.5	4.1	0.5	1.0	0.2	17.9	1.1	4,239

6.14 Difficulty with 'seeing an object at arm's length or in reading' in last 30 days - q2072

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	60.9	1.7	10.2	1.1	5.5	0.7	2.3	0.5	1.1	0.4	19.9	1.6	1,993
Female	56.5	1.6	12.5	0.9	7.9	0.6	5.4	0.8	1.6	0.4	16.1	1.2	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	62.7	2.0	9.8	0.9	6.0	0.6	3.1	0.6	0.7	0.3	17.6	1.9	1,397
Rural	56.6	1.6	12.2	0.9	7.2	0.7	4.3	0.7	1.7	0.4	18.1	1.4	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	48.5	3.1	17.2	2.3	9.7	1.2	5.1	1.2	1.7	0.7	17.7	2.2	820
Q2	59.2	3.0	10.2	1.4	6.5	1.2	3.5	1.0	2.3	0.8	18.3	2.3	842
Q3	60.8	2.8	8.5	1.6	6.0	1.1	4.9	1.1	0.8	0.4	19.0	2.1	937
Q4	57.6	2.8	12.1	1.7	5.5	0.9	4.0	1.1	0.7	0.4	20.0	2.6	698
Q5(richest)	67.6	2.5	10.3	1.5	6.1	1.0	2.3	0.8	0.4	0.2	13.3	1.9	725
M	58.0	4.2	9.0	2.9	6.4	1.8	2.4	1.0	3.8	2.8	20.3	3.6	219
Age													
18-29	65.6	1.7	8.1	1.0	4.0	0.6	1.4	0.4	0.6	0.2	20.4	1.6	1,748
30-44	65.3	1.9	9.1	1.1	5.6	0.8	1.4	0.4	0.4	0.2	18.2	1.6	1,292
45-59	48.8	2.7	20.9	2.2	7.8	1.2	6.2	1.2	1.4	0.7	14.9	1.7	696
60-69	37.5	3.1	15.3	2.6	18.3	3.1	11.4	2.6	4.8	2.0	12.7	3.0	293
70-79	19.7	4.1	18.4	3.8	18.5	3.3	24.6	5.6	8.6	3.3	10.2	3.0	160
80+	23.8	5.3	13.4	3.8	15.7	3.9	17.5	4.9	10.8	3.9	18.8	5.3	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	58.6	1.3	11.4	0.7	6.8	0.5	3.9	0.5	1.4	0.3	17.9	1.1	4,239

6.15 Problem with 'sleeping' in last 30 days - q2080

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	59.5	1.9	14.9	1.3	10.1	1.0	4.8	0.7	0.8	0.3	10.0	1.3	1,993
Female	52.0	1.6	19.3	1.1	12.1	0.9	7.5	0.7	1.5	0.4	7.5	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	64.6	1.4	13.2	1.0	8.9	1.2	4.6	0.6	0.7	0.3	7.9	1.0	1,397
Rural	51.0	1.7	19.2	1.2	12.3	1.0	7.0	0.8	1.4	0.4	9.0	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	48.4	2.4	19.9	1.8	17.1	1.8	8.7	1.7	1.1	0.5	4.8	1.1	820
Q2	55.5	2.7	19.8	2.0	10.4	1.6	6.2	1.2	1.6	0.7	6.5	1.4	842
Q3	52.7	2.6	17.5	1.6	11.3	1.5	6.3	1.4	1.6	0.7	10.8	1.8	937
Q4	57.2	2.6	14.8	1.8	9.1	1.4	5.4	1.1	1.1	0.6	12.4	2.3	698
Q5(richest)	66.5	2.3	13.0	1.7	8.3	1.5	3.6	0.8	0.5	0.3	8.1	1.5	725
M	52.5	4.0	17.6	3.7	7.9	2.0	8.3	2.2	0.8	0.7	12.9	3.0	219
Age													
18-29	61.5	1.8	14.2	1.2	8.0	1.0	4.8	0.8	0.7	0.4	10.8	1.3	1,748
30-44	60.2	1.9	15.9	1.3	9.4	1.0	4.9	0.8	0.8	0.3	8.9	1.4	1,292
45-59	49.4	2.9	20.3	2.4	15.9	2.1	7.1	1.4	1.0	0.6	6.2	1.2	696
60-69	34.3	3.7	27.7	3.5	18.1	2.7	12.3	3.0	2.4	1.1	5.1	1.8	293
70-79	27.4	4.7	24.3	5.2	23.8	4.1	16.1	3.8	7.1	3.7	1.3	0.9	160
80+	25.3	5.1	29.5	5.9	22.0	6.2	13.8	3.6	4.4	2.1	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	55.5	1.2	17.2	0.8	11.2	0.8	6.2	0.6	1.2	0.2	8.7	0.8	4,239

6.16 Problem with 'feeling rested and refreshed' in last 30 days - q2081

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	58.7	1.9	13.6	1.1	11.0	1.1	5.7	0.8	0.9	0.4	10.1	1.4	1,993
Female	51.7	1.6	18.7	1.2	14.2	1.1	6.8	0.8	1.0	0.3	7.6	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	65.0	1.4	13.2	0.9	9.5	0.9	3.8	0.5	0.4	0.2	8.1	1.1	1,397
Rural	50.0	1.7	17.8	1.1	14.2	1.1	7.5	1.0	1.2	0.4	9.2	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	45.6	2.4	20.6	1.9	17.7	1.6	9.7	1.2	1.6	0.8	4.8	1.1	820
Q2	56.2	2.6	16.7	1.6	13.0	1.6	6.8	1.2	1.1	0.6	6.3	1.4	842
Q3	52.8	2.6	15.1	1.5	13.8	1.9	5.9	1.1	1.1	0.5	11.3	1.9	937
Q4	58.3	2.6	13.4	1.4	10.4	1.6	4.9	1.1	0.5	0.4	12.5	2.3	698
Q5(richest)	65.8	2.4	13.8	1.7	7.8	1.2	3.7	1.2	0.4	0.3	8.4	1.7	725
M	48.0	4.2	21.1	4.5	11.2	3.8	6.8	1.9	0.0	0.0	12.9	3.0	219
Age													
18-29	62.6	1.9	12.4	1.0	8.8	1.0	4.7	0.8	0.8	0.5	10.8	1.3	1,748
30-44	59.3	2.0	15.0	1.2	11.0	1.2	5.2	1.2	0.6	0.2	8.9	1.4	1,292
45-59	48.4	2.6	20.4	2.2	16.3	2.2	7.0	1.2	0.5	0.4	7.3	1.5	696
60-69	32.2	3.3	26.7	3.5	22.9	3.7	11.3	3.0	1.5	1.0	5.3	1.8	293
70-79	18.4	3.6	27.6	4.3	29.0	4.7	18.7	4.2	5.1	3.2	1.3	0.9	160
80+	17.2	4.5	32.4	7.6	28.1	5.7	13.0	3.4	4.2	2.1	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	55.0	1.2	16.3	0.8	12.7	0.8	6.3	0.7	0.9	0.3	8.8	0.9	4,239

6.17 Problem with 'feeling sad, low, or depressed' in last 30 days - q2090

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	58.4	2.0	14.8	1.1	10.0	1.2	5.8	0.9	1.1	0.3	9.9	1.3	1,993
Female	51.6	1.6	19.8	1.2	11.6	0.9	7.4	0.7	2.0	0.4	7.7	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	62.7	1.5	13.4	0.8	9.5	0.9	5.1	0.7	1.4	0.3	7.9	1.0	1,397
Rural	50.9	1.9	19.4	1.2	11.5	1.1	7.4	0.8	1.7	0.4	9.1	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	42.7	2.9	23.2	2.0	16.8	2.1	10.1	1.4	2.3	0.9	4.8	1.1	820
Q2	57.3	2.7	17.7	1.8	9.3	1.2	7.6	1.2	1.6	0.6	6.5	1.4	842
Q3	51.7	2.8	18.5	1.9	10.7	1.6	6.2	1.3	1.8	0.6	11.0	1.8	937
Q4	59.7	2.5	13.5	1.6	8.4	1.2	4.6	0.9	1.4	0.6	12.4	2.3	698
Q5(richest)	65.2	2.6	12.8	1.4	9.1	1.5	4.0	0.8	0.9	0.4	8.1	1.5	725
M	53.7	5.4	18.1	5.0	8.2	3.0	6.3	2.1	0.9	0.7	12.9	3.0	219
Age													
18-29	58.1	1.9	16.3	1.2	8.6	1.0	5.3	0.9	0.9	0.3	10.9	1.3	1,748
30-44	58.2	2.3	15.9	1.4	10.4	1.4	5.7	1.1	1.0	0.3	8.9	1.4	1,292
45-59	47.5	2.8	20.9	2.4	14.3	1.9	8.4	1.5	2.6	1.0	6.2	1.2	696
60-69	48.0	3.4	22.6	3.3	13.9	2.7	8.6	2.0	1.8	0.9	5.1	1.8	293
70-79	40.3	4.4	17.3	4.4	14.6	3.2	16.0	4.2	9.4	3.8	2.4	1.4	160
80+	38.0	6.3	19.8	5.4	23.1	4.4	11.2	4.6	2.9	1.5	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	54.8	1.4	17.4	0.8	10.8	0.8	6.6	0.6	1.6	0.3	8.7	0.8	4,239

6.18 Problem with 'worry or anxiety' in last 30 days - q2091

	None	SE	Mild	SE	Moderate	SE	Sever	SE	Extreme/ Cannot do	SE	Missing	SE	N
Gender													
Male	59.7	1.8	13.7	1.0	9.4	1.1	5.8	0.8	1.1	0.4	10.3	1.3	1,993
Female	52.3	1.5	18.7	1.1	11.8	1.0	7.5	0.8	1.7	0.3	7.9	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	62.8	1.6	13.6	1.1	9.7	0.9	4.8	0.6	0.8	0.3	8.2	1.0	1,397
Rural	52.3	1.7	17.7	1.1	11.1	1.0	7.7	0.8	1.7	0.4	9.5	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	44.1	2.6	22.1	1.9	15.0	1.9	11.9	2.1	1.5	0.5	5.5	1.2	820
Q2	58.8	2.6	17.2	1.8	8.4	1.2	7.3	1.3	1.6	0.6	6.8	1.4	842
Q3	54.9	2.6	15.9	2.0	10.1	1.6	6.4	1.1	1.6	0.6	11.1	1.8	937
Q4	59.0	2.7	13.1	1.6	10.7	1.4	3.4	0.8	1.3	0.6	12.5	2.3	698
Q5(richest)	65.6	2.6	11.4	1.6	9.0	1.4	4.3	0.9	1.2	0.8	8.6	1.5	725
M	49.6	5.4	20.4	3.5	10.7	4.1	5.7	1.9	0.7	0.7	13.0	3.0	219
Age													
18-29	59.6	1.8	15.0	1.2	7.7	0.8	5.5	0.8	1.1	0.4	11.2	1.3	1,748
30-44	58.8	2.1	15.1	1.4	10.5	1.2	5.5	0.9	0.8	0.3	9.3	1.4	1,292
45-59	49.5	2.6	17.8	2.2	14.9	1.9	8.5	1.6	2.6	1.0	6.7	1.3	696
60-69	48.0	3.4	24.0	2.8	13.0	2.4	8.0	1.7	1.8	1.0	5.2	1.8	293
70-79	36.4	4.8	20.3	4.9	19.4	4.4	17.9	4.5	3.7	2.2	2.4	1.4	160
80+	40.1	6.2	18.1	4.5	19.8	4.5	15.1	4.2	1.8	1.0	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	55.8	1.2	16.4	0.8	10.6	0.7	6.7	0.6	1.4	0.3	9.1	0.9	4,239

7.1 Self-reported utilization of health services and perceived unmet need

	<u>Ambulatory Adult</u>		<u>Ambulatory Child</u>		<u>Inpatient Adult</u>		<u>Inpatient Child</u>		<u>Informal Care</u>		<u>Unmet Care</u>		<u>Non use</u>	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female	22.7	1.3	10.1	0.9	4.7	0.6	2.3	0.4	1.0	0.2	0.8	0.2	58.4	1.5
Male	20.0	1.3	5.4	0.8	2.6	0.4	0.9	0.3	1.4	0.4	0.2	0.1	69.5	1.6
Area of Residence														
Urban	22.2	1.5	9.9	1.2	4.0	0.6	2.0	0.4	0.7	0.3	0.5	0.2	60.7	1.9
Rural	21.2	1.3	7.0	0.7	3.7	0.5	1.6	0.4	1.4	0.3	0.5	0.2	64.7	1.4
Age Group														
18-29y	17.9	1.4	8.0	1.0	3.5	0.6	1.9	0.5	0.6	0.2	0.8	0.3	67.2	1.8
30-44y	21.5	1.5	12.4	1.2	3.7	0.6	2.0	0.4	1.1	0.4	0.3	0.2	58.9	1.9
45-59y	25.1	2.5	4.6	1.0	2.7	1.0	1.6	0.6	1.9	0.8	0.3	0.2	63.8	2.5
60-69y	29.0	3.8	4.6	1.9	5.4	1.9	1.4	0.9	2.6	1.3	0.4	0.4	56.7	4.2
70+y	29.1	3.3	0.7	0.4	6.2	1.7	0.0	0.0	1.8	1.1	0.5	0.4	61.8	3.8
Education Years														
0	20.7	2.0	6.0	1.1	3.7	0.8	1.4	0.5	1.0	0.7	0.2	0.2	66.9	2.4
1-5	26.9	2.5	8.8	1.6	2.9	0.9	1.3	0.5	1.8	0.8	0.2	0.2	58.1	2.5
6-11	19.0	1.4	8.4	1.1	4.4	0.7	2.1	0.6	0.9	0.3	0.8	0.3	64.4	1.7
12+	21.9	2.0	9.1	1.2	3.5	0.7	2.1	0.6	1.0	0.5	0.4	0.2	61.9	2.3
Income Quintiles														
Q1	25.7	2.2	8.2	1.4	4.2	0.9	1.2	0.4	1.0	0.4	0.4	0.2	59.3	2.2
Q2	18.7	2.2	6.5	1.3	4.4	1.0	2.2	0.7	2.2	0.8	0.9	0.4	65.1	2.5
Q3	18.6	2.2	6.6	1.1	2.4	0.6	1.7	0.6	1.2	0.5	0.6	0.3	68.9	2.4
Q4	21.4	2.2	8.1	1.5	4.7	1.0	1.1	0.4	1.0	0.6	0.8	0.4	62.9	2.8
Q5	23.5	2.4	10.8	1.8	3.2	0.8	2.2	0.6	0.6	0.3	0.2	0.1	59.7	2.8
Ownership														
Government	32.8	1.8	12.7	1.4	7.6	0.9	2.3	0.5	1.7	0.5	0.9	0.3	41.9	2.0
Private	46.4	4.5	16.9	3.3	2.9	1.5	2.6	1.8	0.9	0.9	0.0	0.0	30.3	4.9
NGO	31.1	8.8	4.8	3.0	15.3	7.5	2.0	1.7	0.0	0.0	0.5	0.5	46.3	6.9
Other	41.2	12.0	7.6	3.2	7.4	6.6	18.9	5.5	0.0	0.0	0.0	0.0	24.8	8.7
TOTAL	21.6	1.0	8.1	0.6	3.8	0.4	1.7	0.3	1.1	0.2	0.5	0.1	63.1	1.2

This table presents the percent of people reporting utilization of health services when they last needed care. It combines information from Q7000, Q7001 and Q7016. Utilization was recorded in the WHS in three separate categories: ambulatory care, hospital inpatient care, and informal care. Time cut-offs for utilization recall used here are 1 year for ambulatory care and informal care, and 3 years for hospital inpatient care. Utilization of ambulatory and inpatient care services

have been further sub-divided into "adult" and "child" categories. In addition to utilization, two other related aspects are being reported here - unmet need and non-use (regardless of expressed need) of health services. Unmet need was estimated on the basis of responses to Q7000 and Q7004. Non-use of health services is simply the percent of respondents who reported no need in Q7000 or did not utilize services in the time cut-offs mentioned above.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, and ownership (of health facility). Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership was constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services.

7.2 Patient assessed characteristics of ambulatory health services

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
Sex																
Female	32.2	3.4	93.5	2.6	93.3	3.2	91.2	3.1	27.9	2.9	33.7	3.5	494.0	411.7	9.6	1.8
Male	42.2	4.1	95.5	1.6	94.5	1.8	95.4	1.4	31.0	3.9	34.4	4.1	63.7	13.0	7.0	2.0
Area of Residence																
Urban	55.0	3.8	93.6	1.6	94.2	1.4	91.5	1.7	18.7	1.8	37.7	4.2	730.6	602.8	9.5	1.8
Rural	24.7	3.6	94.9	2.4	93.5	3.1	94.1	2.9	36.2	3.6	31.6	3.5	40.8	8.3	7.8	2.0
Age Group																
18-29y	31.7	3.9	92.4	2.6	90.5	3.6	90.4	2.7	29.2	3.6	29.4	4.1	44.6	10.1	6.3	2.0
30-44y	43.0	5.1	93.5	2.8	94.5	2.4	94.4	2.4	25.5	3.5	40.1	4.8	923.1	771.6	13.1	3.5
45-59y	36.7	5.6	99.3	0.8	98.7	1.2	93.7	3.0	25.5	3.8	39.1	6.0	49.0	17.9	9.3	3.5
60-69y	45.5	9.3	95.1	3.6	95.9	3.5	94.8	3.6	34.9	5.2	35.5	8.4	46.4	12.0	3.2	1.6
70+y	29.1	9.0	97.2	2.0	95.7	2.3	96.1	2.2	40.2	9.3	25.3	7.0	43.8	20.3	6.3	3.6
Education Years																
0	25.4	5.7	97.0	1.5	96.6	1.7	96.3	1.7	36.7	4.8	30.8	5.4	33.8	10.0	10.3	3.3
1-5	40.8	6.9	95.4	2.6	93.1	2.8	89.8	3.7	32.5	6.4	18.7	4.9	34.6	11.6	7.2	2.9
6-11	33.2	4.5	91.0	3.8	91.3	3.7	89.8	3.6	31.1	3.9	31.1	4.4	45.2	11.6	9.2	2.7
12+	48.9	5.7	95.4	2.3	94.5	2.6	96.9	1.7	21.9	3.1	50.0	5.5	1,007.1	839.7	7.1	2.5
Income Quintiles																
Q1	22.1	4.8	98.7	0.8	95.5	2.4	94.6	2.4	39.4	5.4	27.8	6.0	42.7	17.2	12.8	4.3
Q2	27.3	4.7	93.8	3.7	95.2	3.0	98.1	1.4	31.5	5.4	18.1	4.2	43.3	14.3	7.7	3.0
Q3	30.8	6.2	90.7	4.9	91.7	4.8	91.0	4.9	37.5	6.3	34.9	6.4	34.0	6.3	1.9	1.3
Q4	40.2	4.9	91.8	3.5	91.5	3.4	90.2	3.6	21.2	2.9	30.9	5.5	72.9	22.2	7.1	3.0
Q5	59.6	6.2	96.2	2.2	94.7	2.9	93.9	2.5	17.3	2.5	59.3	6.2	1,421.2	1,213.8	9.2	2.8
Condition																
Child-Visit	26.2	5.4	98.6	1.4	100.0	0.0	95.3	3.1	22.2	4.4	30.5	7.4	40.4	10.3	8.1	3.6
Preventive	37.5	12.7	99.6	0.4	89.7	9.2	99.6	0.4	27.2	10.5	53.5	11.4	4,749.6	4,154.8	7.2	7.0
Childbirth	41.5	25.0	100.0	0.0	100.0	0.0	100.0	0.0	17.4	4.7	54.7	25.7	34.1	6.0	0.0	0.0
Acute	32.4	5.4	91.5	4.9	90.7	4.9	89.9	5.0	26.2	3.2	34.4	5.7	58.7	19.8	12.0	3.4
Injury	29.8	9.0	90.0	7.2	96.9	3.0	86.3	7.5	27.7	7.2	34.8	10.0	46.7	12.0	0.0	0.0
Chronic	50.8	9.8	90.8	5.6	92.1	5.5	90.8	5.6	24.9	4.7	31.1	8.8	69.9	23.5	8.1	4.3

	Patients visiting a medical doctor, of those visiting any health care provider		Patients rating characteristics of health care provider as adequate/more than adequate						Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Mean cost for an ambulatory care visit (local currency)		Percent of population perceiving discrimination	
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	mean cost	SE	%	SE
Other	38.1	4.9	94.5	2.4	92.0	2.7	94.5	1.9	33.7	5.1	34.9	4.5	143.1	86.2	9.7	2.7
Ownership																
Government	26.4	3.0	93.5	1.9	92.2	2.5	92.0	2.2	31.0	2.9	26.4	2.8	35.2	5.9	7.9	1.5
Private	81.9	4.6	97.3	2.7	100.0	0.0	100.0	0.0	22.1	3.9	70.9	5.1	1,712.6	1,372.1	10.6	3.8
NGO	50.4	17.6	100.0	0.0	100.0	0.0	81.8	16.4	21.6	7.3	24.8	16.5	32.3	8.4	18.2	16.4
Other	86.9	15.4	100.0	0.0	100.0	0.0	100.0	0.0	25.5	5.1	72.1	23.2	30.0	2.6	0.0	0.0
Self-Assessed Health																
Very-Good	30.7	3.9	95.4	2.0	94.5	3.1	93.6	2.8	24.5	2.8	38.1	4.2	56.8	14.0	5.4	1.4
Good	43.5	4.8	97.0	2.2	96.2	2.3	93.1	2.8	33.3	4.4	31.8	4.8	841.7	722.7	9.4	2.7
Moderate	36.9	5.5	90.0	3.3	89.5	3.2	92.7	2.2	32.2	4.8	30.7	5.1	57.7	15.3	12.6	3.6
Bad	24.8	7.7	93.3	4.7	92.1	4.7	92.1	4.7	25.7	4.5	36.6	10.4	49.5	26.4	4.9	3.4
Very-Bad	71.2	15.1	85.6	13.8	100.0	0.0	85.6	13.8	22.7	7.7	23.7	16.4	92.2	38.2	8.1	8.3
TOTAL	36.6	2.9	94.4	1.6	93.8	2.0	93.0	1.8	29.3	2.5	34.0	2.7	308.2	235.5	8.5	1.4

This table looks at various characteristics of a visit to an ambulatory care facility, for those who utilized such services in the previous 12 months (Q7204 and Q7205). Characteristics of a visit reported here include:

- (1) The percent of patients visiting a medical doctor (of those visiting any ambulatory provider) (Q7302).
- (2) Patients' rating of *skills*, *equipment*, and *drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7304, Q7305 and Q7306, respectively.
- (3) Barriers or facilitators to care access have been reported here for those who utilized ambulatory care services in the last 12 months. They include:
 - (a) *Mean time taken to reach facility*: patient reported times from Q7307 were used to calculate averages across sample sub-groups.
 - (b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7308.
 - (c) *Mean cost of visit*: patient reported costs in Q7309-Q7313 were summed up for an individual and then averaged across sample sub-groups.
 - (d) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination listed in Q7328-Q7334.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include: sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents could select one of the

following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.3.1 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex												
Female	25.9	3.0	30.0	3.1	16.6	2.3	19.0	2.8	23.8	3.1	28.1	3.2
Male	28.1	3.6	35.7	4.1	16.6	2.6	17.4	2.8	21.5	3.8	25.8	4.3
Area of Residence												
Urban	19.3	3.0	27.5	3.1	17.2	2.8	15.9	2.7	19.2	3.2	26.5	3.4
Rural	31.9	3.6	35.8	3.7	16.2	2.2	19.9	2.8	25.1	4.0	27.4	4.2
Age Group												
18-29y	27.5	4.1	27.4	4.4	13.4	2.6	19.3	3.6	27.4	4.4	32.3	4.4
30-44y	27.4	4.5	32.3	4.4	19.4	3.6	17.8	3.7	22.1	4.0	27.6	4.2
45-59y	24.5	5.7	38.3	5.8	18.7	4.8	21.5	5.4	17.3	4.4	18.2	4.6
60-69y	23.6	6.4	21.4	4.9	15.6	5.0	15.5	5.3	12.9	5.7	12.9	5.7
70+y	28.9	7.7	52.6	9.2	18.4	5.6	14.2	5.3	23.9	8.3	30.5	8.5
Education Years												
0	39.8	5.9	48.5	6.8	20.4	5.4	20.2	5.2	26.0	6.6	32.4	6.9
1-5	21.8	5.7	35.7	7.0	17.4	4.3	18.0	4.4	26.0	5.2	32.8	5.3
6-11	29.6	4.2	34.5	4.8	22.8	3.9	22.5	4.1	24.6	4.7	28.9	4.8
12+	17.5	4.4	14.2	3.9	5.9	2.1	11.9	3.6	16.7	3.8	19.6	4.1
Income Quintiles												
Q1	43.1	7.2	52.4	5.7	24.0	4.6	30.8	4.9	42.0	6.6	41.1	6.7
Q2	36.2	6.0	38.4	5.1	20.7	4.1	25.4	5.2	15.0	4.4	21.2	5.0
Q3	24.8	6.0	22.9	5.5	14.9	4.8	15.2	4.9	18.3	5.0	25.3	5.5
Q4	18.4	4.3	31.8	5.6	11.0	3.2	11.5	3.2	23.1	5.1	30.0	5.4
Q5	11.0	3.0	16.0	4.4	11.3	4.3	8.1	4.1	11.5	4.2	15.0	4.4
Condition												
Child-Visit	19.5	7.5	26.8	7.9	18.1	6.3	23.1	7.0	22.0	5.8	20.2	5.7
Preventive	31.9	12.4	31.0	10.6	27.1	9.7	28.8	12.6	8.4	5.8	20.9	8.2
Childbirth	0.0	0.0	0.0	0.0	31.6	25.2	31.6	25.2	31.6	25.2	31.6	25.2
Acute	32.1	5.0	39.2	5.3	22.8	4.2	20.5	4.2	32.7	5.4	33.6	5.7
Injury	23.1	8.4	32.6	10.2	20.2	8.4	4.9	2.6	36.5	11.2	53.4	10.9
Chronic	23.7	8.9	26.3	7.8	15.8	7.0	11.6	5.5	11.1	5.3	15.5	5.8
Other	22.6	3.8	22.4	3.6	11.1	2.4	13.5	3.0	13.8	3.0	21.5	3.6
Ownership												
Government	28.6	3.0	36.5	3.0	19.4	2.1	22.1	2.4	26.4	3.1	31.4	3.3
Private	12.8	4.2	11.8	3.8	3.0	1.5	1.6	1.2	4.7	2.0	7.3	2.3
NGO	54.7	18.3	36.5	18.3	26.3	16.9	18.2	16.4	18.2	16.4	18.2	16.4
Other	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Self-Assessed Health												
Very-Good	21.9	3.6	24.2	4.0	16.5	3.2	16.3	3.4	21.6	4.0	24.6	4.1
Good	23.8	4.7	28.0	5.1	7.2	2.0	11.6	2.7	20.1	4.4	25.3	4.8
Moderate	35.8	4.8	49.0	4.9	28.1	4.8	29.4	4.8	31.6	5.8	35.1	5.5
Bad	31.7	9.9	45.2	10.9	23.0	9.6	22.0	9.1	10.6	4.8	16.2	8.4
Very-Bad	62.5	19.5	36.3	19.2	14.4	13.8	22.3	19.3	0.0	0.0	28.9	20.0
TOTAL	26.9	2.5	32.5	2.6	16.6	1.7	18.3	2.0	22.8	2.7	27.1	2.9

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.3.2 Patient assessment of responsiveness domains for ambulatory services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Autonomy				Confidentiality				Choice of Healthcare Provider		Basic Amenities			
	Treatment Information		Involvement		Talk Privately		Confidentiality of Records				Cleanliness		Space	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE		
Sex														
Female	27.1	3.1	27.8	3.3	21.1	3.0	24.4	3.2	27.6	3.2	21.9	2.6	24.0	2.7
Male	23.5	3.6	26.0	4.1	18.6	3.2	19.7	3.1	25.1	3.8	19.9	3.4	23.6	3.7
Area of Residence														
Urban	24.1	3.5	23.7	3.1	21.1	3.0	19.8	3.0	25.4	3.4	17.3	2.5	23.2	3.4
Rural	26.4	3.8	29.1	4.5	19.2	3.3	24.0	3.8	27.1	4.0	23.5	3.4	24.2	3.4
Age Group														
18-29y	25.7	3.6	32.5	4.6	21.4	3.6	25.3	4.0	29.0	4.0	24.3	3.8	26.6	3.7
30-44y	28.0	4.2	26.0	4.2	24.1	3.8	23.4	4.1	24.7	4.0	21.6	3.8	24.8	4.0
45-59y	18.2	4.6	19.4	4.8	15.0	4.1	21.4	5.0	29.0	5.9	16.0	4.6	13.7	4.0
60-69y	17.6	6.3	15.8	6.0	15.8	6.0	12.9	5.7	18.8	5.9	14.0	5.6	18.2	6.6
70+y	34.2	8.6	29.2	7.6	13.2	5.2	17.6	6.5	25.6	7.7	20.3	6.7	29.5	8.2
Education Years														
0	33.5	6.6	30.5	6.4	21.3	5.4	26.7	6.0	29.3	6.8	25.8	5.4	26.8	5.8
1-5	33.4	5.8	32.4	6.0	25.1	4.4	27.7	5.1	27.5	5.2	19.2	4.8	27.0	4.9
6-11	26.7	4.8	32.1	5.1	22.6	4.6	24.3	4.4	29.6	4.3	22.2	3.5	25.2	4.2
12+	14.1	3.8	16.4	4.2	10.5	3.1	16.0	4.1	19.7	4.1	17.4	4.1	16.3	4.0
Income Quintiles														
Q1	39.2	6.0	42.2	6.6	25.1	5.4	33.9	6.1	44.3	5.8	33.9	5.0	32.7	5.0
Q2	19.8	4.9	20.0	5.5	22.1	5.7	20.8	5.2	22.1	5.4	17.4	4.4	22.6	4.0
Q3	24.6	5.3	32.3	6.8	20.0	5.2	22.3	5.2	24.5	5.6	23.8	5.7	22.0	5.7
Q4	28.3	5.8	25.6	4.8	21.9	3.8	19.0	3.7	23.5	4.6	15.0	3.4	23.5	5.2
Q5	13.1	4.4	12.9	4.3	11.5	4.4	14.3	4.8	14.8	4.5	9.6	2.9	14.3	4.5
Condition														
Child-Visit	16.5	4.9	17.4	4.9	20.9	6.1	17.1	5.4	19.7	5.2	11.8	4.1	16.3	4.8
Preventive	18.2	7.8	13.2	7.3	21.7	8.1	29.2	10.8	19.3	8.2	14.1	7.4	11.1	6.3
Childbirth	31.6	25.2	31.6	25.2	45.3	25.7	0.0	0.0	45.3	25.7	31.6	25.2	31.6	25.2
Acute	34.9	5.5	34.3	5.5	29.2	5.1	30.7	5.1	30.9	4.8	27.1	4.8	31.2	5.0
Injury	54.0	11.0	49.0	11.3	23.0	7.9	30.6	10.8	47.8	11.4	21.7	10.2	31.4	11.2
Chronic	17.8	6.9	24.5	8.3	13.6	6.6	6.9	3.8	17.3	6.8	17.9	6.6	12.8	5.2
Other	18.0	3.1	23.0	4.2	16.6	3.2	17.4	3.3	22.0	3.8	14.1	2.7	19.5	3.2
Ownership														
Government	29.7	3.0	31.7	3.4	23.0	2.7	25.9	2.8	30.7	3.1	23.9	2.6	26.9	2.9
Private	4.0	1.9	4.5	1.9	6.7	2.9	4.9	2.4	6.1	2.7	4.8	2.4	9.0	3.7
NGO	30.3	18.4	26.3	16.9	26.3	16.9	22.3	16.2	30.3	18.4	30.3	18.4	18.2	16.4
Other	13.1	15.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	13.1	15.4	27.9	23.2
Self-Assessed Health														
Very-Good	22.4	4.0	22.5	4.1	16.4	3.8	23.3	4.2	27.8	4.1	22.1	3.4	22.4	3.6
Good	19.9	3.8	24.3	4.8	17.1	3.3	15.3	3.2	20.8	3.8	16.3	3.6	17.2	3.7
Moderate	38.2	5.6	38.3	5.7	30.0	5.1	30.4	5.7	31.9	5.6	25.8	5.0	34.9	5.1
Bad	18.0	8.5	21.5	8.9	8.9	3.8	20.5	8.6	33.9	10.1	23.9	8.8	23.0	8.8
Very-Bad	42.7	20.8	28.6	17.4	43.3	20.8	28.9	20.0	8.1	8.3	9.3	9.4	31.6	20.3
TOTAL	25.5	2.7	27.0	3.0	19.9	2.3	22.3	2.6	26.5	2.7	21.0	2.3	23.8	2.4

The following table reports the percent of people perceiving poor responsiveness in ambulatory care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in Q7315-Q7327 which have 5 response options (including "Good" and "Very good"). Ratings for each of the seven domains of ambulatory care responsiveness (including 13 questions) by patients have been presented here. There are two questions each for six of the seven domains; the domain choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7315) and wait time (Q7316).
- (2) *Dignity*: talked respectfully (Q7317) and privacy (Q7318).
- (3) *Communication*: clear explanation (Q7319) and time for questions (Q7320).
- (4) *Autonomy*: treatment information (Q7321) and involvement (Q7322).
- (5) *Confidentiality*: talk privately (Q7323) and confidentiality of records (Q7324).
- (6) *Choice of health care provider*: Q7325.
- (7) *Basic amenities*: cleanliness (Q7326) and space (Q7327).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used and regrouped information from Q7003 ("Which reason best describes why you [your child] last needed health care?") into 6 categories. The six categories include preventive care (immunization, antenatal consultation and family planning); childbirth; acute conditions (high fever, severe diarrhoea, or cough); injury; chronic conditions (arthritis, asthma and heart disease); and other (dental care, minor surgery and other). Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.4 Patient assessed characteristics of hospital inpatient health services

	Patients staying for more than 6 days on last hospital visit		Patients rating characteristics of health care provider as adequate / more than adequate				Mean time taken to reach health facility (in minutes)		Patients using private car or motorcycle to reach facility		Patients waiting more than a month to be admitted		Mean number of patients sleeping in the same room		Mean cost for a hospital visit		Percent of population perceiving discrimination			
	%	SE	%	SE	%	SE	%	SE	Minutes	SE	%	SE	%	SE	No	SE	mean cost	SE	%	SE
Sex																				
Female	33.0	2.6	95.4	1.0	93.1	1.4	92.0	1.4	36.5	6.8	42.3	3.3	1.1	0.6	5.1	0.7	272.2	59.0	18.2	2.2
Male	42.9	4.7	94.9	1.6	91.7	2.2	91.2	2.8	42.0	6.2	38.8	4.5	1.0	0.7	5.0	0.9	213.3	60.6	13.7	2.8
Area of Residence																				
Urban	34.8	2.9	94.1	1.2	92.6	1.5	92.2	1.6	26.7	3.4	43.4	3.7	0.9	0.6	4.5	0.7	329.7	68.4	20.3	2.5
Rural	37.2	3.4	96.0	1.4	92.7	1.6	91.4	1.8	45.3	8.3	39.7	3.6	1.2	0.6	5.5	0.8	205.2	56.4	14.5	2.4
Age Group																				
18-29y	30.6	3.5	95.7	1.5	93.0	2.0	90.7	2.6	31.0	4.4	36.3	3.6	1.0	0.6	4.9	0.8	200.8	75.1	17.2	2.5
30-44y	37.3	3.9	93.7	1.7	90.4	2.2	92.0	2.0	36.9	5.4	47.8	3.8	1.2	0.9	4.9	0.8	282.6	62.4	16.6	2.5
45-59y	42.6	6.8	99.1	0.6	95.5	3.0	93.3	3.3	36.8	7.2	38.3	6.6	1.4	1.4	5.3	1.0	399.9	165.9	20.0	5.7
60-69y	44.3	8.0	92.7	4.5	94.9	4.3	94.3	4.1	53.8	17.7	37.5	8.7	0.6	0.6	6.9	1.5	127.0	70.7	13.2	6.1
70+y	42.4	8.7	94.5	3.3	94.5	3.3	89.7	4.4	77.1	30.0	46.2	9.6	0.0	0.0	5.1	1.0	178.3	104.3	10.0	4.0
Education Years																				
0	38.8	6.4	96.5	1.8	90.9	3.3	92.4	3.0	51.7	10.4	34.3	6.2	0.2	0.2	6.0	1.0	82.8	37.1	18.5	4.6
1-5	39.0	6.2	96.5	2.0	93.1	3.4	91.0	3.8	44.6	5.8	44.2	5.6	1.7	1.4	5.3	1.0	236.3	111.9	18.0	4.7
6-11	37.0	3.9	95.0	1.4	92.8	1.7	90.6	2.6	36.2	8.0	39.1	4.1	1.3	0.9	5.0	0.7	181.9	48.4	19.4	2.5
12+	27.5	4.1	92.6	2.7	92.2	2.8	91.8	2.7	29.4	6.9	53.0	4.5	1.3	0.8	4.2	0.8	560.2	157.2	9.1	2.4
Income Quintiles																				
Q1	36.1	5.9	97.1	1.3	92.2	2.9	89.9	3.1	37.7	7.3	36.4	6.7	0.2	0.2	6.0	0.9	135.3	77.7	18.3	5.1
Q2	32.4	5.7	99.0	0.6	94.5	2.3	97.2	1.5	45.4	6.0	40.7	5.2	1.1	1.1	5.4	0.9	94.6	33.3	16.3	3.8
Q3	47.2	4.6	91.8	2.7	90.0	2.6	86.7	3.8	46.1	14.9	26.0	3.9	0.3	0.3	5.4	0.9	186.5	71.4	14.0	3.1
Q4	32.8	5.1	93.5	3.2	91.3	3.6	91.1	3.7	39.1	8.0	43.0	5.0	2.8	2.1	4.6	0.8	165.0	50.4	17.5	4.6
Q5	24.3	3.4	93.6	2.0	93.9	2.6	92.9	2.8	22.9	3.6	64.8	5.3	0.5	0.5	3.6	0.8	792.4	212.2	15.0	3.7
Condition																				
Child-Visit	44.5	7.6	97.7	2.4	93.2	3.7	95.8	2.9	41.6	9.3	33.9	7.7	1.3	1.4	6.7	1.2	409.4	250.2	20.0	5.9
Fever	24.6	4.3	98.1	1.0	94.5	2.2	91.0	2.8	37.3	4.7	44.9	5.4	0.3	0.3	5.5	0.9	52.7	19.9	19.5	3.9
Childbirth	11.2	3.0	93.9	2.2	94.0	2.2	95.8	1.8	20.2	2.7	49.6	5.0	1.5	1.5	4.4	0.8	387.8	160.6	26.8	4.5
Arthritis	71.8	23.1	100.0	0.0	100.0	0.0	96.2	4.1	56.0	42.7	52.6	23.9	0.0	0.0	2.4	1.0	27.1	2.7	33.9	24.2
Asthma	80.2	12.9	100.0	0.0	85.8	11.6	71.9	8.7	37.6	8.6	16.7	11.1	0.0	0.0	7.8	1.7	44.8	16.4	12.0	9.4
Heart-Disease	24.5	5.6	97.9	1.7	97.9	1.7	97.9	1.7	30.3	6.3	41.1	7.4	1.7	1.7	4.7	0.9	272.0	172.3	12.4	4.3
Injury	56.2	6.4	98.0	1.1	93.3	3.8	92.5	3.9	49.3	15.3	42.7	8.1	0.0	0.0	5.0	1.1	219.7	84.9	5.2	2.3
Minor-Surgery	43.3	6.6	95.4	2.7	97.8	1.5	97.8	1.6	32.2	7.1	53.9	6.4	1.6	1.2	4.3	1.0	730.0	265.8	10.5	3.5
Other	45.3	4.3	90.6	3.2	86.7	3.4	87.5	3.4	45.1	12.2	34.5	4.8	1.2	1.1	5.4	0.8	288.0	83.0	16.2	3.1
Ownership																				
Government	36.6	2.8	95.0	1.0	91.9	1.3	90.4	1.4	39.5	6.0	38.1	2.7	1.3	0.5	5.4	0.7	123.4	24.8	18.5	2.0
Private	31.2	5.3	97.4	2.6	97.4	2.6	97.1	2.6	25.2	4.5	72.8	6.4	0.0	0.0	3.0	0.9	1,212.5	353.8	6.0	2.6
NGO	37.8	10.7	94.0	5.1	93.7	5.7	100.0	0.0	44.1	9.3	21.4	8.9	0.0	0.0	5.5	1.2	400.9	202.0	5.5	5.4
Other	40.1	8.4	100.0	0.0	100.0	0.0	100.0	0.0	29.1	5.0	66.5	9.8	0.0	0.0	3.2	1.0	547.1	454.8	27.5	12.0
Self-Assessed Health																				
Very-Good	25.5	3.3	96.7	1.2	97.6	0.8	96.7	1.1	34.4	5.6	51.5	4.1	0.3	0.3	4.7	0.7	349.8	102.8	12.9	2.4
Good	28.3	3.3	94.5	1.7	89.5	2.7	91.7	2.4	39.8	5.1	40.4	4.1	0.5	0.3	4.7	0.8	217.1	65.8	19.9	2.9
Moderate	44.6	4.2	94.5	2.1	89.9	2.7	88.4	2.8	43.9	10.9	32.5	4.1	1.9	1.3	5.3	0.8	187.4	63.5	18.0	3.3
Bad	60.3	6.9	98.7	0.8	98.1	1.0	87.6	8.0	34.6	9.4	39.4	7.7	2.1	2.1	6.8	1.2	273.5	161.3	13.9	4.4
Very-Bad	70.1	11.2	80.2	11.0	80.2	11.0	81.5	10.6	17.9	9.4	33.8	15.2	5.8	5.9	5.5	1.8	162.5	74.8	22.8	11.9
TOTAL	36.2	2.4	95.3	1.0	92.7	1.2	91.7	1.3	38.2	5.6	41.2	2.7	1.1	0.4	5.1	0.7	252.9	43.7	16.7	1.8

This table looks at various characteristics of a visit to a hospital inpatient care facility, for those who utilized such services in the past 3 years (Q7200, Q7201 and Q7402). Characteristics of a visit reported here include:

(1) The percent of patients who stayed 6 days or more at an inpatient facility during their last visit (Q7404).

(2) Patients' rating of *skills, equipment, and drug supplies* of the provider as "adequate": percent of patients who reported "Yes" to questions Q7405, Q7406 and Q7407, respectively.

(3) Barriers or facilitators to care access have been reported here for those who utilized inpatient care services in the last 3 years. They include:

(a) *Mean time taken to reach facility*: patient reported times from Q7408 were used to calculate averages across sample sub-groups.

(b) *Patients using private motor vehicles*: percent of patients who reported using a "private car or motorcycle" to get to the facility in Q7409.

(c) *Patients waiting more than a month to be admitted*: percent of patients who reported waiting more than a month to be admitted in Q7410.

(d) *Mean number of patients sleeping in same room*: averages across sample sub-groups have been constructed for numbers reported in Q7417.

(e) *Mean cost of visit*: patient reported costs in Q7411-Q7415 were summed up for an individual and then averaged across sample sub-groups.

(f) *Perceived discrimination*: percent of patients who reported "Yes" to any form of discrimination asked in Q7433-Q7439.

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for seeking care (condition), ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.5.1 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 1

	Prompt Attention				Dignity				Communication				Autonomy			
	Travel Time		Wait Time		Talked Respectfully		Privacy		Clear Explanation		Time for Questions		Treatment Information		Involvement	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex																
Female	30.0	2.7	28.6	2.7	29.4	2.7	25.7	2.7	31.2	2.9	32.6	2.7	31.4	2.7	30.3	3.1
Male	31.6	4.1	27.2	3.5	25.6	3.7	22.3	3.4	30.9	3.7	32.4	3.9	31.5	4.0	35.1	4.0
Area of Residence																
Urban	24.7	2.9	29.3	3.4	26.8	3.4	23.1	2.8	29.6	3.1	31.1	3.4	28.9	3.3	28.4	2.9
Rural	34.3	3.2	27.4	2.8	29.1	3.4	25.6	3.1	32.0	3.4	33.4	3.0	33.1	3.0	34.0	3.7
Age Group																
18-29y	27.7	3.0	27.6	3.2	30.5	3.8	23.0	3.1	30.3	3.7	33.9	3.5	30.1	3.5	30.3	3.9
30-44y	28.6	3.6	28.3	3.3	28.7	3.9	23.6	3.3	33.8	3.4	32.3	3.5	33.6	4.0	30.7	3.7
45-59y	35.2	6.6	26.6	5.2	23.5	4.9	33.9	7.3	36.1	7.8	38.4	7.8	36.4	7.8	43.4	7.3
60-69y	37.1	9.0	31.7	9.0	22.0	7.6	23.2	7.5	20.5	7.4	23.2	7.6	21.4	7.5	23.4	7.6
70+y	41.3	8.8	31.1	8.5	28.5	8.4	19.5	7.6	20.3	6.6	20.7	6.6	26.5	6.9	28.6	7.6
Education Years																
0	40.4	7.3	34.4	6.2	34.0	6.1	31.3	6.5	38.9	7.5	35.3	7.5	38.3	6.4	43.8	7.2
1-5	38.3	5.5	29.5	5.1	23.5	4.4	26.6	5.3	35.7	6.2	35.9	5.8	36.0	5.5	33.5	5.1
6-11	26.6	3.7	27.2	3.7	31.6	4.0	26.5	3.9	30.4	4.1	37.4	3.8	32.5	3.6	31.8	3.6
12+	24.2	3.5	22.5	3.7	21.8	4.1	16.8	3.3	25.3	4.0	21.2	3.7	20.4	3.5	20.9	3.4
Income Quintiles																
Q1	42.0	5.2	34.9	4.6	36.3	5.2	32.4	4.7	43.4	6.7	40.2	6.2	42.2	6.1	47.6	6.2
Q2	37.2	5.3	31.7	5.1	25.8	4.8	20.1	4.0	27.9	4.4	26.7	4.2	27.7	4.3	28.0	4.6
Q3	31.1	4.6	26.4	4.5	31.4	6.3	27.9	5.2	29.9	5.7	37.8	6.0	37.8	5.2	29.6	5.9
Q4	26.3	4.4	21.7	4.4	22.8	4.7	21.9	4.7	25.6	4.5	26.6	4.8	26.0	4.8	28.5	5.0
Q5	18.5	4.6	26.0	5.1	25.4	5.6	20.8	4.1	30.0	5.2	30.4	5.2	22.5	3.8	26.3	4.9
Condition																
Child-Visit	26.6	6.4	33.6	7.6	35.1	7.8	35.0	8.5	38.8	8.9	44.0	8.7	41.1	9.0	26.2	6.4
Fever	31.5	5.0	39.0	4.8	42.6	7.0	28.2	5.1	42.5	5.6	46.9	6.2	41.2	5.4	40.3	5.9
Childbirth	19.3	4.3	18.0	3.7	23.1	4.3	21.2	4.7	29.7	4.9	30.1	5.4	27.8	5.3	24.6	4.5
Arthritis	32.1	23.0	36.4	23.1	32.1	23.0	32.1	23.0	3.8	4.1	0.0	0.0	28.2	23.1	62.1	21.2
Asthma	43.3	11.2	15.2	11.2	34.8	7.6	56.0	8.7	62.3	7.1	55.5	9.0	55.6	8.9	40.0	7.4
Heart-Disease	39.7	8.2	25.9	6.4	23.2	5.5	13.8	4.7	20.7	7.7	20.7	7.7	25.1	7.8	21.5	5.3
Injury	47.8	7.7	27.8	5.2	26.5	5.0	23.2	5.0	36.3	7.7	42.5	6.2	43.6	8.8	39.1	8.8
Minor-Surgery	21.0	5.7	32.8	6.2	24.6	5.8	16.4	4.6	26.3	5.9	26.4	6.0	25.9	6.0	28.1	6.0
Other	26.2	4.4	24.8	3.7	19.3	3.6	25.6	4.6	24.0	4.5	26.5	4.4	25.2	4.5	29.7	4.9
Ownership																
Government	33.4	2.6	30.3	2.5	30.3	2.8	27.4	2.5	34.2	2.7	36.0	2.5	35.2	2.6	35.7	2.9
Private	18.8	5.4	21.2	6.0	20.6	6.1	8.1	3.8	18.2	5.3	16.1	4.9	13.7	3.7	16.5	4.9
NGO	17.3	6.9	14.8	8.3	15.9	8.2	17.2	7.9	16.3	8.2	17.0	8.5	15.7	6.6	9.7	5.6
Other	9.2	3.6	9.2	3.6	10.6	3.3	10.6	3.3	9.2	3.6	13.7	3.3	13.7	3.3	13.7	3.3
Self-Assessed Health																
Very-Good	22.7	3.9	21.7	3.1	25.3	3.8	15.6	3.2	23.4	3.6	25.3	3.4	24.8	3.6	26.8	4.4
Good	31.9	3.1	35.7	4.1	30.3	4.5	26.1	3.6	34.7	4.4	36.4	4.3	34.5	4.1	32.4	4.0
Moderate	38.0	4.8	29.6	4.0	29.7	4.4	26.2	4.2	30.5	3.4	30.5	3.4	27.4	3.4	33.8	4.4
Bad	29.6	6.6	20.5	5.6	29.7	8.5	39.9	8.9	39.9	7.5	41.0	7.5	46.0	8.6	37.2	7.3
Very-Bad	29.4	10.7	17.9	9.1	19.9	9.8	44.2	17.4	57.6	18.2	57.6	18.2	55.6	18.2	34.7	13.1
TOTAL	30.5	2.3	28.2	2.1	28.2	2.4	24.6	2.2	31.1	2.4	32.5	2.3	31.5	2.2	31.8	2.5

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.5.2 Patient assessment of responsiveness domains for hospital inpatient services: percent reporting "Moderate", "Bad" or "Very bad" - 2

	Confidentiality				Choice of health care provider		Basic Amenities				Social Support			
	Talk Privately	Confidentiality of records					Cleanliness	Space			Family Visit	External Contact		
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex														
Female	30.5	2.6	29.8	2.4	36.3	2.8	28.4	2.7	26.7	2.3	24.5	2.6	30.5	2.8
Male	26.7	3.2	29.6	3.5	33.6	4.4	24.3	3.5	20.2	3.0	19.8	2.9	27.7	3.6
Area of Residence														
Urban	25.4	3.0	25.4	3.0	31.2	3.5	27.9	2.9	23.5	2.5	20.7	2.6	25.9	3.3
Rural	31.6	2.6	32.4	2.9	38.0	3.2	26.5	3.0	25.2	2.1	24.4	3.0	31.9	3.5
Age Group														
18-29y	31.2	4.0	28.6	3.6	32.2	3.5	27.9	3.4	27.3	3.0	25.1	3.4	33.4	4.0
30-44y	30.9	3.7	28.7	3.6	38.9	4.0	31.7	3.5	22.7	2.7	22.7	2.8	29.4	3.6
45-59y	30.0	7.4	34.5	6.4	39.5	7.9	19.9	4.5	23.8	6.1	15.1	4.2	26.5	6.3
60-69y	16.5	7.1	29.8	8.8	33.1	8.1	22.0	7.2	23.8	7.3	22.6	7.5	22.2	7.5
70+y	18.6	5.3	30.7	8.0	29.1	7.7	18.2	5.0	20.3	5.6	30.7	8.0	21.6	5.8
Education Years														
0	33.5	5.5	41.2	5.1	43.8	7.6	28.9	4.9	26.1	5.1	24.4	5.3	26.8	6.0
1-5	32.1	5.3	29.2	5.1	33.5	5.8	21.2	4.6	22.8	5.1	25.4	5.1	37.9	6.2
6-11	32.7	3.7	30.7	3.5	37.2	3.6	31.1	4.1	27.8	3.2	23.7	3.3	32.9	4.8
12+	21.9	3.4	20.5	3.7	26.8	3.9	24.6	3.8	20.8	3.4	19.2	3.5	21.6	3.9
Income Quintiles														
Q1	37.2	5.1	33.1	4.1	47.5	6.3	29.3	3.9	26.5	3.9	29.9	5.3	35.8	5.5
Q2	27.3	4.3	29.6	4.4	37.9	4.4	30.5	5.5	26.2	4.4	20.8	4.6	27.3	5.1
Q3	35.2	4.7	35.9	5.5	34.9	4.0	25.8	4.4	26.1	3.8	23.5	3.5	35.2	4.9
Q4	25.9	4.8	29.6	5.2	30.1	4.9	28.3	4.7	26.0	4.0	26.9	4.7	29.3	5.4
Q5	18.5	3.3	16.4	3.6	24.6	5.1	24.2	3.9	16.6	3.6	13.0	3.1	17.2	3.6
Condition														
Child-Visit	42.2	8.7	39.1	8.4	45.1	9.2	32.4	8.2	35.2	8.1	19.2	7.5	47.4	8.4
Fever	32.5	4.7	33.3	4.2	49.3	5.8	22.6	3.9	18.8	3.5	19.3	4.7	27.6	5.0
Childbirth	24.9	4.6	23.4	5.2	34.8	5.6	31.7	5.6	28.7	4.8	23.7	4.7	28.6	5.2
Arthritis	28.2	23.1	32.1	23.0	6.3	5.2	0.0	0.0	31.7	23.0	31.7	23.0	31.7	23.0
Asthma	59.9	7.5	51.5	10.2	36.3	9.7	38.9	11.0	32.9	8.7	17.7	12.5	47.9	11.5
Heart-Disease	17.1	5.0	19.6	5.2	21.4	5.3	15.0	4.7	17.3	4.9	20.0	5.4	22.0	5.7
Injury	27.1	5.3	31.2	5.7	45.3	8.4	33.5	7.8	29.3	5.6	19.6	4.7	28.1	5.4
Minor-Surgery	26.3	6.1	24.2	5.8	30.6	6.6	27.8	6.4	20.6	5.3	23.9	5.8	28.8	6.2
Other	28.8	4.9	27.6	4.3	30.2	4.7	29.2	4.1	28.9	4.5	24.3	3.7	31.3	4.8
Ownership														
Government	31.9	2.2	33.5	2.4	39.2	2.7	29.4	2.5	26.9	1.8	25.2	2.4	32.0	2.8
Private	18.7	5.5	10.7	3.8	18.3	5.2	11.4	4.0	9.9	3.8	7.9	3.6	11.0	3.9
NGO	13.0	5.8	11.7	5.8	21.0	7.4	20.5	7.8	19.4	7.8	20.1	7.9	32.2	9.1
Other	16.4	3.4	21.8	6.6	13.5	3.7	30.5	7.4	18.4	5.8	18.4	5.8	16.9	6.1
Self-Assessed Health														
Very-Good	21.9	3.3	20.8	3.2	25.4	4.3	19.0	3.2	18.7	2.8	20.1	3.6	24.1	3.7
Good	33.6	3.6	31.2	3.8	40.3	4.2	29.4	3.7	28.3	3.3	22.5	3.2	32.5	3.9
Moderate	26.5	3.7	29.4	4.4	38.7	4.2	30.3	3.7	26.8	3.5	26.8	4.4	29.8	4.0
Bad	40.1	7.7	46.4	8.7	34.8	8.5	29.7	9.0	22.1	5.6	24.8	5.4	36.4	9.5
Very-Bad	42.2	17.5	62.7	18.1	56.3	18.5	53.9	18.3	39.4	17.6	18.3	9.5	36.8	17.7
TOTAL	29.2	2.0	29.7	2.1	35.4	2.4	27.1	2.2	24.6	1.6	23.0	2.1	29.6	2.5

The following table reports the percent of people perceiving poor responsiveness in hospital inpatient care facilities, where "poor" is defined as a rating of "Moderate", "Bad" or "Very bad" in questions Q7418-Q7432 which have 5 response options (including "good" and "very good".. Ratings for each of the eight domains of hospital inpatient care responsiveness (including 15 questions) by patients have been presented here. There are two questions each for seven of the eight domains; the domain of choice has only one question. The following questions from the survey have been used to determine patient assessment of each responsiveness domain:

- (1) *Prompt attention*: travel time (Q7418) and wait time (Q7419).
- (2) *Dignity*: talked respectfully (Q7420) and privacy (Q7421).
- (3) *Communication*: clear explanation (Q7422) and time for questions (Q7423).
- (4) *Autonomy*: treatment information (Q7424) and involvement (Q7425).
- (5) *Confidentiality*: talk privately (Q7426) and confidentiality of records (Q7427).
- (6) *Choice of health care provider*: Q7428.
- (7) *Basic amenities*: cleanliness (Q7429) and space (Q7430).
- (8) *Social support*: family visit (Q7431) and external contact (Q7432).

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, reason for visit, ownership (of health facility), and self-assessed health. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, condition and self-assessed health sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. The sub-group "Condition" used information from Q7403 ("Which of the following best describes the reason for your [child's] last overnight stay?"). The eight categories of Q7403 include high fever, diarrhoea, or cough; childbirth; arthritis; asthma; heart disease; bodily injury; minor surgery; and other. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad".

7.6 Importance of different domains of responsiveness: percent reporting "Extremely Important"

	Prompt Attention		Dignity		Communication		Autonomy		Confidentiality		Choice		Basic Amenities		Social Support	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
Sex																
Female	43.0	1.4	45.3	1.5	44.7	1.5	38.1	1.3	44.2	1.5	38.0	1.4	46.6	1.4	42.9	1.4
Male	43.8	1.8	47.7	1.8	47.8	1.8	38.4	1.6	48.9	1.8	37.2	1.9	48.0	1.7	46.4	1.8
Area of Residence																
Urban	45.2	2.0	50.0	2.1	48.5	2.2	41.7	1.8	50.0	2.0	41.0	2.0	47.8	2.0	45.1	2.2
Rural	42.2	1.5	44.0	1.4	44.5	1.5	36.1	1.3	43.7	1.7	35.6	1.6	46.8	1.4	43.9	1.6
Age Group																
18-29y	43.7	1.7	46.3	1.9	46.9	1.8	38.8	1.6	45.0	1.8	39.1	1.8	47.5	1.8	44.7	1.9
30-44y	42.0	1.9	46.5	1.9	45.7	1.9	38.1	1.8	48.2	1.9	37.4	1.8	46.8	2.0	43.4	2.0
45-59y	47.3	2.6	49.0	2.7	49.8	2.9	40.8	2.8	48.0	2.9	35.4	2.8	49.1	2.5	42.8	2.9
60-69y	41.0	4.5	47.0	4.0	43.3	4.0	34.2	3.8	50.5	4.3	36.9	4.3	43.6	4.1	48.8	4.7
70+y	41.6	4.1	39.9	3.5	38.0	4.0	35.0	3.6	35.7	3.7	36.1	3.2	46.2	4.4	45.3	3.8
Education Years																
0	42.9	2.5	46.4	2.5	46.1	2.8	36.4	2.2	48.4	2.7	36.2	2.3	48.1	2.6	43.3	2.9
1-5	40.8	2.6	46.0	2.8	40.4	2.8	30.4	2.5	41.5	2.7	31.2	2.4	42.1	2.8	42.2	2.7
6-11	44.0	1.9	45.1	2.0	46.2	1.9	36.7	1.7	45.8	1.8	36.6	1.8	47.3	2.0	44.0	2.0
12+	43.3	2.3	46.4	2.1	47.0	2.5	43.4	2.2	46.2	2.3	41.6	2.2	46.4	2.2	45.2	2.3
Income Quintiles																
Q1	46.1	2.7	50.5	2.6	50.1	2.6	38.8	2.4	48.9	2.6	38.4	2.8	54.0	2.4	48.5	2.8
Q2	45.8	2.6	43.9	2.6	44.9	2.8	37.2	2.3	46.7	2.6	37.0	2.4	46.0	2.4	45.1	2.6
Q3	38.4	2.6	43.0	2.8	39.7	3.0	32.4	2.7	40.6	2.6	35.5	2.4	42.3	3.0	40.3	3.0
Q4	41.4	2.9	41.8	2.8	42.0	3.1	37.5	2.6	42.0	2.8	36.8	2.8	42.1	2.8	38.2	2.8
Q5	42.1	2.9	50.4	3.0	51.4	2.9	43.9	2.7	50.8	2.9	39.0	2.7	48.9	2.8	46.0	3.0
Ownership																
Government	49.7	2.0	53.8	2.2	50.7	2.0	40.0	1.8	51.7	2.1	39.5	1.9	50.9	2.2	49.1	2.1
Private	49.0	5.0	55.7	4.9	54.5	4.6	44.4	4.5	50.8	4.9	41.5	4.6	49.2	5.0	49.1	5.0
NGO	41.3	11.6	37.8	11.6	35.0	10.4	29.0	8.2	37.3	11.8	33.7	8.9	38.2	10.6	35.0	10.4
Other	44.5	12.0	59.9	9.5	47.0	12.4	53.7	7.8	59.6	8.8	47.4	11.0	63.5	7.1	42.0	11.9
Self-Assessed Health																
Very-Good	44.2	1.9	46.6	1.7	48.5	1.8	41.9	1.6	48.9	1.7	41.6	1.6	47.0	1.7	46.5	1.8
Good	42.2	1.8	46.4	2.1	43.7	2.2	35.4	1.9	40.2	2.1	34.8	1.9	45.6	2.0	41.2	2.1
Moderate	40.5	2.5	43.8	2.5	42.9	2.5	36.0	2.6	46.0	2.4	35.2	2.6	47.8	2.6	43.8	2.5
Bad	52.1	4.8	52.0	5.5	48.6	5.0	36.8	4.8	59.6	5.2	35.7	4.7	52.1	4.9	49.9	4.9
Very-Bad	55.6	10.2	53.4	9.4	64.7	9.1	37.0	8.9	52.1	9.7	35.7	9.4	60.0	9.5	45.8	9.6
Type of Care																
Ambulatory-Adult	50.5	2.5	54.3	2.5	54.8	2.6	44.0	2.2	51.0	2.3	40.0	2.7	55.2	2.2	52.5	2.7
Ambulatory-Child	44.6	3.8	46.9	3.9	49.1	4.0	38.8	3.6	48.6	3.8	38.3	3.7	48.1	3.8	46.9	4.2
Inpatient-Adult	50.0	5.3	60.0	5.7	51.6	5.4	36.5	5.0	51.1	5.4	39.7	5.2	51.0	5.6	42.7	4.9
Inpatient-Child	42.0	6.3	45.7	6.4	33.3	6.7	26.8	6.5	41.7	6.6	39.9	6.4	37.9	6.5	37.1	7.1
Non-Users	41.2	1.4	43.3	1.3	43.1	1.5	36.6	1.4	44.4	1.6	37.1	1.4	44.7	1.4	41.9	1.5
TOTAL	43.3	1.2	46.3	1.2	46.0	1.2	38.3	1.1	46.1	1.3	37.7	1.2	47.2	1.2	44.4	1.3

This table shows the percent of people who rated a responsiveness domain as "Extremely Important". All survey respondents were queried on their perception of the importance of each of the eight responsiveness domains in section 7100 of the WHS where each domain was described in detail (Q7100-Q7107).

Respondents were asked to say whether the domain was "Extremely Important", "Very Important", "Moderately Important", "Slightly Important" or "Not

Important".

Results are reported for the total WHS sample (category: "TOTAL"), and by the following sample sub-groups: sex, area of residence, age group, years of education, income quintiles, ownership (of health facility), self-assessed health, and type of care. Mean values and standard errors (SE) are reported for each category.

Sub-groups common to other tables were constructed in the same manner. Common sub-groups include sex (Q1001), area of residence (Q0104), age group (regrouping of reported age in Q1002), years of education (Q1010) and income quintiles. To determine income and subsequently income quintiles, an asset index was constructed for each household using responses to section 0700 on "Permanent Income Indicators" from the WHS household questionnaire.

Ownership, self-assessed health and type of care sub-groups were constructed uniquely for the responsiveness section's tables. There are four categories of ownership: "Operated by the government" (Government), "Privately operated" (Private), "NGO" and "Other". To construct the ownership sub-group, information was combined from (a) Q7204, Q7205 and Q7301 for those utilizing ambulatory services, and (b) Q7200, Q7201 and Q7401 for those utilizing hospital inpatient services. Self-assessed health was constructed using information from Q2000 ("In general, how would you rate your health today as?") in the WHS. Respondents were asked to select one of the following five categories: "Very good", "Good", "Moderate", "Bad" and "Very bad". Type of care is a constructed sample sub-group combining information from Q7000, Q7001 and Q7016. There are five categories of type of care: "Ambulatory adult", "Ambulatory child", "Inpatient adult", "Inpatient child" and "Non users".

8.0 Expenditure Background Indicators (2002)

GDP per capita (\$)	1,481
Total health expenditure per capita (\$)	99
Total population (000)	1,961
Life expectancy at birth (Both sexes)	49.3
Life expectancy at birth (male)	48.1
Life expectancy at birth (female)	50.5
Healthy life expectancy at birth (Both sexes)	43.3
Healthy life expectancy at birth (Male)	42.9
Healthy life expectancy at birth (Female)	43.8
Government funded health expenditure as a percentage of total health expenditure	70.1
Social security funded health expenditure as a percentage of total health expenditure	1.1
Private health insurance premiums as a percentage of total health expenditure	22.4
Out-of-pocket payment as a percentage of total health expenditure	6.1

8.1 Distribution of health payments across different population groups

	Mean Household expenditure (EXP) in local currency	% Poor	% Impoverished	% Catastrophic	OOP % EXP	OOP % non-subsistence spending	Mean OOP in local currency
Catastrophic							
No	978	52.2	0.3		0.8	1.5	16
Yes	1,566	65.6	9.8		29.3	77.1	1002
Poor							
No	1,868		1.6	3.6	2.8	4.1	132
Yes	240		0.0	6.1	1.8	6.1	4
Insurance							
No	723	61.8	0.7	5.1	2.0	5.2	56
Yes	2,318	11.8	0.8	4.2	3.4	5.3	103
Residence							
rural	688	67.9	0.9	6.0	2.3	6.1	69
urban	1,649	22.6	0.5	2.8	2.1	3.4	54
Income_Quintile							
1st(poorest)	102	100.0	0.0	6.5	2.4	6.5	2
2nd	264	100.0	0.0	6.3	1.5	6.3	4
3rd	467	64.3	2.4	4.7	1.4	4.8	8
4th	842	0.0	1.0	2.7	1.8	3.4	20
5th(richest)	3,362	0.0	0.4	4.4	4.2	5.0	289
total	1,007	52.9	0.8	4.9	2.2	5.2	64

OOP: Out of pocket

Household consumption expenditure (EXP) comprises both monetary and in-kind payment on all goods and services and the money value of the consumption of home-made products.

Poor - a household is considered poor when its total household expenditure is smaller than its subsistence spending
Impoverishment - a non-poor household is impoverished by health payment when it becomes poor after paying for health services.

Catastrophic health expenditure occurs when a household's total out-of-pocket health payments equals or exceeds 40% of the households capacity to pay or non-subsistence spending.

Out of pocket health payments (OOP) refer to payments made by households at the point they receive health services. It includes doctor's consultation fees, purchases of medication and hospital bills. Spending on alternative and/or traditional medicine is included. Expenditure on health related transportation and special nutrition is excluded. Insurance reimbursements are also excluded.

Subsistence spending - household subsistence spending is the minimum requirement to maintain a basic life in a society. A poverty line is used in the analysis as subsistence spending.

Food expenditure is the amount spent on all foodstuffs by the household plus the value of the family's own food production consumed within the household. It excludes expenditure on alcoholic beverages, tobacco and food consumption outside the home (e.g hotel and restaurants).

Household capacity to pay - is defined as a household's non-subsistence spending.

8.2 Percent distribution of structure of out-of-pocket health payments

	Inpatient	Outpatient	Traditional	Drugs	Other	Total
Catastrophic						
No	18.5	35.2	3.1	25.7	17.4	100
Yes	44.7	36.7	3.7	6.7	8.2	100
Poor						
No	38.4	36.7	3.4	11.0	10.4	100
Yes	44.7	23.6	10.1	14.7	6.9	100
Insurance						
No	46.1	38.4	2.8	6.9	5.9	100
Yes	20.0	31.2	5.7	21.6	21.5	100
Residence						
rural	41.0	39.3	3.3	8.2	8.2	100
urban	32.4	28.6	4.4	18.5	16.0	100
Income_Quintile						
1st(poorest)	50.5	24.9	0.3	14.1	10.3	100
2nd	43.8	22.3	11.0	19.5	3.4	100
3rd	40.2	29.9	8.0	10.9	10.9	100
4th	14.0	20.3	33.6	21.8	10.2	100
5th(richest)	40.1	37.9	1.4	10.2	10.4	100
total	38.6	36.3	3.6	11.1	10.3	100

8.3 Financial sources used by households for paying for health services

	Savings	Sold Items	Borrow from relatives &	Borrow from others	Health insurance	Current Income	Other
OOP%non_subsistence_spending							
less than 10%	21.6	13.1	16.4	12.2	4.6	44.7	5.7
10-20%	32.9	18.7	23.5	8.2	10.4	67.7	8.1
20-40%	32.2	19.2	21.1	16.5	8.0	60.1	15.4
above 40%	30.8	30.3	24.3	19.8	5.3	56.0	9.9
Hospitalization							
No	22.2	13.2	16.3	12.3	4.3	44.4	5.9
Yes	25.0	19.7	21.4	13.8	8.0	56.9	8.0
Insurance							
No	19.2	16.1	18.7	14.1	2.7	43.7	6.6
Yes	38.8	5.7	9.8	5.2	14.9	58.8	4.7
Residence							
rural	20.5	17.4	19.0	14.0	3.6	41.8	6.7
urban	27.0	8.0	13.2	9.6	7.5	55.4	5.3
Income_Quintile							
1st(poorest)	11.2	18.4	21.0	18.6	2.0	37.2	8.9
2nd	17.9	18.6	23.9	18.5	2.8	41.5	6.6
3rd	21.0	15.2	17.4	12.2	3.8	43.2	4.6
4th	29.8	12.1	13.0	8.9	5.5	52.3	5.8
5th(richest)	33.5	6.9	10.1	4.4	10.3	57.6	5.4
total	22.7	14.2	17.1	12.5	4.9	46.4	6.2

8.4 Insurance Coverage

	Insured	SE	Mandatory Insurance	SE	Voluntary Insurance	SE	Insured as dependent	SE	Mandatory and Voluntary	SE
Q1(poorest)	1.3	0.6	1.3	0.7	0.5	0.2	1.7	0.8	0.6	0.4
Q2	3.7	1.1	2.4	0.9	4.4	1.6	4.2	1.5	2.9	1.6
Q3	7.5	1.6	3.8	0.9	8.9	2.5	7.2	1.9	3.3	1.1
Q4	21.1	2.6	20.6	3.0	20.5	3.1	18.1	2.7	19.1	3.9
Q5(richest)	66.3	3.2	71.9	3.5	65.7	3.6	68.8	3.4	74.0	4.3
Total	100		100		100		100		100	
Number	1,960.0		1,298.0		1,104.0		1,138.0		586.0	

10.1 Inability to control the important things in life - q8000

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
Gender													
Male	35.6	1.6	12.0	1.1	23.5	1.5	7.4	0.8	10.0	0.8	11.5	1.4	1,993
Female	31.2	1.5	12.2	0.9	27.2	1.2	8.4	0.9	11.8	1.0	9.2	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	38.4	1.4	13.7	1.3	27.1	1.2	6.1	0.9	6.3	0.8	8.5	1.0	1,397
Rural	30.8	1.7	11.3	0.9	24.7	1.3	8.8	0.9	13.2	0.9	11.2	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	28.0	2.3	14.2	1.6	28.8	1.8	11.5	1.6	10.4	1.6	7.1	1.3	820
Q2	37.4	3.3	9.4	1.5	25.4	2.1	8.3	1.3	11.5	1.6	8.1	1.5	842
Q3	32.6	2.6	10.2	1.8	20.9	2.2	7.1	1.2	16.8	1.7	12.4	1.8	937
Q4	31.4	2.5	12.6	1.7	26.8	2.3	6.1	1.2	10.4	2.2	12.8	2.3	698
Q5(richest)	39.1	2.3	13.8	1.5	26.0	1.7	7.2	1.8	4.6	1.0	9.3	1.6	725
M	27.2	4.4	15.3	5.1	26.8	3.8	5.3	2.0	8.6	2.0	16.8	3.4	219
Age													
18-29	36.4	1.8	8.8	1.0	23.5	1.6	5.7	0.7	13.4	1.1	12.2	1.4	1,748
30-44	34.0	1.9	10.7	1.1	28.1	1.7	7.0	1.1	9.5	1.1	10.6	1.5	1,292
45-59	29.0	2.4	21.4	1.9	22.7	2.1	10.8	1.8	7.5	1.4	8.6	1.5	696
60-69	28.8	3.8	14.6	2.6	24.3	2.8	14.4	2.7	11.6	2.7	6.3	2.0	293
70-79	26.3	4.6	14.4	3.3	36.5	4.9	11.3	3.3	9.2	3.4	2.4	1.4	160
80+	14.9	4.2	11.0	4.1	37.6	5.2	20.0	5.9	9.8	3.8	6.8	3.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	33.3	1.2	12.1	0.7	25.5	1.0	7.9	0.7	10.9	0.7	10.3	0.9	4,239

10.2 Inability to cope - q8001

	Never	SE	Almost never	SE	Sometimes	SE	Fairly often	SE	Very often	SE	Missing	SE	N
Gender													
Male	33.3	1.7	10.5	0.9	28.6	1.5	8.8	1.0	7.3	0.8	11.5	1.4	1,993
Female	25.9	1.4	14.8	1.1	31.9	1.4	8.9	0.7	9.1	0.8	9.4	1.0	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	34.0	1.6	14.5	1.2	30.1	1.3	7.5	0.8	5.4	0.7	8.5	1.0	1,397
Rural	27.1	1.6	11.9	0.9	30.5	1.5	9.5	0.8	9.6	0.8	11.4	1.3	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	25.2	2.1	13.9	1.5	33.5	2.5	11.6	1.4	8.6	1.4	7.1	1.3	820
Q2	31.5	2.8	10.3	1.5	30.1	2.2	10.5	1.5	9.1	1.5	8.5	1.6	842
Q3	30.2	2.4	10.6	1.5	28.8	2.6	7.9	1.5	9.8	1.4	12.6	1.8	937
Q4	28.7	2.3	14.2	2.0	28.2	2.5	7.8	1.5	8.3	1.5	12.8	2.3	698
Q5(richest)	33.2	2.1	14.5	1.5	31.3	1.9	6.5	1.2	5.1	1.4	9.4	1.6	725
M	22.5	3.9	17.4	3.8	29.8	4.2	6.7	2.2	6.7	2.1	16.8	3.4	219
Age													
18-29	32.6	1.6	10.3	1.0	28.6	1.7	8.0	0.9	8.4	0.9	12.2	1.4	1,748
30-44	29.7	1.9	13.0	1.4	30.4	1.8	9.2	1.2	7.0	0.8	10.7	1.5	1,292
45-59	27.3	2.8	16.1	1.7	32.2	2.9	7.2	1.1	8.0	1.6	9.1	1.6	696
60-69	24.3	3.3	14.4	3.2	32.2	3.6	14.3	2.8	8.5	1.9	6.3	2.0	293
70-79	14.3	3.7	21.0	4.4	34.1	4.2	11.7	4.2	15.3	3.7	3.6	1.8	160
80+	15.7	4.4	9.6	4.0	42.8	7.8	11.1	5.1	14.0	5.4	6.8	3.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	29.4	1.2	12.8	0.7	30.4	1.1	8.8	0.6	8.2	0.6	10.4	0.9	4,239

10.3 Satisfaction with your health - q8002

	Very dissatisfied	SE	Dissatisfied	SE	Neither satisfied nor dissatisfied	SE	Satisfied	SE	Very Satisfied	SE	Missing	SE	N
Gender													
Male	7.5	1.0	5.9	0.8	10.3	1.1	34.3	1.8	30.6	1.5	11.5	1.4	1,993
Female	5.5	0.7	10.6	0.9	14.5	0.9	35.4	1.3	24.5	1.0	9.4	1.0	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	9.0	1.2	7.6	0.9	11.8	0.9	33.1	1.5	30.0	1.3	8.5	1.0	1,397
Rural	5.2	0.9	8.8	0.9	12.9	1.1	35.7	1.4	26.1	1.2	11.3	1.3	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	4.6	1.1	11.1	1.5	18.4	2.1	39.0	2.1	19.9	2.1	7.1	1.3	820
Q2	6.9	1.7	8.2	1.4	14.6	1.8	35.4	2.3	26.5	2.2	8.4	1.6	842
Q3	4.7	1.1	7.7	1.5	10.0	1.4	32.9	2.8	32.0	2.4	12.6	1.8	937
Q4	5.1	1.1	7.3	1.4	9.5	1.5	33.6	2.2	31.9	2.3	12.8	2.3	698
Q5(richest)	10.7	2.0	5.8	1.0	10.3	1.4	34.3	2.3	29.6	2.3	9.3	1.6	725
M	8.9	2.3	13.9	3.1	11.4	3.1	31.6	4.1	17.0	3.2	17.1	3.3	219
Age													
18-29	6.9	0.9	6.6	0.8	8.3	1.1	33.7	1.6	32.3	1.4	12.2	1.4	1,748
30-44	7.1	1.2	5.9	0.7	13.1	1.3	35.3	1.8	27.8	1.6	10.7	1.5	1,292
45-59	4.9	1.0	14.2	1.9	12.6	1.8	37.3	2.4	21.9	2.2	9.0	1.6	696
60-69	5.7	2.1	11.1	2.1	24.3	3.6	35.8	4.0	16.5	3.1	6.6	2.0	293
70-79	4.6	2.1	16.4	3.4	29.2	4.3	30.6	5.1	16.8	4.2	2.4	1.4	160
80+	3.9	2.2	12.0	4.0	24.9	6.6	35.3	6.1	17.1	5.6	6.8	3.0	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	6.4	0.7	8.4	0.6	12.6	0.8	34.8	1.1	27.4	0.9	10.4	0.9	4,239

10.4 Feeling safe when alone at home - q8011

	Completely safe	SE	Very safe	SE	Moderately safe	SE	Slightly safe	SE	Not safe at all	SE	Missing	SE	N
Gender													
Male	17.5	1.3	23.4	1.5	16.7	1.3	12.2	1.0	19.6	1.3	10.6	1.4	1,993
Female	12.5	0.9	21.9	1.3	17.2	1.0	13.4	1.0	26.6	1.2	8.4	0.9	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	12.2	1.0	19.1	1.2	23.0	1.7	17.4	1.2	19.6	1.4	8.8	1.0	1,397
Rural	16.2	1.1	24.3	1.3	14.0	0.9	10.6	0.9	25.1	1.2	9.8	1.1	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	21.5	2.4	25.7	2.4	13.6	1.5	11.2	1.7	21.7	1.7	6.2	1.2	820
Q2	14.4	1.9	25.3	2.1	12.6	1.5	11.5	1.5	29.5	2.4	6.7	1.4	842
Q3	12.8	1.4	24.0	2.2	14.3	1.7	11.8	1.6	26.1	2.2	11.0	1.8	937
Q4	13.7	1.6	17.0	1.6	20.7	2.5	14.8	1.8	20.9	2.2	12.8	2.3	698
Q5(richest)	11.1	1.3	19.3	2.1	26.2	2.2	16.3	1.7	17.6	2.2	9.5	1.6	725
M	17.2	3.4	23.1	4.2	15.4	3.4	11.0	2.6	18.8	3.6	14.5	3.1	219
Age													
18-29	14.3	1.2	22.1	1.4	15.7	1.3	12.9	1.1	23.4	1.4	11.5	1.3	1,748
30-44	15.9	1.4	20.2	1.4	17.5	1.3	13.8	1.3	23.0	1.4	9.7	1.3	1,292
45-59	15.4	1.7	24.8	2.4	20.3	2.2	12.2	1.9	19.6	2.0	7.6	1.3	696
60-69	14.3	2.6	30.1	3.7	14.8	2.4	7.4	1.8	28.0	3.6	5.5	1.8	293
70-79	11.5	3.2	23.4	4.3	19.1	4.0	17.7	4.2	26.8	4.0	1.6	0.9	160
80+	17.2	5.3	22.7	6.3	9.2	3.7	7.1	2.8	38.8	5.8	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	14.9	0.8	22.6	1.0	17.0	0.8	12.8	0.8	23.3	0.9	9.4	0.8	4,239

10.5 Feeling safe when walking down the street alone after dark - q8012

	Completely safe	SE	Very safe	SE	Moderately safe	SE	Slightly safe	SE	Not safe at all	SE	Missing	SE	N
Gender													
Male	15.0	1.1	16.4	1.2	14.7	1.1	12.0	1.0	29.7	1.5	12.2	1.4	1,993
Female	10.3	0.9	16.4	1.1	15.3	1.0	10.7	0.7	37.2	1.3	10.2	1.0	2,246
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Residence													
Urban	6.9	0.9	12.7	1.0	16.8	1.3	17.4	1.1	36.3	1.7	9.9	1.1	1,397
Rural	15.2	1.1	18.2	1.2	14.1	1.0	8.3	0.7	32.3	1.3	11.8	1.2	2,842
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Income													
Q1(poorest)	19.6	2.0	19.2	1.6	14.0	1.8	6.9	1.1	32.1	2.2	8.1	1.3	820
Q2	14.3	2.1	20.2	1.9	13.6	1.4	9.3	1.2	33.7	2.5	9.0	1.5	842
Q3	11.5	1.5	15.8	1.9	14.2	2.0	10.3	1.4	35.0	2.3	13.2	1.9	937
Q4	8.2	1.2	13.6	1.8	15.0	1.8	12.9	1.4	36.2	2.5	14.2	2.4	698
Q5(richest)	7.0	1.3	12.2	1.5	19.6	1.8	18.3	1.6	32.8	2.4	10.1	1.6	725
M	14.7	3.0	16.2	3.1	13.2	2.8	11.9	2.6	28.2	4.2	15.8	3.3	219
Age													
18-29	11.7	1.4	15.3	1.2	15.6	1.2	11.1	1.0	33.9	1.6	12.4	1.3	1,748
30-44	13.7	1.3	15.4	1.4	13.3	1.0	12.0	1.0	34.6	1.8	11.0	1.4	1,292
45-59	12.2	1.6	17.0	1.9	19.2	2.4	12.2	1.7	28.5	2.3	10.9	1.6	696
60-69	11.8	2.6	25.2	3.1	10.3	2.0	10.6	2.2	32.9	3.7	9.3	2.7	293
70-79	13.0	3.5	16.9	4.0	13.5	3.2	6.7	1.7	45.4	5.8	4.6	2.3	160
80+	14.8	5.2	21.8	5.4	15.1	4.4	4.5	2.0	38.7	6.7	5.0	2.5	49
M	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
All	12.5	0.8	16.4	0.8	15.0	0.8	11.3	0.6	33.6	1.0	11.1	0.9	4,239